

THE CLEAVAGE OF (PERHALOARYL)DIMETHYLSILANES BY ORGANOLITHIUM COMPOUNDS


MILTON R. SMITH JR. and HENRY GILMAN

Department of Chemistry, Iowa State University, Ames, Iowa 50010 (U.S.A.)

(Received October 12th, 1971)

SUMMARY

(Perhaloaryl)dimethylsilanes undergo a cleavage reaction with organolithium compounds, using very moderate conditions, to produce (perhaloaryl)lithium derivatives in good yield.




The corresponding trimethylsilyl compounds [except for 2,5-bis(trimethylsilyl)dichlorothiophene] and phenyldimethylsilane were not cleaved significantly.


INTRODUCTION

We observed, some years ago¹, that phenyllithium was formed in a cleavage reaction when *n*-butyllithium was allowed to react with triphenylsilanol or with hexaphenyldisiloxane. The cleavage of carbon-silicon bonds by an organolithium reagent has been shown to take place with a variety of cyclic silanes². Methyllithium effects some cleavage of a series of the $\text{Ph}_{4-n}\text{SiMe}_n$ type (but not with PhSiMe_3) to give $\text{PhLi}^{3,*}$.

More pertinent to our present studies was the cleavage of the pentachlorophenyl-silicon bond by methyllithium (and by phenyllithium), as in the following illustration⁴:



Then we observed⁵ the cleavage of the C_6F_5 -Si linkage by MeLi (and by BuLi) to yield $\text{C}_6\text{F}_5\text{Li}$:



In all these four cases, the (perhalophenyl)lithium was characterized by reaction with ClSiMe_3 to yield $\text{C}_6\text{X}_5\text{SiMe}_3$.

It was a short step to using ClSiMe_2H in place of the less accessible ClSiPh_2H to give $\text{C}_6\text{X}_5\text{SiMe}_2\text{H}$ types. These, as well as other (perhaloaryl)dimethylsilanes, were

* The compounds were unreactive to filtered MeMgI , but were later shown to be cleaved to a small extent by Mg/RMgX combinations^{3b}.

TABLE 1

REACTION OF ARYL- AND (POLYHALOARYL)DIMETHYL- AND -TRIMETHYLSILANES WITH ALKYL METALLIC REAGENTS IN ETHER

Silane (Mole)	RM (Mole)	Max. (°C)	Time	Deriv- atizing reagent	Product ^a and yield ^b
2-C ₄ Cl ₃ SSiMe ₂ H (0.025)	BuLi (0.025)	-70	10 min	CO ₂	2-C ₄ Cl ₂ SCOOH 5.2 g, 90%
2-C ₄ Cl ₃ SSiMe ₂ H (0.02)	BuMgBr	60 ^c	48 h		2-C ₄ Cl ₃ SSiMe ₂ H 4.05 g, 83%
2-(Me ₂ SiH)C ₄ Cl ₂ S- (Me ₂ SiH)-5 (0.02)	BuLi (0.02)	-70	2 h ^d	CO ₂	2-C ₄ Cl ₂ HSCOOH 3.0 g, 72%
2-(Me ₂ SiH)C ₄ Cl ₂ S- (Me ₂ SiH)-5 (0.02)	BuLi (0.04)	0	4.5 h	CO ₂	2-(COOH)C ₄ Cl ₂ S(COOH)-5 4.2 g, 87%
2-(Me ₂ SiH)C ₄ Cl ₂ S- (Me ₂ SiH)-5 (0.02)	PhLi (0.04)	0	6 h	Me ₃ SiCl	2-(Me ₃ Si)C ₄ Cl ₂ S(SiMe ₃)-5 45% ^e
2-(Me ₂ SiH)C ₄ Cl ₂ S- (Me ₂ SiH)-5 (0.02)	EtMgBr (0.04)	30	80 h		2-(Me ₂ SiH)C ₄ Cl ₂ S- (Me ₂ SiH)-5 ^f
C ₆ F ₅ SiMe ₂ H (0.02)	BuLi (0.02)	-70	1 h	CO ₂	C ₆ F ₅ COOH 3.3 g, 80%
4-C ₅ Cl ₄ NSiMe ₂ H (0.02)	BuLi (0.02)	-70	40 min	CO ₂	4-C ₅ Cl ₄ NCOOH 5.2 g, 86% ^g
C ₆ Cl ₅ SiMe ₂ H (0.02)	BuLi (0.02)	-15	3.5 h	CO ₂	Mixtures of tars and some acid products
C ₆ Cl ₅ SiMe ₂ H (0.02)	BuLi (0.04)	25	2 h	Me ₂ HSiCl	Mixture of tars. IR shows only weak Si-H stretch.
C ₆ Cl ₅ SiMe ₂ H (0.02)	BuLi (0.02)	0	5 h	Me ₃ SiCl	Mixture of C ₆ Cl ₅ SiMe ₃ 62% <i>p</i> -Me ₃ SiC ₆ Cl ₄ SiMe ₂ H- <i>p</i> 27% 5.8 g ^h
C ₆ Cl ₅ SiMe ₂ H (0.02)	MeLi (0.02)	-70	2 h	Me ₃ SiCl	C ₆ Cl ₅ SiMe ₃ 4.9 g, 76%
C ₆ H ₅ SiMe ₂ H (0.04)	BuLi (0.04)	10	16 h		C ₆ H ₅ SiMe ₂ Bu 6.6 g, 86% ⁱ
C ₆ H ₅ SiMe ₂ Bu (0.02)	BuLi (0.02)	30	24 h		No reaction
2-C ₄ Cl ₃ SSiMe ₃ (0.025)	BuLi (0.025)	-78	1 min	CO ₂	2-(SiMe ₃)C ₄ Cl ₂ S(COOH)-5 6.0 g, 91% ^j
2-C ₄ Cl ₃ SSiMe ₃ (0.02)	BuLi (0.04)	25	87 h	CO ₂	2-(SiMe ₃)C ₄ Cl ₂ S(COOH)-5 2.6 g, 48%
2-C ₄ Cl ₃ SSiMe ₃ (0.02)	BuMgBr (0.02)	60 ^c	20 h		2-C ₄ Cl ₃ SSiMe ₃ 4.7 g, 91%
2-(Me ₃ Si)C ₄ Cl ₂ S- (SiMe ₃)-5 (0.02)	BuLi (0.04)	10	4 h	H ₂ O	2,5-C ₄ Cl ₂ H ₃ S 5% ^k 2-HC ₄ Cl ₂ SSiMe ₃ -5 2%
2-(Me ₃ Si)C ₄ Cl ₂ S- (SiMe ₃)-5 (0.02)	BuLi (0.04)	-70	48 h	H ₂ O	2,5-C ₄ Cl ₂ H ₃ S 15% ^{c,h} 2-HC ₄ Cl ₂ SSiMe ₃ -5 8%

(For Footnotes see p. 37)

cleaved very smoothly and in high yields by BuLi. A similar cleavage has been reported recently⁶ by Pinkerton and Thames of the C-SiMe₃ linkage, particularly in some non-halogenated heterocycles.

RESULTS AND DISCUSSION

The cleavage of the carbon-silicon bond in the (perhaloaryl)dimethylsilanes proceeded smoothly and cleanly under very moderate conditions (-70°), and in relatively short reaction times (see Table 1). The products were [except for (pentachlorophenyl)dimethylsilane] the (polyhaloaryl)lithium compound and the alkyl- (usually *n*-butyl-) dimethylsilane or phenyldimethylsilane. All reactions proceeded with little or no apparent decomposition. The resulting perhaloaryllithium reagents were colorless.

This appears to be the first preparation of isomer-free tetrachloro-4-pyridyl-lithium in ether. Previous studies⁷ have produced mixtures of 3- and 4-lithio derivatives in low yields when ether was employed as the solvent. Furthermore, an extremely clean colorless solution is produced in contrast to the formerly observed "clean but colored" solutions.

There seemed to be some differences in the ease of cleavage of the bonds in the several compounds. Of the (perhaloaryl)dimethylsilanes, (trichloro-2-thienyl)dimethylsilane appeared to be cleaved most easily (less than 10 min). Similarly, the first dimethylhydrosilyl group was cleaved from 2,5-bis(dimethylsilyl)dichlorothiophene in less than 20 min. The second of these did not react until the temperature reached about 0° , at which time the reaction proceeded smoothly. The first of these cleavages takes place under conditions which fail to effect Cl-Li exchange in tetrachlorothiophene⁸, whereas the second -SiMe₂H group appears to be cleaved under essentially the same conditions required for the preparation of 2,5-dilithiodichlorothiophene⁹. The trimethylsilyl group can be cleaved from 2,5-bis(trimethylsilyl)dichlorothiophene, but not so readily nor in such good yields as is the SiMe₂H moiety. With trimethylsilyl derivatives examined, the reaction takes an alternate path^{5,10} (Table 1).

In the case of (pentachlorophenyl)dimethylsilane, a competing metal-halogen exchange reaction at the *p*-carbon atom was observed. When the reaction was re-run using methyllithium, which is known to be a poor reagent for the halogen-metal exchange reaction¹¹, the reaction proceeded "normally" to give (pentachlorophenyl)-lithium in good yield.

^a Products were identified by comparison with authentic samples (mixed m.p. and IR spectra).

^b Based on mole of starting silane.

^c Solvent was THF.

^d Reaction was complete in 20 min (VPC). The second SiMe₂H was cleaved by hydrolysis during work-up.

^e The yield was determined by VPC. Also present was 35% of 2-(Me₃Si)C₄Cl₂S(SiMe₂H)-5. From the mixture was isolated 1.12 g (21%) of PhSiMe₂H.


^f Product had other materials present. These were of longer retention time in the VPC and may have been the ethylated silanes.

^g No evidence of other isomers.

^h Percentages determined by VPC.


ⁱ n_D^{20} 1.4920; cited¹⁴ n_D^{20} 1.4920. Identical retention time with material prepared on a small scale from PhMe₂SiCl and BuLi.

^j Analysis found: Cl 26.40%. Calcd. for C₈Cl₂H₁₀O₂SSi: Cl 26.34%.



Even with the easily cleaved (trichloro-2-thienyl)dimethylsilane, no significant cleavage by a Grignard reagent was observed. In the case of ethylmagnesium bromide a product (ca. 05%) of longer retention time in the VPC may have been (trichloro-2-thienyl)ethyldimethylsilane. This is consistent with the lesser reactivity of most Grignard reagents, and also with known reactions of Grignard reagents and the Si-H bond^{12,13}.

The mechanism of the cleavage reaction can be visualized as a nucleophilic attack on the silicon atom, *viz.*:



Such a mechanism would be favored by electron-withdrawing groups (Cl, F) on the aryl nucleus which would increase the positive nature of the silicon atom. The absence of a cleavage reaction with PhMe₂SiH may be attributed largely to a lack of the strongly electron-withdrawing halogens. The reaction proceeded instead to give the known PhBuMe₂Si in 86% yield¹⁴. The fact that C₆Cl₅SiMe₂H is only partially cleaved by BuLi is probably due largely to the ease of the halogen-metal exchange reaction with the *p*-chlorine atom. We were unable to cleave a trimethylsilyl group from the trichlorothiophene nucleus. Instead there was a rapid (1 min) chlorine-lithium exchange reaction to give 2-lithio-5-(trimethylsilyl)dichlorothiophene in 91% yield at -78°.



The difference in behavior of the trimethylsilyl *vs.* the dimethylhydrosilyl types may be steric in origin. The use of more severe conditions might result in cleavage of (2-C₄Cl₃S)SiMe₃ by a second mole of alkyl lithium reagent. However, the first mole of alkyl lithium would still probably follow the reaction path of least resistance, *i.e.* the halogen-metal exchange on the α -position. The above exchange reaction did not take place when BuMgBr was used. It has been previously reported that 2-thienyltriphenylsilane in a similar reaction underwent metalation at the α -position of the thiophene nucleus¹.

We will report later on applications of this conversion or cleavage reaction for syntheses leading to R_FM and M(CF₂)_nM types.

EXPERIMENTAL

All reactions were effected under an atmosphere of dry nitrogen in a three-necked round-bottomed flask equipped with a nitrogen inlet tube, dropping funnel

and sealed stirrer. The equipment was dried at 120°, assembled while hot and flushed with dry nitrogen prior to charging the flask with the reactants. n-Butyllithium (1.6 N in hexane) phenyllithium and methyllithium (1.7 N in ether/benzene) were from the Foote Mineral Co.. Ether was distilled from sodium benzophenone ketyl. All other solvents were used without further purification. (Pentachlorophenyl)-¹⁵, (tetrachloro-4-pyridyl)-¹⁶, (pentafluorophenyl)-¹⁷, phenyl-¹⁸, (trichloro-2-thienyl)dime-thylsilanes⁸, 2,5-bis(dimethylsilyl)dichlorothiophene⁹, 2,5-bis(trimethylsilyl)dichloro-thiophene⁹, and (trichloro-2-thienyl)trimethylsilane⁸ were prepared by standard procedures.

General procedure

The reaction flask was charged with ca. 80 ml of dry ether and the appropriate silane and cooled to ca. -70° with a Dry Ice/acetone bath. To this solution, with vigorous stirring, was added an equivalent quantity of the organolithium compound. The reactions were monitored by inspection of VPC curves (4½ ft. Silicone Gum Rubber, 15% on Chromosorb W) of small aliquots after either hydrolysis and/or treatment with Me₃SiCl. When the particular reaction appeared complete, the products were derivatized. Details of the various reactions are contained in Table 1.

ACKNOWLEDGEMENT

This research was supported by the United States Air Force under contract F33615-70-C-1046 monitored by the Air Force Materials Laboratory, Wright-Pat-terson Air Force Base, Ohio. We are grateful to Dr. K. Leyshon and Dr. David H. Ballard for assistance with an unpublished reference.

REFERENCES

- 1 H. Gilman, R. A. Benkeser and G. E. Dunn, *J. Amer. Chem. Soc.*, 72 (1950) 1689.
- 2 H. Gilman and R. D. Gorsich, *J. Amer. Chem. Soc.*, 80 (1958) 1883, 3243; D. Wittenberg and H. Gilman, *J. Amer. Chem. Soc.*, 80 (1958) 2677; H. Gilman and W. H. Atwell, *J. Amer. Chem. Soc.*, 86 (1964) 5589; 87 (1965) 2678; K. Leyshon, Ph.D. thesis, Univ. of Leicester, 1962; K. Leyshon, private communication; see also, C. Eaborn and R. W. Bott, in A. G. MacDiarmid (Ed), *Organometallic compounds of the Group IV elements*. Vol. 1, Part I, Marcel Dekker, New York, 1968.
- 3 (a) H. Gilman, *Bull. Soc. Chim. Fr.*, (1963) 1356; (b) J. Grobe and V. Möller, *Chem. Ber.*, 103 (1970) 1302.
- 4 P. Morris, F. W. G. Fearon and H. Gilman, *J. Organometal. Chem.*, 9 (1967) 427.
- 5 F. W. G. Fearon and H. Gilman, *J. Organometal. Chem.*, 10 (1967) 409, 535.
- 6 F. H. Pinkerton and S. F. Thames, *J. Organometal. Chem.*, 29 (1971) C4; S. F. Thames, private communication.
- 7 J. D. Cook and B. J. Wakefield, *J. Organometal. Chem.*, 13 (1968) 15.
- 8 M. R. Smith Jr. and H. Gilman, *Organometal. Chem. Syn.*, 1 (1971) 265.
- 9 M. R. Smith Jr. and H. Gilman, *J. Organometal. Chem.*,
- 10 K. Shiina, T. Brennan and H. Gilman, *J. Organometal. Chem.*, 11 (1968) 471; I. Haiduc and H. Gilman, *J. Organometal. Chem.*, 14 (1968) 79.
- 11 H. Gilman and F. W. Moore, *J. Amer. Chem. Soc.*, 62 (1940) 1843; R. G. Jones and H. Gilman, *Organic Reactions*, 6 (1951) 339.
- 12 H. Gilman and E. A. Zuech, *J. Amer. Chem. Soc.*, 79 (1957) 4560; 81 (1959) 5925.
- 13 H. Gilman and S. P. Massie, *J. Amer. Chem. Soc.*, 68 (1946) 1128; R. N. Meals, *J. Amer. Chem. Soc.*, 68 (1946) 1880.
- J. Organometal. Chem.*, 37 (1972)

- 14 V. D. Tyurin, N. V. Ushakov, S. P. Gubin and N. S. Nametkin, *Izv. Akad. Nauk SSSR, Ser. Khim.*, (1968) 1407; *Chem. Abstr.*, 69 (1968) 77424y.
- 15 T. Brennan and H. Gilman, unpublished studies.
- 16 S. S. Dua and H. Gilman, *J. Organometal. Chem.*, 12 (1968) 234.
- 17 T. Brennan and H. Gilman, *J. Organometal. Chem.*, 16 (1969) 63.
- 18 R. A. Benkeser and D. J. Foster, *J. Amer. Chem. Soc.*, 74 (1952) 5314.

J. Organometal. Chem., 37 (1972)