


Bis-adamantylidenemethane

By J. STRATING, A. H. ALBERTS, and HANS WYNBERG*

(Department of Organic Chemistry, The University, Bloemsingel 10, Groningen, The Netherlands)

Summary The β -lactone dimer (II), derived from the keten 2-carboxyladamantane (I), gave the allene (III) on pyrolysis.

KETENS embodying the adamantane skeleton were recently prepared in our laboratory.¹ One of the ketens, (I), dimerized readily to the cyclobutane-1,3-dione derivative (Ia). We now report the direct dimerization of the keten (I) to the β -lactone (II) and the transformation of the latter into the allene (III).^{2,3}



The dimerization was effected by adding a catalytic amount of aluminium chloride to a refluxing dilute ethereal solution of 2-carboxyladamantane (I) and concentrating it by distilling the solvent slowly until a concentration of about 1 g of substance in 3 ml of ether was reached. Treating the mixture with water, and extraction with chloroform, furnished yellow solid material. Crystallizations from ethanol and octane yielded the β -lactone dimer [50%; m.p. $195\text{--}196^\circ$; i.r. (KBr): 1861 and 1720 cm^{-1} ; M.W. (mass spectr.) 324 ; analysis correct for $\text{C}_{22}\text{H}_{28}\text{O}_2$].

On pyrolysis, this compound easily lost CO_2 and a solid, the bisadamantylidenemethane (III), was produced in yields up to 95% as colourless needles, m.p. $224\text{--}226^\circ$, from acetone. The structure of (III) has been assigned on the following evidence (in addition to a correct elemental analysis): the i.r. spectrum shows a weak band at 1970 cm^{-1} due to the allenic linkage, and the absence of olefinic absorption. The n.m.r. spectrum consists of two peaks $\tau\ 7.5\text{--}7.7$, a singlet of 4 protons adjacent to the allene system, and $\tau\ 7.9\text{--}8.2$, a multiplet of 24 adamantyl protons. The mass spectrum exhibits a parent peak at $m/e\ 280$.

In view of the unique properties of a double bond between the adamantane skeletons⁴ a study of the reactivity of this allene is of considerable interest.

(Received, May 4th, 1970; Com. 662.)

¹ J. Strating, J. Scharp, and H. Wynberg, *Rec. Trav. chim.*, 1970, **89**, 23.

² Houben-Weyl, *Methoden der Organische Chemie*, Band VII/4, p. 276.

³ R. N. Lacey, *Adv. Org. Chem.*, 1960, **2**, 213; D. G. Farnum, I. R. Johnson, R. E. Hess, T. B. Marshall, and B. Webster, *J. Amer. Chem. Soc.*, 1965, **87**, 5191. The suggestion by Farnum and his co-workers concerning the mechanism of the β -lactone formation appears eminently sound in view of the large steric requirements of the adamantane group.

⁴ J. Strating, J. H. Wieringa, and H. Wynberg, *Chem. Comm.*, 1969, 907.