

Tetrahedron Vol. 51, No. 36, pp. 10033-10040, 1995 Copyright © 1995 Elsevier Science Ltd Printed in Great Britain. All rights reserved 0040-4020/95 \$9.50+0.00

0040-4020(95)00575-7

High-Pressure Diels-Alder Reactions of Bicyclic Dienones. Short Syntheses of Hydrophenanthrenones

Lucio Minuti*a, Aldo Taticchi*a, Eszter Gacs-Baitz^b, Ernest Wenkert^c

^aDipartimento di Chimica, Università di Perugia,via Elce di Sotto 8, 06100 Perugia, Italy; ^bCentral Research Institute of Chemistry, Hungarian Academy of Sciences, Budapest, Hungary; ^cDepartment of Chemistry (0506). University of California-San Diego, La Jolla, California, USA.

Abstract. Diels-Alder reactions of a heteroannular bicyclic dienone with acyclic dienes under Lewis acid catalysis at atmospheric and 7 Kbar pressure are presented. The pressure reactions are fast, high-yielding and highly diastereoselective en route to hydrophenanthrenones. As a consequence of some post-cycloaddition enolization, both cis and trans products are available.

The stereocontrolled construction of six-membered, polycarbocyclic compounds has been in the limelight for a long time chiefly due to the favorable, medicinal properties of many naturally occurring and unnatural representatives of this class of substances. In an attempt to develop short routes to synthesize these materials, based mainly on Diels-Alder reaction chemistry, recent studies were performed on the preparation of hydrophenanthrenones by cycloaddition of bicyclic, heteroannular dienones.^{1,2} Whereas the latter were poor dienophiles³, they afforded reasonable product yields on reaction with electron-rich dienes and on Lewis acid catalysis. In view of the known, strong acceleration of cycloadditions under high pressure⁴ and the recent experience of the promotion of a Diels-Alder reaction of β -methyl- α , β -unsaturated cycloalkenones (dienophiles totally unreactive at atmospheric pressure) under Lewis acid catalysis and high pressure⁵, an investigation of the pressurized and atmospheric cycloadditions of unsubstituted, bicyclic dienone **1a** with dienes **2a**, **b** and **c** was undertaken.



The dienone $(1a)^6$ needed for the study was prepared in the following manner. Treatment of $\Delta^{1,9}$ -octal-2-one⁷ with acetic anhydride and acetyl chloride in pyridine afforded its enol acetate, whose halogenation

with N-bromosuccinimide in dimethylformamide-water furnished a bromoketone. Exposure of the latter to a lithium bromide-carbonate mixture in the same medium produced dienone **1a** in a 75 % overall yield.⁸

RESULTS AND DISCUSSION

The cycloadditions were carried out in dry methylene chloride solutions of dienone **1a** under atmospheric or elevated (7 Kbar) pressure in the presence of ethylaluminum dichloride catalyst. The reaction with 2,3-dimethyl-1,3-butadiene (**2a**) at 7 Kbar led to a 9:1 mixture (70 % yield) of *anti-cis* and *syn-trans* tricyclic ketones, **3a** and **5a**, respectively, whereas at atmospheric pressure, the reaction produced the same product mixture (55 %) but in 1:9 ratio, respectively, a clear indication of the marked influence of pressure on the composition of the reaction mixture. When dienone **1a** was allowed to interact with (E)-piperylene (**2b**) at 7 Kbar, a 19:1 mixture (80 %) of *anti-endo-cis* and *syn-endo-trans* ketones **3b** and **5b**, respectively, was obtained. When the reaction was executed at atmospheric pressure, a 1:3 mixture (30 %) of the same ketones, respectively, was observed. As expected⁹, both anti and syn additions were regioselective and *endodiastereoselective*. Finally, the cycloaddition with 1,3-butadiene (**2c**) at 7 Kbar failed because of predominant diene polymerization, but at atmospheric pressure an 8.1 mixture (50 %) of *anti-cis* and *syn-cis* ketones, **3c** and **4c**, respectively, was produced. In this case, surprisingly, no *trans*-product was observed.



a, R = Me, R' = H; **b**, R = H, R' = Me; **c**, R = R' = H

The formation of unusual trans adducts (5a and b) in two of the three cycloadditions prompted a more detailed study of the Diels-Alder reaction between dienone 1a and diene 2a. Careful glc monitoring of the reaction at atmospheric pressure showed adduct 3a to be an early product and the 3a/5a product ratio to change throughout the course of the reaction. Furthermore, exposure of pure adduct 3a to the 1a-2a reaction conditions yielded the identical product mixture observed in the initial Diels-Alder reaction.¹⁰ These results indicate that both the Diels-Alder reactions of 2,3-dimethyl-1,3-butadiene (2a) and (E)-piperylene (2b)¹⁰ under atmospheric pressure are thermodynamically controlled and suggest that the formation of *trans*-compounds 5 is probably due to i) the conversion of the *anti-endo-cis* cycloadducts 3 into the short-lived *syn-endo-cis* adducts 4, and ii) the tautomerization (via a dienol or its aluminum complexes) of the elusive *sin-cis* cycloadducts 4 (cf. equations 1).

It seems reasonable that this explanation may be extended to the cycloadditions performed under high pressure conditions, with the different positions of equilibria depending on the effects of pressure⁴.



In principle, the *trans*-products 5 could also be ascribed to non-concertedness of the cycloaddition; this hypothesis, however, seems to be untenable in view of i) the observed *cis*-bridgehead products obtained in the cycloaddition of methylated dienone $1b^2$ where the angular methyl group precluded any post-cycloaddition tautomerization, and ii) the vast array of data on thermal and Lewis acid-catalyzed Diels-Alder reactions of conjugated cycloalkenones, none of which required interpretation in terms of a two-step mechanism^{3,5,11}.

Whereas the formation of *trans*-adduct in the Diels-Alder reactions of dienes 2a and 2b has been reasonably justified, the different behaviour of 1,3-butadiene (2c) is still not understood.

Finally the observed marked effect of pressure on the Diels-Alder reactions of hexalone 1a induced us to re-examine at 7 Kbar the cycloadditions of hexalones 1b and 1c with dienes 2a and 2b, previously studied at atmospheric pressure^{1,2}.

The **1b-2b** cycloaddition (70% yield) exhibited a 1.5 ± 1 *anti-endo/syn-exo* product ratio (vs. at atmospheric pressure²: 45%; 1 \pm 1 ratio); the **1c-2b** reaction (80%) gave only the *anti-endo* product (vs. 62%; 9 ± 1 ratio²) and the **1c-2a** Diels-Alder process (80%) afforded exclusively the *anti-endo* product (vs. 35%; the same compound¹).

In conclusion, several interesting facts emerge from the present study. Diels-Alder reactions of hexalones 1 under pressure are fast, high-yielding, and highly diastereoselective. The cycloadditions of hexalones 1 with acyclic dienes (2) constitute facile, one-step syntheses of hydrophenanthrenones from readily available starting materials. Hence they have the potential for yielding ready access to natural products of especially the di- and tri-terpenic or steroid types. The formation of trans compounds (5) by acid-induced post-cycloaddition isomerization makes the method more flexible and thus more useful in organic synthesis.

Demonstration of the utility of this synthetic strategy will be reported in due course.

STRUCTURE ANALYSIS

The structure and stereochemistry of the tricyclic ketones were inferred from the analysis of their highfield ¹H and ¹³C NMR spectra recorded in CHCl₃. Some of the ¹H spectra were repeated in C_6D_6 and/or in CD₃COOD to avoid overlapping of some proton resonances. The full assignment of the proton and carbon spectra followed from the examination of ¹H-¹H and ¹H-¹³C connectivities (COSY, HETCOR and INEPT spectra). Stereochemical and regiochemical assignments were based on the values of the J_{H,H} coupling constants and selective ¹H-¹H NOE experiments. The pertinent data are collected in the Experimental Section.

The cis-stereochemical relationship of H(4a), H(8a) and H(10a) for ketones 3a, 3b and 3c followed from NOE effects observed on the resonances of H(8a) and H(10a) upon irradiation of the resonance attributed to the H(4a) proton. The ${}^{3}J_{4a,10a}$ value of 5 Hz for these ketones confirmed a *cis*-stereochemistry at the C(4a)-C(10a) ring junction; by contrast, the same vicinal coupling $({}^{3}J_{4a,10a})$ in ketones 5a and 5b assumed the value of 11 Hz, indicating a trans-stereochemistry at C(4a)-C(10a) ring junction for both compounds. Furthermore the cis relationship between hydrogens H(4a) and H(8a) in both ketones 5a and 5b was revealed by ¹H-¹H NOE experiments. Selective pre-irradiation of the resonance due to H(8a) resulted in signal enhancement of the resonance attributed to H(4a), while no NOE effect was observed on H(4a) and H(8a) by selective pre-irradiation of H(10a). This stereochemical assignment was further supported by the similarity of the chemical shifts of carbon C(4a) and C(8a) of ketones 3a and 3c with those of the same carbon atoms of compound 5a. The regiochemistry and stereochemistry of the methyl group for ketones 3b and 5b followed from selective NOE experiments, as well as from the ${}^{3}J_{4,4a}$ value of 4.5 and 9.5 Hz, respectively. Selective pre-irradiation of the H(4) resonance of 3b resulted in signal enhancement of the resonances attributed to H-3, H(4a), and H(10a), revealing spatial proximity of these hydrogens. In the case of ketone 5b (see figure), saturation of the methyl group resonance gave rise to signal enhancement of H(4a); in a similar manner, saturation of the H(10a) resonance resulted in signal enhancement of H(4) signal. This indicated the position of the C(4) methyl group and the stereochemistry depicted in the figure for ketone 5b.



Figure. Minimized energy conformations of products 4c and 5b; the arrows indicate observed n.O.e's.

A cis relationship between H(4a) and H(10a) for ketone 4c was assured by mutual dipolar contact observed between the two hydrogens, as well as their coupling value of 5.5 Hz. Furthermore, selective preirradiation of the H(8a) resonance gave no NOE effect on the signal attributed to H(4a) and H(10a), while strong enhancement occurred at the resonances attributed to axial protons on carbon atoms C(4), C(7) and C(10). This indicated a *trans*-relationship between H(10a), H(4a) and H(8a). Two further facts are in consonance with the stereochemistry depicted in the figure for ketone 4c: i) the NOE effect observed between H(5) and H(4a), showing that proton H(4a) is in the plane of the C(4b)-C(5) double bond, and ii) the coupling constants of the hydrogens at C(4a) (${}^{3}J_{4a,4\alpha} = 11.5$, ${}^{3}J_{4a,10a} = 5.5$, ${}^{3}J_{4a,4\beta} = 4.5$ Hz) and at

10036

C(8a) $({}^{3}J_{8a,8\beta} = 9, {}^{3}J_{8a,8\alpha} = 5, {}^{3}J_{8a,9\beta} = 13, {}^{3}J_{8a,9\alpha} = 6$ Hz) indicating a conformation as depicted in the figure [e.g. H(4a) assumes a *trans*-axial orientation with respect to H(4)].

EXPERIMENTAL SECTION12

4,4a,5,6-Tetrahydro-2(3H)-naphthalenone (1a). Acetyl chloride (7 mL) was added under nitrogen to a solution of $\Delta^{1,9}$ -octal-2-one⁷ and then refluxed in an atmosphere of nitrogen for 3 h. The reaction mixture was cooled, diluted with water and worked up as usual, to afford 6 g of enol acetate.

A solution of the enol acetate in dimethylformamide (32 mL) and water (0.8 mL) was treated with NBS (6 g) at 0°C and under nitrogen. The addition of NBS was complete within 0.5 h. The reaction mixture was stirred at 0°C for 0.5 h. Then LiBr (2.7 g) and Li_2CO_3 (5.4 g) were added and the reaction mixture was heated at 70°C for 2.5 h under stirring. It was cooled, poured into ice-water (150 mL) and acetic acid (10 mL) and extracted with ether. After the usual workup a residual oil was obtained, which was purified by column chromatography. Elution with 4:1 hexane-ether gave 2.6 g of pure dienone **1a** in 75% overall yield.

Colorless oil; IR 1653 (s, C=O), 1617 (m, C=C) cm⁻¹; ¹H NMR δ 1.20-2.60 (m, 9H), 5.68 (s, 1H, H-1), 6.20 (m, 2H, H-7, H-8); ¹³C NMR δ 26.11 (C-6), 29.14 (C-4 or C-5), 30.12 (C-5 or C4), 35.43 (C-4a), 37.81 (C-3), 123.76 (C-1), 128.27 (C-8), 139.21(C-7), 158.21 (C-8a), 199.96 (C-2).

Diels-Alder Reactions of Dienone 1a. General Procedure.

The mixture of the starting materials was prepared in a dry-box by adding the hexane solution of $EtAlCl_2$ to a dichloromethane solution of dienone. The mixture was kept at 25°C for 40 min.¹³ and then the diene was added. Thereafter (in the case of the Diels-Alder reaction being carried out at atmospheric pressure) the mixture was heated in an oil bath and then worked up by following a previously reported set procedure^{3a}. In the case of the Diels-Alder cycloadditions being performed under pressure, all operations were executed according to the recently described procedure⁵.

Diene	Diene/	Reaction	Pressure	Products	Product
	Ketoneo	Time (h)	<u>(K</u> bar)		<u>Yield (%)</u>
2a	3	22	7	9:1 3a/5a	70
2a	5	240	-	1:9 3a/5a	55d
2b	3	65	7	19:1 3b/5b	80
2b	3	90	-	1:3 3b/5b	30d
2c	10	90	-	8:1 3c/4c	50d

Table I. Conditions of the Diels-Alder Reactions of Hexalone 1a with Dienes 2a

^aComplexation time¹³: 40 min; complexation temperature¹³: 25° C; reaction temperature: 35° C; ketone concentration: 0.2 M; catalyst/ketone equiv. ratio: 0.5. ^bRatio of equivs. ^cBased on isolated compounds. ^dPlus product(s) (up to 3 %) of unknown constitution.

1a-2a Reactions. Tricyclic ketone **3a** was isolated from the reaction mixture of the cycloaddition performed under pressure (elution with 7:3 petroleum ether-ether). Tricyclic ketone **5a** was isolated from the reaction mixture of the cycloaddition accomplished at atmospheric pressure (same elution).

2,3-Dimethyl-1,4,4af,8,8af,9,10,10af-octahydro-6(7H)-phenanthrenone (3a): m.p. 67-68°C; IR 1658 (s, C=O), 1615 (w, C=C) cm⁻¹; ¹H NMR δ 1.54 (br.s, 3H, 3-Me), 1.64 (br.s, 3H, 2-Me), 1.65 (m, 3H, H-8, H-9, H-10), 1.75-1.85 (m, 4H, Hs-1, H-9, H-10), 2.06 (d, 1H, J = 18.0, 1.5 Hz, H-4), 2.14 (ddd, 1H, J = 13.5, 8.5, 5.5 Hz, H-8), 2.24 (m, 1H, J = 9.0, 7.0, 5.0, 4.5, 2.6 Hz, H-10a), 2.26 (m, 1H, H-4), 2.28 (m, 2H, J = 16.0, 11.0, 9.0, 5.0, 1.5 Hz, H-7, H-8a), 2.39 (ddd, 1H, J = 16.0, 8.5, 5.0 Hz, H-7), 2.53 (m, 1H, J = 5.0, 5.0, 1.5, 1.0 Hz, H-4a), 5.63 (dd, 1H, J = 1.5, 1.0 Hz, H-5); ¹³C NMR δ 18.58 (C-2 Me), 28.38 (C-8), 29.51 (C-9), 30.17 (C-1 or C-10), 32.20 (C-1 or C-10), 33.67 (C-4), 35.33 (C-7), 35.67 (C-10a), 38.34 (C-8a), 42.20 (C-4a), 122.15 (C-2 or C-3), 122.74 (C-5), 124.33 (C-2 or C-3), 168.02 (C-4b), 200.35 (C-6); MS, m/e (rel intensity) 230 (M⁺, 30), 157 (10), 150 (11), 149 (base), 120 (62), 119 (24). Anal. Calcd for C₁₆H₂₂O: C, 83.43; H, 9.63. Found: C, 83.49; H, 9.61.

2,3-Dimethyl-1,4,4aβ,8,8aβ,9,10,10aα-octahydro-6(7H)-phenanthrenone (5a): m.p. 100-101°C; IR 1662 (s, C=O), 1617 (w, C=C) cm⁻¹; ¹H NMR δ 1.24 (m, 1H, J = 13.0, 12.5, 11.0, 4.0 Hz, H-10), 1.30 (m, 1H, j = 13.5, 12.5, 12.0, 4.0 Hz, H-9), 1.43 (m, 1H, J = 11.5, 11.0, 11.0, 4.0, 4.0 Hz, H-10a), 1.62 (br.s, 3H, 3-Me), 1.66 (br.s, 3H, 2-Me), 1.67 (m, 1H, J = 13.5, 12.0, 8.5, 5.0 Hz, H-7), 1.83-1.86 (m, 2H, H-9, H-10), 1.96 (m, 1H, J = 11.0, 7.5, 7.5, 1.4 Hz, H-4a), 1.98 - 2.06 (m, 4H, Hs-1, Hs-4), 2.08 (ddd, 1H, J = 13.5, 5.5, 5.0 Hz, H-8), 2.27 (ddd, 1H, J = 16.0, 12.0, 5.0 Hz, H-7), 2.28 (m, 1H, J = 12.0, 8.5, 5.0, 5.0, 5.0, 1.5 Hz, H-8a), 2.37 (ddd, 1H, J = 16.0, 5.5, 5.0 Hz, H-7), 5.88 (br.s, 1H, H-5); ¹³C NMR δ 18.57 (C-3 Me), 19.04 (C-2 Me), 29.06 (C-8), 32.72 (C-10), 33.73 (C-9), 34.22 (C-4), 36.05 (C-7), 38.47 (C-8a), 39.78 (C-10a), 40.64 (C-1), 43.37 (C-4a), 122.10 (C-5), 124.17 (C-2), 124.61 (C-3), 169.00 (C-4b), 200.27 (C-6); MS, m/e (rel. intensity) 230 (M⁺, base), 202 (20), 187 (17), 159 (17), 157 (18), 132 (17). Anal. Calcd for C₁₆H₂₂O: C, 83.43; H, 9.63. Found: C, 83.54; H, 9.59.

la-2b Reactions. Tricyclic ketones **3b** and **5b** were purified by column chromatography of the crude mixtures of reactions performed at atmospheric and elevated pressure, eluting with gradient petroleum ether to 7:3 petroleum ether-ether.

4 α -Methyl-1,4,4 $a\beta$ 8,8 $a\beta$ 9,10,10 $a\beta$ -octahydro-6(7H)-phenanthrenone (3b): m.p. 77-78°C; IR 1660 (s, C=O), 1625 (m, C=C) cm⁻¹; ¹H NMR & 0.99 (d, 3H, J = 7.3 Hz, 4-Me), 1.15-1.44 (m, 5H, H-8, Hs-9, Hs-10), 1.52 (ddd, 1H, J = 18.0, 5.5, 2.5 Hz, H-1), 1.60 (m, 1H, H-8a), 1.67 (ddd, 1H, J = 18.0, 11.5, 4.5 Hz, H-1), 1.73 (m, 1H, H-8). 1.91 (m, 1H, J = 11.5, 5.0, 4.5, 4.5, 2.5 Hz, H-10a), 2.01 (ddd, 1H, J = 4.5, 4.5, 1.0 Hz, H-4a), 2.14 (ddd, 1H, J = 16.0, 5.0, 5.0 Hz, H-7), 2.19 (ddd, 1H, J = 16.0, 11.0, 5.0 Hz, H-7), 2.31 (m, 1H, J = 7.3, 4.5, 4.5, 3.0 Hz, H-4), 5.36 (m, 1H, J = 10.0, 4.5, 3.0 Hz, H-2), 5.39 (ddd, 1H, J = 10.0, 4.5, 2.5 Hz, H-3), 6.16 (br.s, 1H, H-5); ¹³C NMR & 17.23 (C-4 Me), 26.36 (C-1), 27.72 (C-8), 30.41 (C-9), 30.55 (C-10), 33.92 (C-4), 33.96 (C-7), 39.03 (C-10a), 39.89 (C-8a), 48.96 (C-4a), 124.65 (C-5), 125.55 (C-2), 130.54 (C-3), 161.86 (C-4b), 197.06 (C-6); MS, m/e (rel. intensity) 216 (M⁺, 39), 149 (base), 120 (61), 105 (29), 91 (79). Anal. Calcd for: C₁₅H₂₀O C, 83.29; H, 9.32. Found: C, 83.40; H, 9.32.

4β-Methyl-1,4,4aβ,8,8aβ,9,10,10aα-octahydro-6(7H)-phenanthrenone (5b): m.p. 102-103°C; IR 1662 (s, C=O), 1621 (m, C=C) cm⁻¹; ¹H NMR δ + 01 (d, 3H, J = 7.0 Hz, 4-Me), 1.34 (m, 1H, J = 13.5,

10038

12.0, 11.0, 4.5 Hz; H-10), 1.51 (m, 1H, J = 11.0, 11.0, 11.0, 4.5, 4.0 Hz, H-10a), 1.64 (m, 1H, J = 13.5, 12.5, 12.0, 4.0 Hz, H-9), 1.73 (ddd, 1H, J = 11.0, 9.5, 1.5 Hz, H-4a), 1.78, (m, 1H, H-8), 1.84 (ddd, 1H, J = 17.0, 11.0, 2.0 Hz, H-1) 1.90 (m, 2H, H-9, H-10), 2.13 (ddd, 1H, J = 17.0, 4.5, 2.0 Hz, H-1), 2.23 (m, 1H, J = 13.5, 12.0, 10.0, 5.0 Hz, H-8), 2.30 (m, 1H, J = 12.5, 10.0, 5.0, 5.0 Hz, H-8a), 2.34 (ddd, 1H, J = 16.0, 5.5, 5.0 Hz, H-7), 2.42 (ddd, 1H, J = 16.0, 12.0, 5.0 Hz, H-7), 2.44 (ddd, 1H, J = 9.5, 7.0, 2.5 Hz, H-4), 5.49 (m, 1H, J = 10.0, 2.5, 1.8, 1.5 Hz, H-3), 5.62 (m, 1H, J = 10.0, 4.5, 2.5, 2.0 Hz, H-2), 5.87 (dd, 1H, J = 1.5, 1.0 Hz, H-5); 13 C NMR δ 20.27 (C-4 Me), 27.87 (C-8), 29.97 (C-4), 33.30 (C-1), 33.54 (C-9), 33.66 (C-10), 34.24 (C-7), 39.87 (C-8a), 40.60 (C-10a), 52.17 (C-4a), 121.12 (C-5), 123.96 (C-2), 132.40 (C-3), 169.21 (C-4b), 200.04 (C-6); MS, m/e (rel. intensity) 216 (M⁺, base), 201 (24), 149 (38), 145 (21), 120 (32), 91 (44). Anal. Calcd for C₁₅H₂₀O⁻C, 83.29, H, 9.32. Found⁻C, 83.16; H, 9.30.

la-2c Reaction. The crude mixture was chromatographed on a column and elution with gradient petroleum ether to 7/3 petroleum ether-ether led to pure ketones **3c** and **4c**.

1,4,4aβ,8,aβ,9,10,10aβ-Octahydro-6(7H)-phenanthrenone (3c): m.p. 60-61°C; IR 1653 (s, C=O), 1613 (m, C=C) cm⁻¹; ¹H NMR δ 1.66 - 1.88 (m, 7H, Hs-1, H-8, Hs-9, Hs-10), 2.14 (m, 1H, J = 13.5 8.5, 5.2, 5.0 Hz, H-8), 2.21-2.27 (m, 2H, Hs-4), 2.28 (m, 1H, J = 9.0, 6.5, 5.0, 4.5, 3.0 Hz, H-10a), 2.29 (ddd, 1H, J = 16.0, 10.0, 4.5 Hz, H-7), 2.30 (m, 1H, J = 11.5, 11.0, 5.0, 4.0, 1.5 Hz, H-8a), 2.39 (ddd, 1H, J = 16.0, 8.5, 5.0 Hz, H-7), 2.58 (m, 1H, J = 5.5, 5.0, 1.4, 1.0 Hz, H-4a), 5.56 (m, 1H, J = 10.0, 4.0, 2.0, 2.0 Hz, H-2), 5.60 (m, 1H, J = 10.0, 4.0, 2.5, 1.5 Hz, H-3), 5.81 (dd, 1H, J = 1.5, 1.4 Hz, H-5); ¹³C NMR δ 25.76 (C-1), 27.19 (C-4), 28.44 (C-8), 29.83 (C-9), 30.26 (C-10), 35.08 (C-10a), 35.31 (C-7), 38.43 (C-8a), 41.51 (C-4a), 122.78 (C-5), 123.78 (C-3), 126.02 (C-2), 167.16 (C-4b), 200.06 (C-6); MS, m/e (rel. intensity) 202 (M⁺, 83), 148 (50), 120 (89), 91 (base), 79 (51); Anal. Calcd for C₁₄H₁₈O: C, 83.12; H, 8.97. Found: C, 83.21; H, 9.01.

1,4,4a α *8,8aβ*,9,10,10a α -Octahydro-6(7H)-phenanthrenone (4c): m.p. 80-81°C; IR 1662 (s, C=O), 1613 (m, C=C) cm⁻¹: ¹H NMR δ 1.24 (m, 1H, J = 13 5, 13.0, 4.0 Hz, H-9), 1.44 (m, 1H, H-10), 1.61 (m, 1H, J = 13.5, 13.0, 9.0, 4.0 Hz, H-8), 1.73 (m, 1H, J = 13.0, 12.5, 3.5 Hz, H-10), 1.86 (m, 1H, H-1), 1.92 (m, 1H, J = 12.5, 5.5, 3.5, 3.0, 3.0 Hz, H-10a), 1.95 (m, 1H, H-4), 2.01 (m, 1H, J = 13.5, 6.0, 3.5, 3.0 Hz, H-9), 2.10 (m, 1H, J = 13.5, 5.0, 4.5, 4.0 Hz, H-8), 2.18 (m, 1H, H-4), 2.30 (ddd, 1H, J = 16.5, 13.0, 4.5 Hz, H-7), 2.38 (m, 1H, H-1), 2.42 (ddd, 1H, J = 16.5, 4.0, 4.0 Hz, H-7), 2.54 (m, 1H, J = 13.0, 9.0, 6.0, 5.0, 2.0 Hz, H-8a), 2.64 (ddd, 1H, J = 11.5, 4.5, 4.5 Hz, H-4a), 5.63 (m, 1H, H-3), 5.64 (m, 1H, H-2), 5.87 (d, 1H, J = 2.0 Hz, H-5); ¹³C NMR δ 26.33 (C-10), 27.26 (C-4), 29.59 (C-8), 30.59 (C-1), 33.43 (C-8a), 34.31 (C-9), 34.72 (C-10a), 37.01 (C-7), 42.26 (C-4a), 124.01 (C-3), 124.69 (C-5), 125.04 (C-2), 170.62 (C-4b), 200.49 (C-6); MS, m/e (rel. intensity) 202 (M⁺, 81), 160 (19), 148 (62), 120 (base), 91 (42); Anal. Calcd for C₁₄H₁₈O: C, 83.12; H, 8.97. Found: C, 83.11; H, 8.93.

Acknowledgement: L.M. and A.T. thank the MURST and C.N.R. for financial support of the work in Perugia. The work in Budapest was supported by OTKA grant n T016715.

REFERENCES AND NOTES

- 1. Guo, M.; Minuti, L.; Taticchi, A.; Wenkert, E. J. Org. Chem. 1989, 54, 6138-6140.
- 2. Guo, M.; Minuti, L.; Selvaggi, R.; Taticchi, A.; Wenkert, E. Can. J. Chem. 1992, 70, 1481-1485.
- (a) Fringuelli, F.; Pizzo, F.; Taticchi, A.; Halls, T.D.J.; Wenkert, E.J. Org. Chem 1982, 47, 5056-5065. (b) Fringuelli, F.; Taticchi, A.; Wenkert, E. Org. Prep. Proced. Int. 1990, 22, 131-165.
- 4. Matzumoto, K.; Acheson, R.M. Organic Synthesis at High Pressure, John Wiley & Sons, Inc., New York, 1991
- (a) Aben, R.W. M.; Minuti, L.; Scheeren, H. W.; Taticchi, A. *Tetrahedron Lett.* 1991, 32, 6445-6448.
 (b) Minuti, L.; Scheeren, H. W.; Selvaggi, R.; Taticchi, A. *Synth. Commun.* 1992, 22, 2965-2969.
- (a) Birch, A. J.; Murray, A. R.; Smith, H. J. Chem. Soc. 1951, 1945-1950. (b) Dolbj, L. J.; Alder, E. Tetrahedron Lett. 1971, 3803-3806.
- Stork G.; Brizzolara, A.; Landesman, H.; Szmuszkovicz, J.; Terrel, R. J. Am. Chem. Soc. 1963, 85, 207-222
- 8. An alternate route to the dienone, dehydrogenation of $\Delta^{1,9}$ -octal-2-one with chloranil failed.
- 9. Fringuelli, F.; Taticchi, A. *Dienes in the Diels-Alder Reaction*, John Wiley & Sons Inc., New York, 1990.
- The same type of results emanated from the exposure of pure adduct 3b to the 1a-2b reaction conditions (under atmospheric pressure). The product mixture (3b and 5b) was the same as that from the original cycloaddition.
- Inter alia: (a) Angell, E. C.; Fringuelli, F.; Pizzo, F.; Porter, B.; Taticchi, A.; Wenkert, E. J. Org. Chem. 1986, 51, 2642-2649. (b) Fringuelli, F.; Guo, M.; Minuti, L.; Pizzo, F.; Taticchi, A.; Wenkert, E. J. Org. Chem. 1989, 54, 710-712. (c) Angell, E. C.; Fringuelli, F.; Pizzo, F.; Minuti, L.; Taticchi, A.; Wenkert, E. J. Org. Chem. 1989, 54, 1217-1218. (d) Guo, M.; Minuti, L., Taticchi, A.; Wenkert, E. J. Org. Chem. 1990, 55, 1366-1368. (e) Gacs-Baitz, E.; Matchytka, D.; Minuti, L.; Selvaggi, R.; Taticchi, A. Synth. Commun. 1991, 21, 2143-2155. (f) Fringuelli, F.; Minuti, L.; Pizzo, F.; Taticchi, A. Acta Chem. Scand. 1993, 47, 255-263.
- (a) For general experimental details see: Gacs-Baitz, E.; Minuti, L.; Taticchi, A., *Tetrahedron*, 1994, 50, 10359-10366. All products were crystallized from hexane. All NMR spectra were run at room temperature (internal Me₄Si) on a Varian Associates VXR-400 multinuclear instrument. (b) Based on the doctoral dissertation of Maravalli Federico.
- 13. Fringuelli, F.; Pizzo, F; Taticchi, A.; Wenkert, E. J. Org. Chem. 1983, 48, 2802-2808

(Received in UK 19 May 1995; revised 7 July 1995; accepted 14 July 1995)

10040