

SYNTHESIS AND HIGH-PRESSURE DIELS-ALDER CYCLOADDITIONS OF 6-METHOXYCARBONYL-3-OXO-2-AZABICYCLO[2.2.0]HEX-5-ENE

Hiroto NAKANO and Hiroshi HONGO*

Tohoku College of Pharmacy, 4-4-1 Komatsushima, Aoba-ku, Sendai 981, Japan







The efficient photochemical synthesis of photopyridone having an electron-withdrawing group and its high-pressure Diels-Alder cycloaddition, the first examples of an application of the technique relating to photopyridone, are reported.

KEYWORDS photoisomerization; photo-2(1*H*)-pyridone; high-pressure Diels-Alder cycloaddition; azapolycyclic ring system; β -lactam

The photoisomers, 3-oxo-2-azabicyclo[2.2.0]hex-5-enes, of 2(1*H*)-pyridones contain β -lactam and cyclobutane, and therefore have great potential as synthetic intermediates. Thus, photo-2(1*H*)-pyridones may become synthons of β -lactam antibiotics¹⁾ and a potent inhibitor of human immunodeficiency virus (HIV).²⁾ In contrast to the substantial amount of experimental work on photoisomerization of 2(1*H*)-pyridones having electron-donating groups in the ring,³⁾ little attention has been focused on similar reactions of 2(1*H*)-pyridones possessing electron-withdrawing substituents, except for the reactions of 3-cyano-4-methoxy-1-methyl-^{3a)} or monochloro-⁴⁾2(1*H*)-pyridones. Herein we report the photoisomerization of 1-unsubstituted 2(1*H*)-pyridones having a methoxycarbonyl substituent in the ring; the reaction of the 2(1*H*)-pyridone (**1a**) gave the photopyridone (**2a**) with no dimer formation in a good yield. The photopyridone (**2a**) obtained is potentially a valuable synthetic intermediate, and may act as a dienophile leading to richly functionalized, polycyclic ring systems fusing β -lactam (**4a-e**), since it contains an electrophilic group. These compounds (**2a** and **4a-e**) are expected to possess interesting chemical properties and pharmacological activities.

Irradiations of the 2(1*H*)-pyridones (**1a,b**) (10^{-2} M) in benzene with a 400-W high-pressure mercury lamp through a Pyrex vessel were carried out for 24h. The reaction of the 2(1*H*)-pyridone (**1a**) gave the stable photopyridone (**2a**)⁵⁾ [52%, mp 68-70°C, IR: 1775 cm^{-1} (β -lactam carbonyl)], which was isolated by column chromatographic separation of the concentrated residue, and the irradiation of 2(1*H*)-pyridone (**1b**) afforded the photodimer (80%, mp 184-186°C).

In general, photopyridones are all liquids or low melting solids, and revert readily to 2(1*H*)-pyridones under appropriate thermal conditions. Likewise, the photopyridone (**2a**) also gradually returned to the 2(1*H*)-pyridone (**1a**) in refluxing benzene. Although a high-pressure strategy has proven extremely useful to surmount the energy barrier imposed by the steric and electronic effects and to reduce the thermal influence in cycloaddition reaction, such as Diels-Alder reaction,⁶⁾ an application of the technique relating to photopyridones has not been reported. Diels-Alder cycloadditions of the photopyridone (**2a**) with the synthetically useful dienes (**3a-d**), (**3a**: buta-1,3-diene, **3b**: 2,3-dimethyl-buta-1,3-diene, **3c**: furan, **3d**: anthracene) (2 eq) were carried out under high-pressure conditions (10Kbar, 50°C, 24h, dichloromethane), and the *cis-anti-cis* adducts (**4a-d**) were stereoselectively obtained in excellent or good yields, respectively.

**1a,b****a:** $R^1 = H$, $R^2 = CO_2Me$ **b:** $R^1 = CO_2Me$, $R^2 = H$ **2a****4a,b****a:** $R = H$, 88%, mp 94-95 °C**b:** $R = Me$, 87%, mp 51-52°C**4c**, 73%, mp 170-171°C**4d**, 70%, mp 209-210°C**4e**, 75%, viscous oil

Furthermore, treatment of the photopyridone (2a) with 1-methoxy-3-(trimethylsilyloxy)buta-1,3-diene (Danishefsky's diene) (2 eq) under high-pressure conditions (6.5Kbar, 50°C, 24h, dichloromethane) followed by acid work-up (camphor-10-sulfonic acid) regio- and stereo-selectively gave the *cis-anti-cis* adduct (4e) in a good yield. The *cis-anti* stereochemistry (C-5 and C-6 positions) was confirmed by the observation of ¹H-NMR NOE enhancement between Ha and Hb, and the absence of any NOE between Ha and Hc. All new compounds (2a and 4a-e) were characterized by IR, ¹H-NMR spectroscopy, elemental analysis, mass and high-resolution mass spectrometry.

REFERENCES AND NOTES

- 1) a) E. J. Corey, J. Streith, *J. Am. Chem. Soc.*, **86**, 950 (1964); b) G. Kretschmer, R. N. Warrener, *Tetrahedron Lett.*, **16**, 1335 (1975); c) W. J. Begley, G. Lowe, A. K. Cheetham, J. M. Newsam, *J. Chem. Soc., Perkin Trans. I*, **1981**, 2620; d) J. Brennan, *J. Chem. Soc., Chem Commun.*, **1981**, 880; e) T. Kametani, T. Mochizuki, T. Honda, *Heterocycles*, **19**, 89 (1982); f) C. Kaneko, T. Naito, A. Saito, *Tetrahedron Lett.*, **25**, 159 (1984); g) N. Katagiri, M. Sato, N. Yoneda, S. Saikawa, T. Sakamoto, M. Muto, C. Kaneko, *J. Chem. Soc., Perkin Trans. I*, **1986**, 1289.
- 2) N. Katagiri, H. Sato, C. Kaneko, *Chem. Pharm. Bull.*, **38**, 288 (1990).
- 3) a) R. C. De Selms, W. R. Schleigh, *Tetrahedron Lett.*, **1972**, 3563; b) H. Furrer, *Chem. Ber.*, **105**, 2780 (1972); c) K. Shiba, C. Kaneko, *J. Chem. Soc., Chem Commun.*, **1980**, 537; d) J. Kurita, T. Yoneda, N. Kakusawa, T. Tsuchiya, *Chem. Pharm. Bull.*, **38**, 2911 (1990).
- 4) W. L. Dilling, N. B. Tefertiller, A. B. Mitchell, *Mol. Photochem.*, **5**(4), 371 (1973).
- 5) 2a: ¹H-NMR (400 MHz, CDCl₃) δ, 7.21 (1H, d, C₅-H), 6.45 (1H, br s, NH), 4.59 (1H, dd, C₁-H), 4.20 (1H, br s, C₄-H). The decoupling experiments gave *J*_{1,4} = 2.2Hz, *J*_{1,5} = 2.9Hz, and *J*_{4,5} = 0.7Hz.
- 6) a) K. Matsumoto, A. Sera, T. Uchida, *Synthesis*, **1985**, 1; b) K. Matsumoto, A. Sera, *ibid.*, **1985**, 999; c) W. J. Ie. Noble (ed.), "Organic High Pressure Chemistry," Elsevier, Amsterdam, 1988.

(Received July 14, 1993)