

Condensation of Phosphonate Anions with 4-Amino-5-nitrosopyrimidines: a New Pteridine Synthesis¹

By EDWARD C. TAYLOR* and BEN E. EVANS


(Department of Chemistry, Princeton University, Princeton, New Jersey 08540)

Summary The condensation of diethyl phosphonate carbanions, prepared from α -bromo-esters, -nitriles, or -ketones, with 4,6-diamino-5-nitrosopyrimidines constitutes a versatile and unequivocal route to pteridin-7(8H)-ones, 7-aminopteridines, and 7-alkyl- or arylpteridines.

A RECENT report² describing a novel method for the synthesis of pteridin-7(8H)-ones by the condensation of triethyl phosphonoacetate with several 2-substituted 4,6-diamino-5-nitrosopyrimidines prompts us to report our independent observation that the reaction has greater scope than previously indicated, and constitutes a versatile, unequivocal synthetic route to pteridin-7(8H)-ones, 7-aminopteridines, and 7-alkyl- or arylpteridines.

Addition of phosphonate carbanions to carbonyl groups to afford olefins is a widely employed modification of the Wittig reaction.³ The analogous addition of such anions to the nitroso-group has also been successful,⁴ and these latter reports prompted our interest in the possible condensation of phosphonate anions with 4-amino-5-nitrosopyrimidines as a general synthetic approach to pteridines.[†]

agreement with results already reported.² A variety of other phosphonate anions, readily prepared by reaction of α -bromo-esters, -nitriles or -ketones with triethyl phosphite followed by addition of base, reacted analogously to give pteridin-7(8H)-ones, 7-amino-, and 7-alkyl- or arylpteridines, respectively, carrying alkyl or aryl groups (or hydrogen) at C-6. Some representative 4-amino-2-phenylpteridines prepared from 4,6-diamino-5-nitroso-2-phenylpyrimidine by these procedures are listed in the Table.



TABLE

Pteridine	m.p. (° C)	Yield (%)
(I) R=H	331—332 (decomp) ^a	90
(I) R=Me	289—291 (decomp) ^b	74
(I) R=Me (H for Ph at C-2)	>350 ^c	78
(II) R=H	295—298 (decomp) ^d	28
(II) R=Ph	368 (decomp) ^e	27
(III) R ¹ =H, R ² =Ph	253.5—254 (decomp)	75
(III) R ¹ =R ² =-CH ₂ CH(Me)·[CH ₂] ₂ -	317 (decomp)	32

^a Lit. m.p. >340° (ref. 2).


^b Lit. m.p. 282—284°: J. Weinstock, R. Y. Dunoff, J. E. Carevic, J. G. Williams, and A. J. Villani, *J. Medicin. Chem.*, 1968, **11**, 618.

^c Lit. m.p. >350°: D. Söll and W. Pfeleiderer, *Chem. Ber.*, 1963, **96**, 2977.

^d Lit. m.p. 299—302°: J. Weinstock, R. Y. Dunoff, and J. G. Williams, *J. Medicin. Chem.*, 1968, **11**, 542.

^e Lit. m.p. >320°: I. J. Pachter and P. E. Nemeth, *J. Org. Chem.*, 1963, **28**, 1187.

Condensation of 4,6-diamino-5-nitroso-2-phenylpyrimidine with the anion of triethyl phosphonoacetate in tetrahydrofuran proceeded rapidly at room temperature to give 4-amino-2-phenylpteridin-7(8H)-one in 90% yield, in good



The condensation of phosphonate anions with 4-amino-5-nitrosopyrimidines may be considered as a special case of the Timmis reaction⁵ involving active methylene components. In the present instance, however, the phosphonate intermediate constitutes a doubly activated methylene compound which loses one of its activating groups (*i.e.*, the phosphonate) during the course of the condensation. The reaction proceeds with a variety of 2-substituents² and should prove a useful addition to the select number of pteridine syntheses which are unequivocal in their placement of substituents in the pyrazine ring.⁶

(Received, November 23rd, 1970; Com. 2023.)

† All attempts to condense 4-amino-5-nitrosopyrimidines with Wittig reagents were unsuccessful.

¹ For the previous paper in this series, see E. C. Taylor and K. Lenard, *Annalen*, 1969, **726**, 100.

² R. D. Youssefeyeh and A. Kalmus, *Chem. Comm.*, 1970, 1371.

³ W. S. Wadsworth, Jun. and W. D. Emmons, *J. Amer. Chem. Soc.*, 1961, **83**, 1733.

⁴ J. A. Maassen, Th. A. J. W. Wajer, and Th. J. de Boer, *Rec. Trav. chim.*, 1969, **88**, 5; H. Zimmer, P. J. Berecz, and G. E. Heuer, *Tetrahedron Letters*, 1968, 171.

⁵ For a review of the Timmis reaction, see T. S. Osdene, in "Pteridine Chemistry", eds. W. Pfeleiderer and E. C. Taylor, Pergamon Press Ltd., London, 1964, pp. 65—73.

⁶ E. C. Taylor and K. Lenard, *J. Amer. Chem. Soc.*, 1968, **90**, 2424.