

Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn19>

4'-Thio-RNA: Synthesis, Base Pairing Properties and Interaction with Dimerization Initiation Site of HIV-1

David Dukhan^b, Florence De Valette^b, Roland Marquet^a, Bernard Ehresmann^a, Chantal Ehresmann^a, François Morvan^b, Jean-Louis Barascut^b & Jean-Louis Imbach^b

^a Institut de Biologie Moléculaire et Cellulaire, 15 rue René, Descartes, 67084, Strasbourg

^b Laboratoire de Chimie Bio-organique, E. Bataillon, 34095, Montpellier, France

Published online: 04 Oct 2006.

To cite this article: David Dukhan, Florence De Valette, Roland Marquet, Bernard Ehresmann, Chantal Ehresmann, François Morvan, Jean-Louis Barascut & Jean-Louis Imbach (1999) 4'-Thio-RNA: Synthesis, Base Pairing Properties and Interaction with Dimerization Initiation Site of HIV-1, Nucleosides and Nucleotides, 18:6-7, 1423-1424, DOI: [10.1080/07328319908044740](https://doi.org/10.1080/07328319908044740)

To link to this article: <http://dx.doi.org/10.1080/07328319908044740>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

4'-THIO-RNA : SYNTHESIS, BASE PAIRING PROPERTIES AND INTERACTION WITH DIMERIZATION INITIATION SITE OF HIV-1


David Dukhan², Florence De Valette², Roland Marquet¹, Bernard Ehresmann¹, Chantal Ehresmann¹, François Morvan², Jean-Louis Barascut² and Jean-Louis Imbach^{*2}

¹Institut de Biologie Moléculaire et Cellulaire, 15 rue René Descartes 67084 Strasbourg

²Laboratoire de Chimie Bio-organique, E. Bataillon, 34095 Montpellier France

ABSTRACT: in the present paper, we describe the synthesis of a modified 9-mer oligonucleotide, 4'-S-r(UGUGCACCU) containing for the first time 4'-thio-guanosine units. This modified 9-mer was found to inhibit *in vitro* genomic RNA dimerization as well as the wild type RNA.

Antisense oligonucleotides constitute a promising class of antiviral drugs that offer a new and highly selective chemotherapeutic strategy to treat human diseases. To be successful, antisense therapeutics have to fulfill some criteria including nuclease resistance and affinity. We have showed that 4'-thio-β-D-oligoribonucleotides (4'-S-RNA) containing three of the four bases such as U, C and A exhibit very good nuclease resistance in comparison with wild-type RNA and bind more tightly to its complementary RNA strand than to its complementary DNA strand¹.



SCHEME

To explore further the biological applications of 4'-S-RNA series, we report the synthesis of 4'-S-r(UGUGCACCU) complementary to the loop of the HIV-1 Mal Dimerization Initiation Site (DIS) (Scheme). RNA dimerization is an ubiquitous property of retroviruses and is required for recombination. Furthermore, the DIS is involved in encapsidation and proviral DNA synthesis. Thus, the DIS represents a promising target for antiviral agents².

We found that 5-O-acetyl-2,3-O-isopropylidene-4-thio-D-ribofuranose is the best candidate for nucleosidic condensations on account of its anomer β orientation. This sugar was obtained in large quantities starting from D-gulono-1,4-lactone. Using Vorbrüggen condensation methods, the four 4'-thio-ribonucleosides (U, N4-bz-C, N6-bz-A, N2-Ac-G) were obtained. Each nucleoside was converted into its phosphoramidite derivatives for solid phase synthesis. Then, the 9-mer 4'-S-r(UGUGCACCU) was automatically synthesized, purified by HPLC and characterized by MALDI-TOF mass spectrometry.

Unlike the unmodified RNA 9-mer, this 4'-thio-RNA oligonucleotide was able to form a stable duplex by a self-association phenomenon (six bases in row), suggesting that Watson-Crick interactions are stabilized by the S \rightarrow O substitution. We studied the ability of this modified series targeted against the DIS to interfere a pre-formed viral RNA dimer. The 4'-thio-RNA was found to inhibit *in vitro* genomic RNA dimerization as the natural 9-mer, provided that the self-association was disrupted by a treatment involving heating and rapid cooling, before addition to the genomic RNA dimer. All these data indicate that 4'S-RNA can be considered as good candidates in antisens approach.

REFERENCES

1. Leydier, C.; Bellon, L.; Barascut, J.-L.; Morvan, F.; Rayner B.; Imbach J.-L. *Antisense Res. Dev.*, **1995**, 5: 167-174.
2. Skripkin, E.; Pallart, J.-C.; Marquet, R.; Blumenfeld, M.; Ehresmann, B.; Ehresmann, C.; *J. Biol. Chem.*, **1996**, 271(46): 28812-28817.