and dried to give 5.20 g (85%, mp 235-248°). Recrystallization from EtOH gave material with mp 249-251° dec. The uv spectral data are given in Table III. Anal. (C₆H₇ClN₆O₂) C, H, N.

Methyl 3-Hydroxy-6-chloropyrazinecarboxylate (XII).—A solution of NaNO₂ (7.0 g, 0.1 mole) in concentrated H₂SO₄ (75 ml) was added with stirring to a mixture of IIa (18.7 g, 0.10 mole) in concentrated H₂SO₄ (75 ml), and the resulting solution was stirred for 1 hr. The reaction mixture was poured over ice $(500~{\rm g})$ and the resulting aqueous solution was extracted with four 250-ml portions of EtOAc. The EtOAc was dried $(MgSO_4)$ and evaporated under reduced pressure to give 18.0 g (96%), mp 122-124°. Recrystallization from methylcyclohexane gave material with mp 127-129°. Anal. (C₆H₃ClN₂O₃) C, H, N. Pyrazinyl-1H-1,2,4-triazoles (VIII).—The general procedures

for the preparation of the (pyrazinecarboxamido)guanidines (I) applies as well to the preparation of the triazoles since, in most instances, a mixture of the two was obtained. This mixture was readily separated by taking advantage of the amphoteric properties of the triazoles. Routes B and E produced the least amount of VIII, undoubtedly due to the milder reaction conditions. Typical examples follow.

3-Amino-5-(3-amino-5-trifluoromethylpyrazinyl)-1H-1,2,4-triazole (VIIId).-Aminoguanidine hydrochloride (15.22 g, 0.137 mole) was added to a solution of Na (2.88 g, 0.125 g-atom) in MeOH (150 ml) and the mixture was stirred at room tem-perature for 1 hr. The mixture was filtered to remove NaCl and the filtrate was evaporated under reduced pressure to a thick paste. II (X = H, Y = CF₃, 5.52 g, 0.025 mole) was added and this mixture was heated on the steam bath for 2 min. H₂O (50 ml) was added and the mixture was filtered. This solid was Ik. The filtrate was neutralized with HOAc and the precipitate was filtered, washed (H_2O) , and dried to give VIIId, 0.97 g.

3-Amino-5-(3-amino-6-chloropyrazinyl)-1H-1,2,4-triazole (VIIIa). Method F.—Ia (4.0 g, 0.0175 mole) was pulverized and placed in a large test tube. A stream of N_2 was admitted and the tube was heated to 290° for 30 min. After cooling, the product was dissolved in 5% HCl and clarified with Darco. This solution was made strongly basic with 10% NaOH and again treated with Darco. This solution, when neutralized with HOAc and cooled to 0°, gave VIIIa.

Method G.—X^{1a} (0.5 g, 0.0016 mole) was dissolved in 5% HCl and warmed on the steam bath for 15 min. Neutralization of the cooled (0°) solution gave 0.20 g of VIIIa.

Pyrazine Diuretics. VII. N-Amidino-3-substituted Pyrazinecarboxamides

JAMES H. JONES, WILBUR J. HOLTZ, AND EDWARD J. CRAGOE, JR.

Merck Sharp & Dohme Research Laboratories, Division of Merck & Co., Inc., West Point, Pennsylvania

Received October 2, 1968

The synthesis of a series of N-amidino-3-substituted pyrazinecarboxamides, principally by the reaction of a methyl 3-substituted pyrazinecarboxylate with guanidine, is described. The intermediate 3-substituted pyrazinecarboxylates were generally prepared by a nucleophilic displacement reaction involving the appropriate 3-bromopyrazinecarboxylates which in turn were prepared from the corresponding 3-aminopyrazinecarboxylates. When the 3 substituent was methoxy, mercapto, methylmercapto, or substituted amino the compounds were generally less active than their 3-amino analogs in the normal or adrenalectomized DOCA loaded rats. The 3-hydroxy compounds were exceptions since they were as potent as their 3-amino analogs in the latter test.

Certain N-amidino-3-aminopyrazinecarboxamides¹ possess interesting and useful diuretic properties; therefore, it was of interest to determine the effect of various substituents in the 3 position on the diuretic activity.

The N-amidino-3-substituted pyrazinecarboxamides (IVa-p) examined in this study were prepared by the reaction of a methyl 3-substituted pyrazinecarboxylate (III) with guanidine according to the method described earlier² (see Scheme I). An exception to this method, noted in Scheme I, involves the synthesis of N-amidino-3-hydroxy-6-chloropyrazinecarboxamide (IVq) by the action of nitrous acid on the corresponding 3-amino analog (V). It is interesting to note that there is no attack on the guanidine moiety, even in the presence of excess nitrous acid.

The most useful method for the preparation of the intermediate methyl 3-substituted pyrazinecarboxylates (III) involved the nucleophilic displacement of the 3-halogen of the methyl 3-bromopyrazinecarboxylates (II). A wide variety of nucleophiles attack the 3-position halogen without affecting the halogen in the 6 position even when the reagent was present in excess. In an attempt to determine if displacement was occurring to

any extent at the 6 position, methyl 3-bromo-6chloropyrazinecarboxylate (IIb) was treated with NH₃ in DMSO. The progress of the reaction was checked readily by the periodic examination of a reaction mixture sample using tlc. The only product that could be detected, and eventually isolated in good yield, proved to be methyl 3-amino-6-chloropyrazinecarboxylate (Ib). Compounds IIId, e, and j were prepared by diazotization³ of the appropriate methyl 3aminopyrazinecarboxylate in concentrated H_2SO_4 followed by treatment of the diazonium salt with methanol or water to introduce a 3-methoxy or 3-hydroxy group. Methyl 6-bromo-3-methylaminopyrazinecarboxylate (IIIh) was prepared via 1,3-dimethyllumazine⁴ which was hydrolyzed to 3-methylaminopyrazinecarboxylic acid.⁵ then esterified, and, finally, brominated.

Ellingson and Henry⁶ have reported the preparation of methyl 3-bromopyrazinecarboxylate (IIa) by diazotization of the 3-amino compound⁷ (Ia) in 48% HBr containing Br₂. This method was readily adapted to the synthesis of compounds Ib-e by adding sufficient acetic acid to assure the dissolution of the ester in the reaction medium.

⁽¹⁾ J. B. Bicking, J. W. Mason, O. W. Woltersdorf, Jr., J. H. Jones, S. F. Kwong, C. M. Robb, and E. J. Cragoe, Jr., J. Med. Chem., 8, 638 (1965), paper I of this series.

⁽²⁾ E. J. Cragoe, Jr., O. W. Woltersdorf, Jr., J. B. Bicking, S. F. Kwong, and J. H. Jones, ibid., 10, 66 (1967).

⁽³⁾ This method has been described by A. E. Erickson and P. E. Spoeri, J. Am. Chem. Soc., 68, 401 (1946).

⁽⁴⁾ A. Albert, D. J. Brown, and H. C. S. Wood, J. Chem. Soc., 2066 (1956). (5) D. J. Brown and N. W. Jacobsen, *ibid.*, 4413 (1961).
(6) R. C. Ellingson and R. L. Henry, J. Am. Chem. Soc., 71, 2800 (1949).

⁽⁷⁾ R. C. Ellingson, R. L. Henry, and F. G. McDonald, ibid., 67, 1711 (1945).



Structure–Activity Relationships.—The N-amidino-3-substituted pyrazinecarboxamides (IV) that were prepared in this study were assayed⁸ for their ability to inhibit the decrease in the urinary ratio of Na/K produced by desoxycorticosterone acetate (DOCA) using the adrenalectomized rat according to the method described previously.^{1,2} The compounds were routinely administered subcutaneously but similar results were obtained when intraperitoneal or oral routes were used. The activity scores, which are presented in Table I, are in accordance with the scoring system described earlier.^{1,2}

Using this assay as the criterion of evaluation it can be seen that substitution of the 3-amino nitrogen of N-amidino-3-amino-6-bromopyrazinecarboxamide (VI) (15 in paper I¹ scored ± 2) results in reduction of activity. Furthermore, the larger groups have a greater effect than the smaller ones; thus, IVf is less

(8) Drs. M. S. Glitzer and S. L. Steelman and their associates supplied part of these data; the remainder was supplied by Dr. J. E. Baer and his associates.

	1 ABLE 1		
	BIOLOGICAL RESULTS		
IV	Rat DOCA-inhibs score ^{a}	Normal rat score ^b	
24		+1	
Ь	+1	0	
С	±i=	0	
\mathbf{d}	0	÷tr	
е	()	+-1	
ſ	<u>.</u> ±.	0	
g	+1	0	
h	+2	+1	
i	4:	t.:	
j	0		
k	and a second sec	()	
1	:1:	0	
m	+ 4	+ 1	
11	-+- 1	+- 1	
0	+3	+2	
р	:ta	+2	
(1	zir.	+3	

713

^a The DOCA-inhibition score¹ is the dose producing reversal of the DOCA Na/K effect: $+4 = <10 \ \mu g/rat$, +3 = 10-50, +2 =51-100, +1 = 101-800, $\pm = 800$, 0 = no activity at 800 μg . Compounds which scored 0 were tested only at a maximum dose of 800 $\mu g/rat$; thus, the possibility exists that activity would be observed at higher doses. ^b Activity¹⁰ based on increase of urinary electrolyte and volume over control values referred to standards: +3 = activity of hydrochlorothiazide, +2 = chlorothiazide, 0 = controls. Compounds with activities between chlorothiazide and controls are scored +1 or \pm .

active than IVg and h. Similarly, a considerable reduction in activity is observed upon substitution of the 3-amino group of N-amidino-3-amino-6-chloropyrazinecarboxamide (V) (**1** in paper I¹ scored +3) with relatively large groups (IVb, k, and l). On the other hand substitution of the 3-amino nitrogen of the more potent 5-amino series, *i.e.*, N-amidino-3-amino-5ethylamino- (or dimethylamino-) 6-chloropyrazinecarboxamide (VII and VIII) (IIa-3 and IIa-14 in paper II² scored +3), with groups such as allyl or ethyl (IVm and o) produced compounds with potencies equal to or greater than their parents. Substituents bearing hydroxy or alkoxy groups were detrimental to activity (IVp and n).



Replacement of the amino group of V by SH (IVc) or the amino group of VI by SCH_3 (IVi) produced a marked reduction in activity. Similar decreases in activity were seen when the amino group of VI. V, or VIII was replaced by OH (IVj, q, and e) or OCH₃ (IVd).

The compounds recorded in Table I also were tested in normal rats using the intraperitoneal route of administration.⁹ The assay and the scoring system have been described previously.¹⁰ The relative activities obtained in this test generally paralleled those observed

⁽⁹⁾ Dr. J. E. Baer and his associates conducted these studies.

⁽¹⁰⁾ J. H. Jones, J. B. Bicking, and E. J. Cragoe, Jr., J. Med. Chem., 10, 899 (1967).

in the DOCA-inhibition assay. The exceptions were the 3-hydroxy compounds (IVe, j, and q) which were markedly more active (comparable to their amino analogs V, VI, and VII) and IVm which was considerably less active in the normal rat assay than in the DOCA-inhibition assay.

Experimental Section¹¹

Details of the syntheses of the new compounds are presented. Where several compounds of one type have been prepared by a particular method, only one example is given. Pertinent data regarding each compound are recorded in Table II.

Method 1. Methyl 3-bromopyrazinecarboxylates (II). Methyl 3-Bromo-6-chloropyrazinecarboxylate (IIb).—A suspension of Ib (18.7 g, 0.1 mole) in a solution of 48% HBr (114 ml) and AcOH (30 ml) was cooled to 0°, stirred, and treated with a solution of Br₂ (15 ml) in AcOH (30 ml) over a period of 45 min. Then a solution of NaNO₂ (17.4 g, 0.1 mole) in H₂O (39 ml) was added while maintaining the temperature at 0°. Stirring was continued for 30 min and then the excess Br₂ was destroyed by the dropwise addition of a 30% aqueous solution of NaHSO₃ (150 ml). The product, which separated, was recovered by filtration, washed well with cold H₂O, dried, and recrystallized.

Method 2. Methyl 3-Substituted Pyrazinecarboxylates (III). Method 2A. Methyl 3-(2-Dimethylaminoethylamino)-6-chloropyrazinecarboxylate (IIIb).—To 2-dimethylaminoethylamine (3.4 g, 0.04 mole) in DMSO (20 ml) was added IIb (5.0 g, 0.92 mole), the solution was stirred at room temperature for 1.5 hr, then poured into H_2O (100 ml), and the product which separated was removed by filtration, dried, and purified by recrystallization.

For the synthesis of related compounds, the pure liquid or gaseous amine was used.

Method 2B. Methyl 3-Mercapto-6-chloropyrazinecarboxylate (IIIc).—A suspension of Na₂S·9H₂O (4.8 g, 0.02 mole) and S (5.0 g, 0.156 g-atom) in EtOH (40 ml) was refluxed for 30 min and cooled. Then IIb (5.0 g, 0.01 mole) was added and the solution was stirred at ambient temperature for 1.5 hr. The reaction mixture was poured into H₂O (100 ml), the precipitated S was removed by filtration, and the filtrate was acidified with HCl to precipitate the product which was removed by filtration and purified by crystallization.

Method 2C. Methyl 3-Methylmercapto-6-bromopyrazinecarboxylate (IIIi),—CH₃SH gas (1.28 g, 0.026 mole) was dissolved in a solution of MeOH (75 ml) containing 20% NaOH (5.3 ml, 0.026 mole). IIc (7.8 g, 0.026 mole) was added and the reaction mixture was stirred at ambient temperature for 1 hr, then poured into H₂O (100 ml). The precipitated product was purified by recrystallization.

Method 2D. Methyl 3-Methoxy-6-chloropyrazinecarboxylate (IIId).—To a solution of Ib (18.7 g, 0.1 mole) in concentrated H_2SO_4 (75 ml) maintained at 0–5° was added a solution of NaNO₂ (9.0 g, 0.13 mole) in concentrated H_2SO_4 (75 ml) and the reaction mixture was stirred at 5–10° for an additional 30 min. The reaction mixture was carefully added to MeOH (11.), refluxed for 1.5 hr, then concentrated *in vacuo* to 500 ml and poured onto ice (1500 g). The aqueous solution was extracted with CHCl₃ (three 300-ml portions), then the organic phase was washed with 2% NaOH (three 200-ml portions), dried, and evaporated *in vacuo* to an oil which was purified by recrystallization.

Method 2E. Methyl 3-Hydroxy-6-bromopyrazinecarboxylate (IIIj).—Diazotization was carried out exactly as described in method 2D using Ic (4.6 g, 0.02 mole), concentrated H₂SO₄ (30 ml), and NaNO₂ (1.4 g, 0.02 mole). Then the reaction mixture was poured directly onto ice (200 g). The resulting mixture was extracted with CHCl₃ (two 200-ml portions), the CHCl₃ layer was extracted with $2\frac{G}{C}$ NaOH (two 100-ml portions), and the aqueous phase was acidified with HCl which precipitated the product. The crude material was purified by recrystallization.

Method 2F. Methyl 3-Methylamino-6-bromopyrazinecarboxylate (IIIh).—1,3-Dimethyllumizine⁴ was hydrolyzed using aqueous NaOH (10%) to give 3-methylaminopyrazinecarboxylic

TABLE II

METHYL 3-BROMOPYRAZINECARBOXYLATES (II), METHYL 3-SUBSTITUTED PYRAZINECARBOXYLATES (III), AND N-AMIDINO-3-SUBSTITUTED PYRAZINECARBOXAMIDES (IV)

	Syn	Recrystn	%		
No.	$method^a$	solvent	\mathbf{yield}	Mp, °C	$\mathbf{Formula}^d$
IIa ⁶					
$_{\rm IIb}$	1	Petr ether	56	38-40	$C_6H_4BrClN_2O_2$
IIc	1	EtOH-H ₂ O	60	66-68	$C_6H_4Br_2N_2O_2$
IId	1	Cyclohexane	51	98-99	C8H9BrClN3O2
IIe	1	C_6H_6	60	160 - 162	C ₈ H ₉ BrClN ₃ O ₂
IIIa	2B	ь	48	124 - 125	$C_6H_6N_2O_2S^b$
IIIb	2A	<i>i</i> -PrOH	33	116-118	C10H16ClN4O2 ^e
IIIc	2B	EtOH	87	80-82	$C_6H_5ClN_2O_2S^f$
IIId	2D	$EtOH \cdot H_2O$	13	45	$C_7H_2ClN_2O_3^b$
IIIe	$2\mathrm{E}$	MeOH	35	140-141	$C_8H_{10}ClN_3O_3$
IIIf	2A	$EtOH \cdot H_2O$	73	88-90	C11H14BrNsO2
IIIg	2A	$EtOH \cdot H_2O$	77	80-82	C8H10BrN3O2
IIIh	2F	i-rPOH	75	181-183	C7H8BrN3O2
IIIi	2C	EtOH	80	135 - 136	C7H7BrN2O2S
IIIj	$2\mathrm{E}$	Hexane	75	120.5 - 121.5	C6H5BrN2O3
IIIk	2A	EtOH	67	152 - 154	$C_{13}H_{11}Cl_2N_3O_2$
1111	2A	EtOH	98	105-106	C14H14ClN3O3
IIIm	2A	i-PrOH	78	100-102	$C_{11}H_{18}ClN_4O_2$
IIIn	2A	Hexane	68	89-90	C11H17ClN4O3
III_0	2A	Cyclohexane	92	93-95	$C_{10}H_{16}ClN_4O_2$
$_{\rm IIIp}$	2A	BuCl	88	103 - 105	$C_{10}H_{1b}ClN_4O_3$
IVa	3	$EtOH \cdot H_2O$	33	244–246 dec	$(C_6H_7N_5OS \cdot HCl)_2 \cdot H_2O$
IVb	3	HCl-NaOH	16	210-212 dec	C10H16ClN7O
IVe	3	HCl-NaOH	54	260 dec	$(C_6H_6ClN_5OS)_2 \cdot H_2O$
IVd	3	Dil HCl	47	214-216 dec	C7H3ClN5O2 · HCl
IVe	3	Dil HCl	20	231.5-233.5	$C_{\$}H_{11}ClN_{6}O_{2}\cdot HCl$
				dec	
IVf	3	HCl-NaOH	81	220222 dec	C11H15BrN5O
IVg	3	HCl-NaOH	30	216-218 dec	CsH11BrN6O
IVh	3	HCl-NaOH	50	230–232 dec	C7H9BrN6O
IVi	3	H_2O	55	275-278 dec	C7H8BrN5OS · HCl
IVj	3	$EtOH-H_2O$	75	>290	C6H6BrN5O2·HCl
IVk	3	$EtOH-H_2O$	40	245-248 dec	$C_{13}H_{12}Cl_2N_6O \cdot HCl$
IVI	3	EtOH-H ₂ O	43	200-202 dec	$C_{14}H_{15}ClN_6O_2 \cdot HCl$
IVm	3	H_2O	97	132 - 135	$C_{11}H_{16}CIN_7O \cdot HCl$
IVn	3	MeOH	80	219 - 220	$C_{11}H_{18}ClN_7O_2$
IVo	3	H_2O	67	241 - 242	$C_{10}H_{16}ClN_7O \cdot HCl$
IVp	3	H_2O	75	242 - 244	$\mathrm{C}_{10}\mathrm{H}_{16}\mathrm{ClN;}\mathrm{O}\cdot\mathrm{HCl}\cdot\mathrm{H}_{2}\mathrm{O}$
IVq	4	HCl-NaOH	81	257-259 dec	C6H6ClN5O2

^a See the Experimental Section for the number and letter that corresponds to each experimental method. ^b This material was used in the next step without further purification. ^c Dr. J. B. Bicking is responsible for this preparation. ^d All compounds were analyzed for C, H, N, except for IIIa and IIId. The analytical results for the elements were within $\pm 0.4\%$ of the theoretical values except where indicated. ^e C: calcd, 46.42; found, 45.95. ^f C: calcd, 35.21; found, 35.62.

acid⁵ which, in turn, was esterified using MeOH saturated with HCl. The ester was not purified but brominated directly by the procedure of Ellingson and Henry⁶ and the product was purified by crystallization.

Method 3. The preparation of the N-amidinopyrazinecarboxamides (IVa-p) was carried out exactly as described previously.²

Method 4. N-Amidino-3-hydroxy-6-chloropyrazinecarboxamide (IVq).—Compound V (6.42 g, 0.03 mole) was dissolved in H₂O (300 ml) by the addition of methanesulfonic acid (7.2 g, 0.075 mole), then stirred and cooled to 5-10°. A solution of NaNO₂ (2.01 g, 0.033 mole) in H₂O (20 ml) was added dropwise over a period of 1 hr. The reaction mixture then was stirred at room temperature for 30 min, heated to 60°, and filtered, and the filtrate was neutralized with NaOH to pH 7 which caused the product to separate. The product was removed by filtration and purified by reprecipitation.

Method 5. Methyl 3-Amino-6-chloropyrazinecarboxylate (Ib) from Methyl 3-Bromo-6-chloropyrazinecarboxylate (IIb).—A solution of IIb (1.0 g, 3 mmoles) in DMSO (3 ml) was stirred and heated on the steam bath while a stream of NH₃ was admitted over a period of 30 min. Addition of H₂O (15 ml) afforded a solid that was recrystallized from H₂O to yield 0.65 g (91%) of Ib melting 158-160°. Mixture melting point with authentic Ib was not depressed; the ir spectra of the compounds were identical.

⁽¹¹⁾ Mr. K B. Streeter, Mr. Y. C Lee, and their staff have provided the analytical data. The melting points are corrected (open capillaries).