

Oxidation of 5-Bromo- and 5-Fluoro-1,3-dimethyluracils
by Peroxomono- and Peroxodisulfate Ions


Toshio ITAHARA* and Akihiro NISHINO
Institute of Chemistry, College of Liberal Arts,
Kagoshima University, Korimoto, Kagoshima 890

Oxidation of 5-bromo-1,3-dimethyluracil by KHSO_5 gave 5,5-dibromo-6-hydroxy- and 5,5,6-trihydroxy-1,3-dimethyl-5,6-dihydrouracils and that of 5-fluoro-1,3-dimethyluracil by $\text{Na}_2\text{S}_2\text{O}_8$ gave a 6,6'-dimeric product.

5-Fluorouracil has been widely used in the chemotherapy of malignant neoplasms but has toxicity such as neurotoxicity.¹⁾ Hence, metabolism of 5-fluorouracil is of interest.¹⁾ Also, 5-bromouracil is a mutagenic compound and cells with DNA containing 5-bromouracil are deactivated by γ - and UV-radiation. The γ -radiation²⁾ and photoreaction³⁾ of 5-bromouracils have, therefore, been investigated. However, little attention has been paid to oxidation of these compounds. As a part of the investigation concerning a comparison of products on oxidation of nucleic acid bases and nucleosides between by peroxomonosulfate ion (HSO_5^-) and by peroxodisulfate ion ($\text{S}_2\text{O}_8^{2-}$),⁴⁾ we studied the oxidation of 5-bromo- and 5-fluoro-1,3-dimethyluracils as model compounds.

A solution of 5-bromo-1,3-dimethyluracil (1a) (1 mmol) and KHSO_5 (1 mmol) in water (50 ml) was heated at 70 °C for 5 h under argon atmosphere. The reaction mixture was submitted to chromatography on ODS-silica gel.⁴⁾ By elution with water, a fraction containing 5,5,6-trihydroxy-1,3-dimethyl-5,6-dihydrouracil (2) was obtained. Further elution with a mixture of water and acetonitrile led to the isolation of 1a (0.04 mmol, 4 %) and 5,5-dibromo-6-hydroxy-1,3-dimethyl-5,6-dihydrouracil (3)⁵⁾ (0.33 mmol, 33 % yield). The fraction containing 2 was further separated by gel-filtration chromatography (Tosoh HW-40) with water to give 2⁶⁾ (32 %). Similarly treatment of 5-fluoro-1,3-dimethyluracil (1b) with KHSO_5 gave 2 (11 %) together with recovered 1b (59 %). On the other hand, reaction of 1a (1 mmol) with $\text{Na}_2\text{S}_2\text{O}_8$ (1 mmol) in water (50 ml) at 80 °C for 5 h under argon atmosphere gave a complex reaction mixture, which was chromatographed on silica gel with ethyl acetate to give 1,3-dimethylparabanic acid (4)⁷⁾ (11

%), amalic acid (tetramethylalloxantine) (5)⁸⁾ (10 %), and recovered 1a (43 %). Under similar conditions, the reaction of 1b with $\text{Na}_2\text{S}_2\text{O}_8$ interestingly gave a 6,6'-dimeric product (6)⁹⁾ (18 %) together with 4 (3 %), 5 (3 %), and recovered 1b (68 %). The structure of 6 resembles that of the dimeric product from γ -radiation of thymidine¹⁰⁾ and the formation of 6 suggests that the oxidation by $\text{Na}_2\text{S}_2\text{O}_8$ is reasonably explained in terms of formation of a radical intermediate.



We thank Professor Kenichi Somekawa, Faculty of Engineering, Kagoshima University, for valuable support throughout the study.

References

- 1) R. B. Diasio, T. L. Beavers, and J. T. Carpenter, *J. Clin. Invest.*, **81**, 47 (1988) and references therein.
- 2) C. von Sonntag, "The Chemical Basis of Radiation Biology," Taylor and Francis, London (1987).
- 3) H. Sugiyama, Y. Tsutsumi, and I. Saito, *J. Am. Chem. Soc.*, **112**, 6720 (1990) and references therein.
- 4) T. Itahara and S. Koga, *Chem. Lett.*, **1991**, 85.
- 5) S. Y. Wang, *J. Org. Chem.*, **24**, 11 (1959).
- 6) 2: Oil; $^1\text{H-NMR}$ (D_2O) δ 3.00 (Me), 3.01 (Me), 5.28 (H-6); (d_6 -DMSO) δ 2.82 (Me), 2.85 (Me), 4.96 (H-6), 6.2-6.7 (broad, OH-5,5,6); $^{13}\text{C-NMR}$ (D_2O) δ 27.29, 28.27, 89.23, 92.13, 160.57, 176.61. The compound 2 was stable in D_2O for more than 2 days but labile in d_6 -DMSO.
- 7) T. L. Patton, *J. Org. Chem.*, **32**, 383 (1967).
- 8) H. Bredereck and W. Pfeleiderer, *Chem. Ber.*, **87**, 1119 (1954).
- 9) 6: mp 290-296 °C; $^1\text{H-NMR}$ (d_6 -DMSO) δ 3.15 (Me), 3.26 (Me), 9.30 (OH); (D_2O) δ 3.27 (Me), 3.39 (Me); ($\text{D}_2\text{O}+\text{NaOH}$, pD 13) δ 3.22 (Me), 3.37 (Me); $^{13}\text{C-NMR}$ (d_6 -DMSO) δ 28.35, 32.47, 123.05, 130.74, 149.91, 159.91; MS, m/z (relative intensity) 310 (M^+ , 16), 309 (100). Anal. Found: C, 46.39; H, 4.59; N, 17.78 %. Calcd for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_6$: C, 46.45; H, 4.55; N, 18.06 %.
- 10) S. Nishimoto, H. Ide, K. Nakamichi, and T. Kagiya, *J. Am. Chem. Soc.*, **105**, 6740 (1983).

(Received September 12, 1991)