in lessening the ventricular fibrillation induced in anesthetized dogs by immersion tank hypothermia was measured. Quinidine was the reference drug used and the active compounds are listed in Table IX.

(20) E. T. Angelakos and A. H. Hegnauer, J. Pharmacol. Exptl. Therap., 127, 137 (1959).

Acknowledgment.—The authors thank Dr. R. E. Bowman for many helpful discussions, Mr. F. H. Oliver for the microanalyses, Miss E. M. Tanner for the physical measurements, Dr. Vandenbelt for the n.m.r. spectra, and Drs. Corne, Chen, and Wheelock, and Mr. C. Schneider for the pharmacological testing.

New Analgesic N-Substituted Carboxamides

OTIS E. FANCHER, SHIN HAYAO, WALLACE G. STRYCKER, AND L. F. SANCILIO

Chemical Therapeutics Research Laboratory, Miles Laboratories, Inc., Elkhart, Indiana

Received April 9, 1964

A series of N-substituted propionanilides was synthesized as potential analgesics. The most active compound, N-[1-methyl-2-(4-hydroxy-4-phenylpiperidino)ethyl]propionanilide oxalate (27), was approximately 150 times more active than morphine by the artery-clip assay method in mice.

In the course of our search for new, potent analgesics, we have synthetized a group of N-substituted carboxamides (I), mainly N-substituted propionanilides, which are structurally related to methadone in that the quaternary carbon atom and a phenyl group of methadone are replaced with a tertiary nitrogen. During the course of our work, two potent analgesics, phenampromid and diampromid,¹ were reported. Both compounds are N-substituted propionanilides. Similar compounds were described later by Shigematsu² and by Carabateas.³ Among our compounds, 1 and 2 of Table I were described in a recent patent.⁴

The compounds reported here are listed in Table I and may be represented by the generic formula R¹- $C_6H_4(CH_2)_nN(COR^2)CHR^3CHR^4+B$ (I). The isomerically pure compounds were prepared readily by reduction of the corresponding amides, with lithium aluminum hydride followed by acylation. When α -(1,2,3,6tetrahydro-4-phenyl-1-pyridyl)propionanilide (II) was reduced in this way, followed by propionylation, the product isolated as the oxalate was unexpectedly the saturated compound (III). The latter was also synthesized by reducing α -(4-phenylpiperidino)propionanilide (IV), followed by propionylation and formation of the oxalic acid salt.

Salts (III), prepared by both routes, were identical



(1) W. B. Wright, Jr., H. J. Brabander, and R. A. Hardy, Jr., J. Am. Chem. Soc., **81**, 1518 (1959); J. Org. Chem., **25**, 1033 (1960); *ibid.*, **26**, 476, 485 (1961).

(3) P. M. Carabateas, W. F. Wetterau, and L. Grumbach, J. Med. Chem., 6, 355 (1963).

(4) O. E. Fancher and S. Hayao, U. S. Patent 3,037,982 (June 5, 1962).

by mixture melting point determination and gave identical ultraviolet spectra which showed two maxima at 257.5 m μ (ϵ 1050) and 264 m μ (ϵ 832).

In contrast, the reduction of 1-(α -anilinopropionyl)-1,2,3,6-tetrahydro-4-phenylpyridine (V), an isomer of II, gave the unsaturated compound VI. In support of the unsaturated structure was a strong absorption maximum at 244.5 m μ (ϵ 14,600),⁵ indicating the presence of a double bond conjugated with the aromatic ring. It is reported that a double bond in the systems ArC=CCO- or ArC=CN-⁶ may be reduced with 1 it is in the grouping C₆H₅C=CCH₂- has not been described previously.



⁽⁵⁾ α -Methylstyrene has $\lambda_{\max}^{E:OB}$ 243 m μ (ϵ 11,500); A. E. Gillam and E. S. Stern, "An Introduction to Electronic Absorption Spectroscopy in Organic Chemistry," 2nd Ed. Edward Arnold Ltd., London, 1957, p. 277. (6) W. G. Brown, Org. Reactions, **6**, 480 (1951).

 ⁽²⁾ N. Shigematsu, Yakugaku Zasshi, 81, 423 (1961); *ibid.*, 81, 815
 (1961); N. Sugimoto, K. Okumara, N. Shigematsu, and G. Hayashi, *Chem. Pharm. Bull.* (Tokyo), 10, 1061 (1962); G. Hayashi, N. Shigematsu, and Y. Kowa, Yakugaku Zasshi, 81, 62 (1963).

		Z	9.68, 0.57	9.54	9.46	8.74	8.74	9.30	8,46 8	9.05	8.87	16°£	$\frac{\infty}{2}$	7.79	7.96	6.32	6.39	6.46	6.48
		Found, % H	7.03	7.21	7.20	6.53		7.18	6.81	6.82	7.13	6.52	3. SN	6.46	7.05	6.65	6, 73	7.24	7 30
		C]	65.1	65.0	65.6	60.4		65.8	64.5	63.6	64.0	+ 22	0110	59.4	6769	68.2	68.4	2.73	87 S
		N	9.52	9.52	9.23	8.83	8.83	9.23	8,45	16.8	8.91	5.97	8.12	7.52	7.92	6.39	6.39	6.36	6.36
		Saled., % H	7.03	7.03	7.25	6.30		7.25	7.04	7.00	7.01	6.61	5.99	6.62	6.96	6.84	6.84	7.28	1.28
		c)	65.3	65.3	65.9	60.5		0.09	65.2	63.6	63.6	53.8	58°.0	60.1	70.0	68.4	68.5	68.2	68.2
		M.p., °C.	150–152 dec.	172.5-173.5 dec.	131–133 dec.	162-163	108–109 dec.	175-176 dec.	132-133.5	143144	166.5–168.5 dec.	209 dec.	203–203 . 5 dec.	216 dec.	216-216.5 dec.	159–160 dec.	193.5–194.5 dec.	152–153 dec.	186-487 dec.
EI	COR?)CHR3CHR4B	Formula	${\rm C}_{22}{\rm H}_{29}{\rm N}_{3}{\rm O}\cdot{\rm C}_{2}{\rm H}_{2}{\rm O}_{4}{\rm a}$	$C_{22}H_{29}N_{3}O\cdot C_{2}C_{2}O_{4}$	$\mathrm{C}_{23}\mathrm{H}_{31}\mathrm{N}_{3}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{22}\mathrm{H}_{28}\mathrm{ClN}_{3}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{22}\mathrm{H}_{28}\mathrm{ClN_3O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{23}\mathrm{H}_{31}\mathrm{N}_{3}\mathrm{O}\cdot\mathrm{C}_{9}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{23}\mathrm{H}_{31}\mathrm{N}_3\mathrm{O}_2\cdot\mathrm{C}_4\mathrm{H}_4\mathrm{O}_4{}^b$	$\mathrm{C}_{23}\mathrm{H}_{31}\mathrm{N}_3\mathrm{O}_2\cdot\mathrm{C}_2\mathrm{H}_2\mathrm{O}_4$	$\mathrm{C}_{23}\mathrm{H}_{31}\mathrm{N}_{3}\mathrm{O}_2\cdot\mathrm{C}_2\mathrm{H}_2\mathrm{O}_4$	$\mathrm{C}_{17}\mathrm{H}_{27}\mathrm{N}_{3}\mathrm{O}\cdot 2\mathrm{C}_{2}\mathrm{H}_{3}\mathrm{O}_{4}$	$\mathrm{C}_{\mathrm{sl}}\mathrm{H}_{\mathrm{sf}}\mathrm{N}_{\mathrm{s}}\mathrm{O}\cdot\mathrm{2}\mathrm{C}_{\mathrm{s}}\mathrm{H}_{\mathrm{s}}\mathrm{O}_{\mathrm{4}}$	$C_{24}H_{38}N_{8}O\cdot 2C_{2}H_{2}O_{4}$	$C_{29}H_{35}N_3O\cdot C_2H_2O_1$	${\rm C}_{23}{\rm H}_{28}{\rm N}_{2}{\rm O}\cdot{\rm C}_{2}{\rm H}_{4}{\rm O}_{4}$	$\mathrm{C}_{23}\mathrm{H}_{28}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{23}\mathrm{H}_{30}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$\mathrm{C}_{23}\mathrm{H}_{30}\mathrm{N}_2\mathrm{O}\cdot\mathrm{C}_2\mathrm{H}_2\mathrm{O}_4$
TABL	CH ₂) " N((u	0	0	1	0	0	0	÷	0	0	c	0	0	0	e	0	0	
	R	ĩ	N NC ₀ H ₅	N NC,H.	N N C,H,	N NC,H4CI-P	NC.,H,CI-0	N NC ₆ H ₁ CH ₇ ·P	N NC ₈ H.	N NC ₈ H.	NC,H.	N NCH	N NCH	N NCH ² CH ² C ⁴ H ²	N NCHIC ₆ H.).	N C.H.	$N = C_6 H_s$		N C ₀ H ₂
		184	Ш	CH_{3}	CH,	CH,	CH_{i}	CH_3	CH.	CH_3	CH_3	CH_z	$OH_{\rm s}$	CH_{s}	CH_{s}	II	CH_3	Н	CH_3
		εIJ	CH ₃	H	II	Ш	Н	Η	H	Η	Н	Η	Н	Η	Н	CH_{s}	Н	CH_3	н
		\mathbb{R}^2	${ m C}_{2}{ m H}_{5}$	C_2H_5	C ₂ H ₅	C_2H_6	C_2H_5	C_2H_5	$C_2 H_5$	C_2H_5	CJII,	$C_{2}H_{5}$	$\mathbf{C}_{\mathbf{b}}\mathbf{H}_{\mathbf{b}}$	$\mathbf{C}_2\mathbf{H}_5$	C_2H_5	$\mathrm{C}_{2}\mathrm{H}_{5}$	C_2H_5	$\mathbf{C}_2\mathbf{H}_5$	C_2H_5
		Rı	Н	II	II	H	Н	Н	p-CH ₃ O	m-CH ₃ O	0-CH3O	H	Н	Н	Ш	Н	Η	Н	Н
		No.	1	57	ŝ	4	ũ	9	l-	x	6	10	11	15	13	t I	15	16	21

Vol. 7

05	33	88	92	72	75		57	07	19 03	5 20	62	65	41	28	12	83	93	68	4.
6.0	6.5	5.5	5.0	5.	5.		5.	9.1	97.1		0.0	8	5.	9	.9	9.9	5.0	8	с 1 , н.
7.56	7.49	6.45	6.55	7.18	7.24	6.74	6.30	7.10	6.91 7.80 6.64	7.14	7.98	6.96		6.86	6.34	6.57	7.27	8.80	nd: He
67.9	68.2	71.1	71.1	67.4	65.8	73.6	71.1	65.3	65.6 68.1 56.3	00.9 65.9	69.3	57.6		67.8	62.6	66.3	63.6	66.1	Four
6.17	6.17	5.74	5.74	5.65	5.96		5.58	6.14	$6.14 \\ 6.95 \\ r_{r_{r_{r_{r_{r_{r_{r_{r_{r_{r_{r_{r_{r$	5.96	6.50	8.76	5.49	6.36	5.95	6.80	5.92	8.68	Cl, 11.3.
7.49	7.49	6.55	6.55	7.25	7.23	6.90	6.77	7.01	7.01 7.69	0.00 7.23	8.13	6.89		6.58	6.43	6.80	7.18	8.37	cd.: H
68.7	68.7	71.4	71.4	67.5	66.4	74.2	71.7	65.8	65.8 68.5 7.7	50.7 66.4	69.7	57.6		68.1	62.8	67.0	63.4	67.0	è. " Cal
142–144 dec.	149–151 dec.	160–161 dec.	198–199 dec.	145-146	120.5–121.5 dec.	198.5–200 dec.	174–175.5 dec.	103–105 dec.	165.5-167 dec. 215-217 dec.	12-04 uec. 142-144 dec.	189–191 dec.	185–185.5	165.5–167 dec.	200–202 dec.	196-198	145.5–147 dec.	149150 dec.	156–157 dec.	Aethiodide. / Sulfamate
¹ 24H ₂₂ N ₂ O · C ₂ H ₂ O ₄	$^{2_4}\mathrm{H}_{32}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$^{\prime}_{27}\mathrm{H}_{30}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$_{27}\mathrm{H}_{30}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	${}_{24}\mathrm{H}_{32}\mathrm{N}_{2}\mathrm{O}_{2}\cdot\mathrm{C}_{4}\mathrm{H}_{4}\mathrm{O}_{4}$	${}_{24}\mathrm{H}_{32}\mathrm{N}_{2}\mathrm{O}_{2}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	${}_{36}\mathrm{H}_{38}\mathrm{N}_{2}\mathrm{O}\cdot\mathrm{HCl}^{c}$	$^{12}_{28} m H_{22} m N_{2} m O\cdot C_{2} m H_{2} m O_{4}$	$_{23}{ m H}_{30}{ m N}_{2}{ m O}_{2}\cdot{ m C}_{2}{ m H}_{2}{ m O}_{4}$	$_{23}^{23}\mathrm{H}_{30}^{20}\mathrm{N}_{2}^{2}\mathrm{O}_{2}\cdot\mathrm{C}_{2}^{2}\mathrm{H}_{2}^{2}\mathrm{O}_{4}$ $_{23}^{23}\mathrm{H}_{30}^{2}\mathrm{N}_{2}\mathrm{O}_{2}\cdot\mathrm{HCl}^{d}$	23H30N2O2 • CH31 24H32N2O2 • C2H2O4	25H34N2O2.HCl	23H20N2O2. NH2SO2H ^J	$^{1}_{27}\mathrm{H}_{36}\mathrm{N}_{2}\mathrm{O}_{2}\cdot\mathrm{C}_{2}\mathrm{H}_{2}\mathrm{O}_{4}$	$_{26}$ H $_{28}$ N $_{2}$ O $_{3}$ ·HCl	$_{33}H_{41}N_{3}O_{3}\cdot 2C_{2}H_{2}O_{4}$	21H26N2O.C2H2O4	$^{23}_{23}H_{32}N_{2}O_{3}\cdot C_{2}H_{2}O_{4}$	18H26N2O · HCl9	md: HCl, 9.04.
0	0	\circ	0	U	5	0	0	Ð	000	00	0	0	0	0	0	0	0	0	Fot
1 C	1 0	0 C	0	0	0	0	0	0	000	1	2 G	0	0	0	0	0	0	0	9.08. Fot
$N O_{G,H_5}$ 1 C	N C ₆ H ₅ 1 C	N -CoHe 0 C	N C.H. 0 C	$N \longrightarrow C_{c}H_{s} = 0$ C	$N \longrightarrow C_{cH_s} 0 C$		$N \longrightarrow C_{0}H_{5} = 0$ C	N Coll 0 C		N CH 1 C	$N \sim C_{c,H_s} $ 2 C	$\sqrt{\sum_{C,H_5}}$ 0 C	$N \longrightarrow C_{G,H_1} 0 C$	$N \sim C_{c,H_1} 0 C$	N CH 0 C		NiCHJACH(CH_J)CH_2 0 CH_3 0 C	$N(CH_2CH=CH_2)_2$ 0 C	: HCl, 8.67. ⁴ Caled.: HCl, 9.08. For
H $N = N = 0.05$ H C	CH_3 $N O_{c_6H_5}$ 1 C	H $N \longrightarrow C_{c}H_{c}$ 0 C	CH_3 $N \longrightarrow -C_0H_5$ 0 C	CH_3 $N \longrightarrow C_0H_5$ 0 C	CH_3 $N - c_3H_5$ 0 C	C_6H_5 $N \longrightarrow 0$ C	C_6H_5 $N \longrightarrow C_6H_5$ 0 C	CH_3 $N \sim C_{cH_3} = 0$ C	H $N \sim C_{\rm c,H}^{\rm OH} = 0$	CH_3 $N \sim CH_2 = 1$ C	H $N \sim C_{C,H_3}$ 2 C	H $(\sim C_{C_{0}H_{5}}^{OH}) = 0$ C	H $N \sim C_{G,H_1} = 0$ C	H $N \sim C_{C_0,H_5}^{OH}$ 0 C	H $N \sim C_{C_0H_5}^{OH}$ 0 C	CH ₃ 0 C	H N(CH.)(CH.)(CH.)(CH.) OCH3 0 C	$CH_3 \qquad N(CH_2CH=CH_2)_2 \qquad 0 C$	Found: HCl, 8.67. d Caled .: HCl, 9.08. For
CH ₃ H $N = N = N = C$	H CH_{3} $N \longrightarrow C_{6}H_{5}$ 1 C	CH ₃ H $N \longrightarrow C_{c}H_{c}$ 0 C	H CH_3 $N - c_sH_s = 0$ C	H CH_3 $N - C_{a}H_5$ 0 C	H CH ₃ $(\sqrt{-c_{H_5}} - 0)$ C	H C_6H_5 N 0 C	H C_6H_s $N \longrightarrow C_6H_s$ 0 C	H CH_3 $N \sim C_{GH_3} = 0$ C	CH ₃ H $N \sim C_{GH_5}^{OH}$ 0 C C	H CH ₃ $N \sim CH_3$ I C	CH_3 H $N \sim C_{C_0H_3} = 2$ C	CH_3 H $N \sim C_{C,H_3} O C$	CH ₃ H $N \sim C_{C,H_1}^{OH}$ 0 C	CH ₃ H $N \sim C_{C_0,H_1}^{OH}$ 0 C	CH_{3} H $N \sim C_{C,H_{5}}^{OH}$ 0 C	H CH_3 0 C	H H N(CH ₃)CH ₃ CH ₄ \rightarrow 0 CH ₃ 0 CH ₃	H CH_3 $N(CH_2CH=CH_2)_2$ 0 C	HCl, 8.68. Found: HCl, 8.67. ^d Caled.: HCl, 9.08. For
C_2H_5 CH_3 H $N_{C_4H_5}$ 1 C	C_2H_6 H CH_3 N C_6H_6 1 C	C_6H_5 CH_3 H $N \longrightarrow C_6H_6$ 0 C	$C_{c}H_{5}$ H CH_{3} N $C_{c}H_{5}$ 0 C	H_3O C_2H_3 H CH_3 $N \longrightarrow C_3H_5$ 0 C	$I_3O C_2H_5 H_1 CH_3 N - C_3H_5 O C$	C_6H_5 H C_6H_5 N 0 C	C_2H_5 H C_6H_5 N C_6H_5 0 C	C_3H_6 H CH_3 $N \sim C_{GH_3} = 0$ C	C_2H_5 CH_3 H $N \sim c_{c,H_5}^{OH}$ $0 \sim C_{c,H_5}^{OH}$ C_{c,H_5}^{OH} C_{c,H_5}^{O	C_2H_5 H CH_3 N $V_{C_0H_5}$ 1 C	C_2H_5 CH_3 H $N \sim C_{C_2H_3}^{OH}$ 2 C	$0C_2H_5$ CH_3 H $N \sim CH_5$ 0 C	C_6H_{11} CH_3 H $N \xrightarrow{OH} C_{GH_3}$ 0 C	$- \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ CH ₃ H N $\times \begin{pmatrix} 0 \\ C_{c,H_3} \end{pmatrix}$ 0 C	$-CH_{a}N$ CH_{a} CH_{a} H N CH_{c} 0 C	C_2H_5 H CH_3 0 C	C_2H_5 H H NICH ₃ CH(CH ₃)CH ₂ $\longrightarrow 0$ C	C_2H_5 H CH_3 N($CH_2CH=CH_2$), 0 C	^b Malcate. ^c Calcd.: HCl, 8.68. Found: HCl, 8.67. ^d Calcd.: HCl, 9.08. For
H C_2H_s CH_s H $\sqrt[N]{C_sH_s}$ 1 C	H C_2H_6 H CH_4 $N O_{G_6H_5}$ 1 C	H C_6H_5 CH_3 H $N \longrightarrow C_6H_6$ 0 C	H $C_{6}H_{5}$ H CH_{3} $N \longrightarrow C_{6}H_{5}$ 0 C	p -CH ₃ O C ₂ H, H CH ₃ $N \longrightarrow C_{c,H_3}$ 0 C	$h_{\rm e}$ CH ₃ O C ₂ H ₆ H CH ₃ O C $N \longrightarrow C_{\rm cH_5}$ O C	H C_6H_6 H C_6H_6 $N \longrightarrow 0$ C	H C_2H_5 H C_6H_5 $N \longrightarrow C_6H_5$ 0 C	H C_2H_6 H CH_3 $N \longrightarrow C_{GH_3}^{OH}$ 0 C	H C_2H_5 CH_3 H $N \sim C_{CH_5}^{OH}$ 0 $C_{CH_5}^{OH}$ 0 C_{C	H C_2H_5 H CH_3 $N \sim CH_6$ 1 C	H C_2H_5 CH_4 H $N \sim C_{C,H_5}^{OH}$ 2 C	H $0C_2H_4$ CH_3 H $N \longrightarrow C_{C_2H_4}^{OH}$ 0 C	$H \qquad C_6 H_{11} \qquad C H_3 H \qquad N \overbrace{C_6 H_4}^{OH} \qquad 0 C$	H $- \left< 0 \right>$ CH ₃ H N $\left< c_{G,H_3} \right>$ 0 C	H $-CH_{s}N$ $\rightarrow CH_{s}$ H N $\sim CH_{s}$ H CH_{s} O C	H C_2H_5 H CH_3 0 C	H C_3H_5 H H NICH, CH, CH, CH, CH, CH, OCH_3 0 C	H C_2H_5 H CH_3 $N(CH_2CH=CH_2)_2$ 0 C	alate. ¹ Maleate. ^c Calcd.: HCl, 8.68. Found: HCl, 8.67. ^d Calcd.: HCl, 9.08. Fo

Reduction of α -(4-hydroxy-4-phenylpiperidino)propionanilide (VII) with lithium aluminum hydride gave the corresponding amine (VIII) which showed near infrared absorption (in chloroform) at 1.47 (OH) and 1.56 μ (NH). Propionylation of VIII and formation of the oxalic salt gave IX.

The near-infrared absorption of IX (free base) at 1.47 μ confirmed the presence of an OH group. However, heating VIII in chloroform with propionyl chloride for 7 hr. resulted in dehydration to give X. Absorption at 240.8 m μ (ϵ 16,800)⁵ showed the presence of a styrene-like double bond.

Pharmacology.—The most active compound in this series is N-[1-methyl-2-(4-hydroxy-4-phenylpiperidino)ethyl]propionanilide oxalate (**27**) which is 150 times more active than morphine sulfate and about 1000 times more active than (+)-propoxyphene hydrochloride. These conclusions are based on the ED_{50} values determined by the artery-clip⁷ method after subcutaneous administration in mice. The structure– activity relationships in this series may be summarized as follows (referring to I).

$$R^1 = H > R^1 = OCH_3(o, m, p); n = 0 > 1 > 2$$

$$R^3 = CH_3, R^4 = H > R^3 = H, R^4 = CH_3$$

$$B = -N \xrightarrow{OH} > -N \xrightarrow{C_6H_5} > -N \xrightarrow{C_6H_5} >$$

Selected analgesic results are given in Table II. These compounds, though in different degrees, led to the development of dependence and were antagonized by nalorphine. For example, **27** produced less tolerance development and was less antagonized by nalorphine than was **16**. The abstinence symptoms in monkeys⁸ addicted to 3 mg./kg. of morphine sulfate were completely suppressed 12–14 hr. after withdrawal by 0.2 mg./kg. of **16**. Additional pharmacological details will be published elsewhere.

Experimental⁹

The detailed procedures are intended as models for the preparation of the compounds in Table I.

N-[2-(4-Phenylpiperidino)propyl]propionanilide Oxalate (III). Method A. (a) Synthesis of II.—1,2,3,6-Tetrahydro-4phenylpyridine (31.8 g., 0.2 mole), α -bromopropionanilide (45.0 g., 0.2 mole), 2-propanol (230 ml.), and anhydrous sodium carbonate (21.2 g., 0.2 mole) were heated under reflux for 15 hr. with vigorous stirring, then were filtered and cooled. The solid that separated was collected, washed with ether, and dried in air, m.p. 138–139°, yield 45.9 g. It was recrystallized from aqueous acetone to give a pale yellow powder (39 g.) of m.p. 139–140°.

TABLE II ANALGESIC ACTIVITY

	E D 50		
	$\langle 95\% m ~confidence$	$\mathrm{LD}_{\mathrm{au}}$	Thera-
	limits)	ba s e,	peutic
Compd.	Base, mg./kg., s.c. ^a	mg./kg., s.c.	index
16	0.20(0.14 - 0.29)	86.0^{b}	430.0
17	0.48(0.37-0.63)	251.0^{b}	556.0
26	0.155(0.076-0.32)	94.6^{5}	610.0
27	0.021(0.015 - 0.035)	89.5^{b}	4260.0
32	0.088(0.058 - 0.13)	71.0^{6}	807.0
34	0.045(0.028-0.065)	91.7^{h}	2040.0
30	$7^{+}5(5,35(10,5))$	45.5^b	6.1
18	$29.5(23.6{ ext{}}36.5)$	117.0^{b}	4.0
19	17.5(14.6-20.0)	54.2^{b}	3.10
1	35.5(23.4 - 54.0)	480.0 °	13.5
(+)-Propoxy-			
phene hydro-			
ahlorido ^d	99 5 (97 B. 18 4)	1.19. Oc	5 95

$chloride^{a}$	$22.5(27.6{ cdot}18.4)$	118.0°	5.25
Morphine sulfate	3.10(2.68 - 3.60)	390.0°	126.0

^a Determined by the method of J. F. Litchfield, Jr., and F. Wilcoxon, J. Pharmacol. Expll. Therap., **96**, 99 (1949), using at least four dose levels of 20 mice/dose. ^b Determined by the method of W. Thompson, Bacteriol. Rev., **11**, 115 (1947), with the tables of C. Brown, Federation Proc. **20**, 169 (1961), using a minimum of four dose levels of 2 mice/dose. ^c Same as footnote b except that 5 mice/dose were used. ^d Darvon^{*}.

Anal. Caled. for $C_{29}H_{22}N_2O$: N, 9.15. Found: N, 9.10.

(b) Reduction of II.—The above amide (34.5 g., 0.112 mole) in 150 ml. of tetrahydrofuran was added dropwise during 30 min. to a slurry of lithium aluminum hydride (6.4 g., 0.17 mole) in 150 ml. of tetrahydrofuran. The mixture was refluxed for 6 hr. and decomposed with 6 ml. of water, 6 ml. of 20% NaOH, and 18 ml. of water. The dried filtrate was concentrated *in vacuo* to a dark brown sirup which distilled to give 17.6 g. of 1-(2-anilinoethyl-1-methyl)-4-phenylpiperidine, b.p. 191-192° (0.3 mm.), $\lambda_{maxi}^{CHCl_2} 1.56 \mu$ (NH).

Anal. Calcd. for $C_{20}H_{24}N_2$: N, 9.60. Found: N, 9.54.

(c) **Propionylation.**—The above amine (14.9 g., 0.05 mole) and 30 ml. of propionic anhydride were kept at 25° for 2 days. The dark solution was concentrated *in vacuo* and the residue dissolved in ether. The cooled solution (ice-water bath) was shaken with aqueous NaOH. The ether layer was dried and treated with anhydrous oxalic acid (4.6 g., 0.05 mole) to give a tan salt (17.1 g.) which was recrystallized from methanol-ether to give 13.8 g. of colorless III, m.p. 185–186° dec., $\lambda_{max}^{CH_3OH}$ 257.5 m μ (ϵ 1050) and 264 m μ (ϵ 832).

Anal. Calcd. for $C_{23}H_{36}N_2O(C_2H_2O_4; N, 6.36)$. Found: N, 6.35.

The melting point of a mixture with a sample from method B (m.p. $186-187^{\circ}$ dec.) was not depressed. The ultraviolet spectra of two samples were identical.

Method B. (a) Synthesis of IV.—4-Phenylpiperidine (51.0 g., 0.316 mole), α -bromopropionanilide (72 g., 0.315 mole), absolute ethanol (250 ml.), and anhydrous sodium carbonate (33.5 g., 0.316 mole) were stirred under reflux for 22 hr. The mixture was filtered while hot and the inorganic salt was rinsed with absolute ethanol. The clear filtrate was heated to boiling, and water was added to cloudiness. A colorless solid separated and was collected by filtration, m.p. 117–120°, yield 77.9 g. (80%). A sample was recrystallized from aqueous methanol to give needles of m.p. 123–124°; $\lambda_{\max}^{CHCi_{2}}$ 2.88 (NH), 5.91 (amide C==O), and 6.5 μ (amide II).

Anal. Caled. for C₂₀H₂₄N₂O: N, 9.09. Found: N, 9.08.

(b) 1-(2-Anilinoethyl-1-methyl)-4-phenylpiperidine. $-\alpha$ -(4-Phenylpiperidino)propionanilide (76.0 g., 0.246 mole) was reduced with lithium aluminum hydride as before. The reaction mixture was processed by standard procedures to give a dark brown liquid, which distilled as a pale yellow liquid at 182-191° (0.15-0.2 mm.), yield 60.2 g. (83.3%).

(c) **Propionylation.**—The above amine (32.5 g., 0.11 mole) was propionylated as before and converted to the oxalate salt. The colorless salt was collected, washed with ether, and dried in air, m.p. 186° dec., yield 45.1 g. (92%). It was recrystallized from aqueous methanol-ether to give 39.7 g. of a colorless powder (17), m.p. 186~187° dec.

⁽⁷⁾ C. Bianchi and J. Franceschini, Brit. J. Pharmacol., 9, 280 (1954).

⁽⁸⁾ Private communication from G. Deneau and M. H. Seevers, Department of Pharmacology, School of Medicine, University of Michigan, Ann Arbor, Mich.

⁽⁹⁾ All melting points were measured in a Büchi apparatus (Switzerland) and are corrected. Infrared spectra were measured with a Perkin-Elmer Infracord Model 137 and Perkin-Elmer 237 grating infrared spectrophotometer. Near infrared and ultraviolet spectra were measured with a Warren Spectracord 4000. Titrations were done with a Sargent recording titrator Model D.

1-(2-Anilinopropyl)-1,2,3,6-tetrahydro-4-phenylpyridine.— The reaction mixture (0.2-mole run) in benzene obtained from α -bromopropionyl chloride and 4-phenyl-1,2,3,6-tetrahydropyridine under Schotten-Baumann conditions was stirred with 0.4 mole of aniline under reflux for 24 hr. Aniline hydrobromide was filtered, washed with benzene, and the dark benzene solution was concentrated *in vacuo* to give a tan solid mass of amide V which, after trituration with water and ether, melted at 120– 127°, yield 40.5 g. (66%).

The amide (74.4 g., 0.24 mole) was reduced with lithium aluminum hydride under conditions identical with those for the reduction of II. The dark brown liquid distilled as a pale yellow viscous liquid of b.p. $192-201^{\circ}$ (0.4 mm.), yield 47.9 g. (67.4%).

 \tilde{N} -[1-Methyl-2-(1,2,3,6-tetrahydro-4-phenyl-1-pyridyl)ethyl]propionanilide Oxalate (VI).—The above amine (47.9 g., 0.164 mole) was propionylated as usual. Anhydrous oxalic acid (15.3 g., 0.17 mole) was added to the crude amide to give a light tan sticky gum which slowly solidified on scratching. The solid was collected, washed with ether, and dried in air, m.p. 150–156°, yield 60.1 g. (83.5%). It was recrystallized twice from methanol-ether (Norit) to give 39.1 g. of a colorless powder (14) of m.p. 159–160° dec., $\lambda_{max}^{CH_3OH}$ 244.5 m μ (ϵ 14,600). Mixed with 16 (m.p. 152–153°), the m.p. was 141–151° (with bubbling).

1-(2-Anilino-1-methylethyl)-4-hydroxy-4-phenylpiperidine (VIII).- $-\alpha$ -(4-Hydroxy-4-phenylpiperidino)propionanilide (VII, 45.0 g., 0.14 mole) was reduced with lithium aluminum hydride as before. The amber sirup distilled at 202-215° (0.25 mm.); yield 30.3 g.; $\lambda_{max}^{CHCl_2}$ 2.79 (OH), 2.99 (NH), and 8.59 μ (CO stretching of tertiary alcohol), but no carbonyl band around 6 μ . The near infrared spectrum (CHCl₃) showed bands at 1.47 and 1.56 μ due to OH and NH, respectively.

Anal. Caled. for $C_{20}H_{25}N_2O$: N (basic), 4.52. Found: N (basic), 4.51 (titration).

N-[2-(4-Hydroxy-4-phenylpiperidino)propyl]propionanilide (IX).—Amine VIII (30.0 g.) was dissolved in 50 ml. of hot benzene, and propionic anhydride (25 ml.) was added. The light amber solution was refluxed for 1 hr., and the solvent was removed *in vacuo*. The residue was taken up in ether, washed with sodium hydroxide solution, and dried (MgSO₄). The filtrate was treated with 8.7 g. (0.1 mole) of anhydrous oxalic acid to give a light tan sticky gum which gradually solidified on scratching. This was collected by suction, washed with ether, and airdried, yield 43.9 g. It was recrystallized from methanol-ether to give a solid which was collected, washed with ether, and dried *in vacuo*, m.p. 103-105° dec., yield 31.7 g. (26).

A small sample was converted to the free base, extracted with ether, and dried. Removal of the solvent *in vacuo* gave a sirup whose near infrared spectrum (CHCl₃) showed an OH band at 1.47 μ , but no NH band at 1.56 μ .

N-[2-(1,2,3,6-tetrahydro-4-phenyl-1-pyridyl)propyl]propionanilide Oxalate (X).—Amine VIII (28.7 g., 0.09 mole), propionylchloride (37.0 g., 0.4 mole), and 100 ml. of chloroform were refluxed for 1.5 hr. and left overnight. The solvent and excess acid chloride were removed *in vacuo* to leave an amber sirup which was dissolved in a small amount of methanol and made basic with sodium hydroxide solution. The free base was extracted with ether and dried (MgSO₄). The solvent was removed *in vacuo* to give a sirup. The infrared spectrum (CHCl₃) showed an ester band at 5.79 (m) and amide band at 6.10μ (s). Esterification seemed to be incomplete. It was again dissolved in 100 ml. of chloroform, and 15 ml. of propionyl chloride was added. The solution was refluxed for 5.5 hr. and left overnight. The reaction mixture was worked up as before. The infrared spectrum of the free base (CHCl₃) showed a weak ester band at 5.80, a strong amide band at 6.10μ , but no OH band. The sirup was dissolved in ether, filtered, and treated with anhydrous oxalic acid (9.0 g., 0.1 mole) to give a sticky gum which soon solidified on scratching, yield 37.3 g. It was recrystallized from methanolether to give 19.7 g. of a colorless solid (15), m.p. 193.5–194.5° dec., λ_{mowB}^{mox} 240.8 m μ (\$16,800).

dec., λ_{\max}^{MaOH} 240.8 mµ (ϵ 16,800). **N-[1-Methyl-2-(4-hydroxy-4-phenylpiperidino)ethyl]pro pionanilide (27). A.** 1-(α -Anilinopropionyl)-4-hydroxy-4-phenlypiperidine.—A toluene solution of α -bromopropionyl chloride (68.6 g., 0.4 mole) was added slowly with stirring to an ice-cold suspension of 4-phenyl-4-piperidinol (70.8 g., 0.4 mole) in 750 ml. of toluene and 130 ml. of 20% sodium hydroxide. Stirring was continued overnight. The layers were separated, and the dried toluene layer was concentrated *in vacuo* to half its volume. This concentrate and aniline (74.4 g., 0.8 mole) were heated under reflux with stirring for 16 hr. The solvent was removed *in vacuo* to leave a solid mixture of amide and aniline hydrobromide. This was suspended in water and left overnight. The amide was collected, washed with water, and dried. The solid was triturated in ether and again collected and dried, yield 112.9 g. (87%), m.p. 147.5–148.5°.

Anal. Calcd. for $C_{20}H_{24}N_2O_2$: N, 8.64. Found: N, 8.65.

B. 1-(2-Anilinopropy)-4-hydroxy-4-phenylpiperidine.— The above amide (113 g., 0.35 mole) was reduced with lithium aluminum hydride as before. The reduction was repeated since the infrared spectrum still showed absorption at 6.1 μ . The crude amine was distilled using a short-path column, b.p. 195-210° (0.2-0.3 mm.), yield 39.5 g. (36.6%). No amide absorption (6.1 μ) appeared in the spectrum of this product.

Anal. Calcd. for C₂₀H₂₆N₂O: N, 9.03. Found: N, 9.04.

C. N-2-(4-Hydroxy-4-phenylpiperidino)-1-methylethyl]propionanilide Oxalate.—A solution of the above compound (39 g., 0.125 mole) was treated with propionic anhydride as before and converted to the oxalate salt. The salt was collected and recrystallized twice from a methanol-ether solution, yield 41 g. (72%), m.p. 165.5–167.0° dec.

Acknowledgment.—The authors wish to express their gratitude to T. J. Leipzig and R. A. Kulp for the preparation of starting materials and to F. R. Bunn and Mrs. E. Kurchacova for analytical and spectral data.