

[Chem. Pharm. Bull.]
26(5)1615-1619(1978)

UDC 547.913.04 : 581.192

Conversion of Grayanotoxin III to Grayanotoxin V, Rhodojaponin III and Rhodojaponin IV

TADAMASA TERAI, MASAOKI KATAI,^{1a)} NOBUYUKI HAMANAKA,^{1b)}
TAKESHI MATSUMOTO,^{1c)} and HARUO MEGURI^{1a)}Department of Applied Chemistry, Osaka Institute of Technology^{1a)}
and Department of Chemistry, Faculty of Science,
Hokkaido University^{1c)}

(Received October 17, 1977)

Treatment of grayanotoxin (G) III with CH_3CHO in the presence of anhydrous CuSO_4 afforded a diethylidene compound (2). Oxidation of 2 (CrO_3) and subsequent ozonolysis and hydrolysis (KOH-MeOH) gave G-V (4) in 34% overall yield. Direct ozonolysis of 2 produced rhodojaponin (R) IV (8) in 68% yield. Δ^2 -Diethylidene compound (10) was oxidized to 11, which on ozonolysis-hydrolysis furnished R-III (12) in 9.2% overall yield.

Keywords—*Leucothoe grayana* MAX.; *Rhododendron japonicum* SURINGER; grayanotoxin V; rhodojaponin III; rhodojaponin IV; ozonolysis; conversion of grayanotoxin III; selective masking and demasking of grayanoids

From the leaves of *Leucothoe grayana* MAX. (Ericaceae), seventeen A-nor-B-homokauranoids, grayanotoxin I—XVII have been hitherto isolated.²⁾

On the other hand, from the flowers of *Rhododendron japonicum* SURINGER, seven toxic diterpenoids with the same skeleton, rhodojaponin I—VII have been isolated.³⁾ These grayanoids are generally highly oxygenated, bearing several hydroxyl or acetoxyl groups extending over the A, B, C and D rings, as exemplified by grayanotoxin III (G-III) (1). More highly oxygenated grayanoids contain a carbonyl group [grayanotoxin V (G-V) (4)] or an epoxide group [rhodojaponin III (R-III) (12)]. Structures of most of the grayanoids have been determined by correlating them chemically with the basic grayanoids, grayanotoxin I and grayanotoxin II.⁴⁾ However, interconversion among the grayanoids is not always simple, since they are acid sensitive, and selective masking and demasking reactions at a particular position are often difficult.

In the present paper, we should like to describe a selective method for masking and demasking of oxygen functions of grayanoids under mild conditions and conversion of G-III to G-V, R-III and rhodojaponin IV (R-IV) by means of the new procedure.

- 1) a) Location: Omiya 5-16-1, Asahi-ku, Osaka 535, Japan; b) Present address: Research Institute, Ono Pharmaceutical Co., Ltd., Sakurai 3-1-1, Shimamoto-cho, Mishima-gun, Osaka 618, Japan; c) Location: Kita 10 Nishi 8, Sapporo 060, Japan.
- 2) S. Takei and J. Miyajima, *Nippon Nogeikagaku Kaishi*, **10**, 1093 (1934); *idem, ibid.*, **12**, 497 (1935); T. Okuno, N. Hamanaka, H. Miyakoshi and T. Matsumoto, *Tetrahedron*, **26**, 4765 (1970); H. Hikino, S. Koriyama, T. Ohta, and T. Takemoto, *Chem. Pharm. Bull.* (Tokyo), **18**, 2357 (1970); *idem, ibid.*, **19**, 1289 (1971); *idem, ibid.*, **20**, 422 (1972); H. Hamanaka, H. Miyakoshi, A. Frusaki and T. Matsumoto, *Chemistry Letters*, **1972**, 779; S. Gasa, R. Ikeda, N. Hamanaka, and T. Matsumoto, *Bull. Chem. Soc. Jpn.*, **49**, 835 (1976).
- 3) H. Hikino, T. Ohta, Y. Hikino, and T. Takemoto, *Chem. Pharm. Bull.* (Tokyo), **17**, 1078 (1969); *idem, ibid.*, **18**, 2357 (1970); *idem, ibid.*, **20**, 1090 (1972); R. Iryye and I. Tomida, *Tetrahedron Lett.*, **1972**, 1381.
- 4) a) Structure: J. Iwasa, Z. Kumazawa and M. Nakajima, *Agr. Biol. Chem.* (Tokyo), **25**, 782 (1961); *idem, ibid.*, **25**, 793 (1961); *idem, ibid.*, **25**, 798 (1961); H. Kakisawa, *Nippon Kagaku Zasshi*, **82**, 1096 (1961); *idem, ibid.*, **82**, 1216 (1961); H. Kakisawa, T. Kozima, M. Yanai and K. Nakanishi, *Tetrahedron*, **21**, 3019 (1965); b) Synthesis: N. Hamanaka and T. Matsumoto, *Tetrahedron Lett.*, **1972**, 3087; S. Gasa, N. Hamanaka, S. Matsunaga, T. Okuno, N. Takeda, and T. Matsumoto, *ibid.*, **1976**, 553.

olysis of **3** and **5** in accordance with the procedure proposed by P. Deslongchamps.⁵⁾ The 3-dehydro-diethylidene derivative (**3**) and 3-dehydro-monoethylidene derivative (**5**) were reacted with ozone to give the already known 6,14-diacetate and 6-monoacetate as oily products respectively. These products gave **4** by hydrolysis with 5% KOH-MeOH. Melting points, IR spectrum and NMR data of **4** were identical with those of natural G-V.

Similarly, hydrolysis of diethylidene G-III (**2**) with dil. methanolic HCl at room temperature afforded selectively 5,6-monoethylidene G-III (**6**), δ (CDCl₃) 4.15 (C-3H), 4.79 (C-6H). The IR spectrum and other physical data of **6** were identical with those of the compound derived from G-1. However, removal of the second ethylidene group under acidic conditions was also difficult in this case. On the contrary ozonolysis of diethylidene G-III (**2**) afforded smoothly diacetate (**8**), mp 221–227°, in 68% yield. The IR and NMR spectra indicated the presence of two acetoxy groups, 1733 and 1240 cm⁻¹, δ (C₅D₅N) 2.01 and 2.30 (each 3H, singlet), which were allocated at C-6 and C-14 on the basis of the characteristic splitting pattern of C-6 and C-14 protons at δ 5.19 and 6.10.

Diacetate (**8**) was identified with an authentic sample of R-IV by the use of thin-layer chromatography (TLC), IR and NMR spectroscopy.

On the other hand, monotosylate (**9**), obtained from **2**, was refluxed with collidine to give Δ^2 -3-deoxy-diethylidene G-III (**10**) in 76% yield, δ (C₅D₅N) 5.54 and 6.15 (each 1H, d.d, $J_{1,2}$ =3.0 Hz, $J_{2,3}$ =6.0 Hz and $J_{3,1}$ =2.0 Hz), which, on epoxidation with *m*-chloroperbenzoic acid afforded stereoselectively 2,3-epoxy-3-deoxy-diethylidene G-III (**11**). Ozonolysis and subsequent hydrolysis of **11** in the same way as mentioned above furnished **12** in an overall yield of 9.2% from G-III. The product was identical with an authentic sample of R-III in TLC, IR and NMR spectra.

Experimental

All melting points are uncorrected. The NMR spectra were obtained on a Hitachi R-22 spectrometer in CDCl₃ or C₅D₅N containing tetramethylsilane (TMS) as internal standard. Chemical shifts are reported as parts per million on the δ scale (s=singlet, d=doublet, t=triplet, q=quartet and m=multiplet). The IR spectra were measured on a Shimadzu IR-400 spectrophotometer.

5,6:14,16-O-Diethylidene G-III (2)—To a solution of 1 g of **1** in 30 ml acetaldehyde, 2 g of anhydrous cupric sulfate was added and the solution was stirred for 1 hr at room temperature. The mixture was filtered and the filtrate was evaporated under reduced pressure. The residue was crystallized from ether-*n*-hexane (1:1) to give 840 mg of **2** as colorless prisms, mp 180–181°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3480 and 3470 (OH), 1139, 1082 and 1000 cm⁻¹, NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.85, 1.07, 1.33 and 1.42 (each 3H, s), 1.30, 1.35 (each 3H, d, J =5.0 Hz, >CH-CH₃), 3.62 (1H, br.s, C-3H), 4.23 (1H, br.s, C-6H), 4.48 (1H, s, C-14H), 4.98 and 5.12 (each 1H, q, J =5.0 Hz, >CH-CH₃). Anal. Calcd. for C₂₄H₃₈O₆: C, 68.22; H, 9.07. Found: C, 68.37; H, 9.04.

5,6:14,16-O-Diethylidene-3-dehydro G-III (3)—To a mixture of pyridine (10 ml) and CrO₃ (500 mg), 500 mg of **2** dissolved in 5 ml pyridine was added and allowed to stand overnight. The mixture was poured into ice water and the resultant solution was shaken 3 times with EtOAc. The combined EtOAc layers were washed with water, dried over anhydrous Na₂SO₄ and evaporated *in vacuo* to give crystals. Recrystallization from ether-*n*-hexane gave 250 mg of **3** as colorless prisms, mp 251–252°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3513 (OH), 1733 (C=O), 1095–1140 cm⁻¹, NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.93, 1.00, 1.25 and 1.34 (each 3H, s), 1.24 and 1.30 (each 3H, d, J =4.8 Hz, >CH-CH₃), 4.26–4.30 (1H, m, C-6H), 4.48 (1H, s, C-14H), 4.90 and 5.13 (each 1H, q, J =4.8 Hz, >CH-CH₃), MS m/e : 420 (M⁺). Anal. Calcd. for C₂₄H₃₆O₆: C, 68.54; H, 8.63. Found: C, 68.58; H, 8.62.

3-Dehydro G-III (=G-V) (4)—Air containing ozone was passed through an ice cooled solution of 50 mg of **3** in 10 ml of EtOAc-AcOH (1:1). After the absorption of ozone had ceased, the solution was treated with 3 ml of water and allowed to stand overnight. The solvent was removed *in vacuo*, and the residue containing diacetyl G-V was tried to crystallize from several solvents without success. After the reaction mixture was treated with 5% KOH-MeOH for 4 hr at 50°, the solution was evaporated under reduced pressure and the residue was extracted with EtOAc. The combined EtOAc layers were washed with water, dried over Na₂SO₄, and evaporated *in vacuo* to give crystals. Recrystallization from EtOAc gave 30 mg of **4** as colorless needles, mp 253–256°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3550–3250 (OH), 1730 (C=O) cm⁻¹, NMR $\delta_{\text{ppm}}^{\text{C}_5\text{D}_5\text{N}}$: 1.16, 1.47, 1.53 and 1.73 (each 3H, s), 4.68 (1H, m, C-6H) and 5.07 (1H, s, C-14H). The IR spectrum and NMR data of **4** were identical with those of natural G-V.

5,6-O-Ethylidene-3-dehydro G-III (5)—To a solution of 100 mg of **3** in 10 ml of MeOH, 0.1 ml of conc. HCl was added and the mixture was allowed to stand at room temperature for 1 hr. The solution

was neutralized with 10% Na_2CO_3 , concentrated *in vacuo* and extracted 3 times with EtOAc. The combined EtOAc layers were washed with water, dried over Na_2SO_4 and evaporated *in vacuo* to give crystals. Recrystallization from EtOAc gave 70 mg of **5** as colorless needles, mp 232–238°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3460 (OH), 3330 (OH), 1734 (C=O) cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.98, 1.06 (each 3H, s), 1.40 (6H, s), 1.34 (3H, d, $J=5.0$ Hz, >CH-CH_3), 4.20 (1H, br. s, C-6H), 4.40 (1H, br. s, C-14H) and 4.48 (1H, q, $J=5.0$ Hz, >CH-CH_3). *Anal.* Calcd. for $\text{C}_{22}\text{H}_{34}\text{O}_6$: C, 66.98; H, 8.69. Found: C, 66.50; H, 8.68.

5,6-O-Ethylidene G-III (6)—To a solution of 100 mg of **2** in 10 ml of MeOH, 0.1 ml of conc. HCl was added and the resultant solution was allowed to stand at room temperature for 2 hr. The mixture was treated by the same method as above. Recrystallization from EtOAc-*n*-hexane gave 50 mg of **6** as amorphous solid, IR $\nu_{\text{max}}^{\text{KBr}}$: 3600–3200 (OH), 1138, 1100, 1060 and 1020 cm^{-1} . The IR spectrum of **6** was identical with that of the derivative from G-I.

5,6-O-Ethylidene G-III 3,14-Diacetate—To a solution of 100 mg of **6** in 1 ml of pyridine, 1 ml of Ac_2O was added and allowed to stand at room temperature overnight. The mixture was poured into ice water and extracted 3 times with EtOAc. The combined EtOAc layers were washed with water, dried over Na_2SO_4 and evaporated *in vacuo* to give crystals. Recrystallization from EtOAc-*n*-hexane gave 90 mg of 5,6-O-ethylidene G-III 3,14-diacetate, mp 152–155°, IR $\nu_{\text{max}}^{\text{KBr}}$: 3600–3250 (OH), 1720 (C=O) and 1245 (C-O-) cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.93 and 0.95 (each 3H, s), 1.37 (6H, s), 1.35 (3H, d, $J=5.0$ Hz, >CH-CH_3), 2.40 (6H, s, $\text{OAc} \times 2$), 4.15 (1H, m, C-3H), 4.79 (1H, m, C-6H), 4.93 (1H, q, $J=5.0$ Hz, >CH-CH_3) and 5.83 (1H, s, C-14H).

G-III 6,14-Diacetate (=R-IV) (8)—Air containing ozone was passed through an ice cooled solution of 200 mg of **2** in 10 ml of EtOAc-AcOH (1:1). After the absorption of ozone had ceased, the solution was treated with 3 ml of water and left overnight. The mixture was neutralized with 10% NaHCO_3 , concentrated *in vacuo* and extracted 3 times with EtOAc. The combined EtOAc layers were washed with water, dried over anhydrous Na_2SO_4 and evaporated *in vacuo* to give crystals. Recrystallization from EtOAc-*n*-hexane gave 150 mg of **8** as colorless needles, mp 221–227°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3578, 3342 (OH), 1733 (C=O) and 1240 (C-O-) cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.96 and 1.70 (each 3H, s), 1.43 (6H, s), 2.01 and 2.30 (each 3H, s, $\text{-OAc} \times 2$), 3.80 (1H, m, C-3H), 5.19 (1H, m, C-6H) and 6.10 (1H, s, C-14H). *Anal.* Calcd. for $\text{C}_{24}\text{H}_{38}\text{O}_8 \cdot 1/2\text{H}_2\text{O}$: C, 62.18; H, 8.48. Found: C, 62.20; H, 8.72.

The IR spectrum and NMR data of **8** were identical with those of natural R-IV. Hydrolysis of **8** with 5% KOH gave G-III. **8** was identical with G-III 6,14-diacetate which were obtained from G-III on partial acetylation with Ac_2O -pyridine.

3-O-Tosyl-5,6:14,16-O-diethylidene G-III (9)—To a solution of 1 g of **2** in 10 ml of pyridine, 1.1 g of *p*-toluenesulfonyl chloride was added and the solution was allowed to stand at 0° for 5 day. The mixture was poured into ice water and the resultant precipitate was washed with water, dissolved in EtOAc, and the organic layer was dried over anhydrous Na_2SO_4 and evaporated *in vacuo* to give crystals. Recrystallization from EtOAc-*n*-hexane gave 950 mg of **9** as colorless needles, mp 162–165°, IR $\nu_{\text{max}}^{\text{Nujol}}$: 3480 (OH), 1598 and 1180–1085 cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.86 and 0.96 (each 3H, s), 1.33 (6H, s), 2.45 (3H, s, arom. -Me), 1.28 and 1.40 (each 3H, d, $J=5.0$ Hz, >CH-CH_3), 4.13 (1H, m, C-3H), 4.42 (2H, m, C-6H, C-14H), 5.00 and 5.40 (each 1H, q, $J=5.0$ Hz, >CH-CH_3), 7.34 and 7.82 (each 2H, d, $J=8.2$ Hz, arom. protons). *Anal.* Calcd. for $\text{C}_{31}\text{H}_{44}\text{O}_5\text{S}$: C, 64.56; H, 7.69. Found: C, 64.87; H, 7.80.

5,6:14,16-O-Diethylidene- Δ^2 -3-deoxy G-III (10)—A solution of 300 mg of **9** in 30 ml of collidine was refluxed for 4 hr. After cooling it was poured into dilute H_2SO_4 and extracted with EtOAc. The combined EtOAc layers were washed with water, dried over anhydrous Na_2SO_4 and evaporated *in vacuo* to give crystalline material. Recrystallization from EtOAc-*n*-hexane gave 160 mg of **10** as colorless needles, mp 179–182°, IR $\nu_{\text{max}}^{\text{KBr}}$: 3520 (OH), 1160–1085 cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.80, 1.08, 1.36 and 1.65 (each 3H, s), 1.38 and 1.40 (each 3H, d, $J=5.0$ Hz, >CH-CH_3), 4.38 (1H, m, C-6H), 5.54 (1H, d.d, $J_{2,1}=3.0$ Hz, $J_{2,3}=6.0$ Hz, C-2H), 6.15 (1H, d.d, $J_{3,1}=2.0$ Hz, $J_{3,2}=6.0$ Hz, C-3H), 5.11 (1H, s, C-14H), 5.00 and 5.40 (each 1H, q, $J=5.0$ Hz, >CH-CH_3). *Anal.* Calcd. for $\text{C}_{24}\text{H}_{36}\text{O}_5$: C, 71.25; H, 8.97. Found: C, 71.43; H, 9.19.

2,3-Epoxy-3-deoxy-5,6:14,16-O-diethylidene G-III (11)—To a solution of 110 mg of **10** in 10 ml of EtOAc, a mixture of 86 mg of *m*-chloroperbenzoic acid and 2 ml of methylene chloride was added and the whole was allowed to stand overnight at room temperature. After the mixture was treated with 10% aq. Na_2S (10 ml), the organic layer was washed with 5% NaHCO_3 and water, dried over anhydrous Na_2SO_4 and evaporated *in vacuo* to give crystalline material. Recrystallization from CHCl_3 -*n*-hexane gave 60 mg of **11** as colorless needles, mp 209–210°, IR $\nu_{\text{max}}^{\text{KBr}}$: 3520 (OH), 1080–1140 cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 0.88, 1.18, 1.30 and 1.53 (each 3H, s), 1.27 and 1.34 (each 3H, d, $J=5.0$ Hz, >CH-CH_3), 3.05 and 3.58 (each 1H, d, $J=4.0$ Hz, C-2H, C-3H), 4.25 (1H, m, C-6H), 4.53 (1H, s, C-14H), 4.86 and 5.12 (each 1H, q, $J=5.0$ Hz, >CH-CH_3). *Anal.* Calcd. for $\text{C}_{24}\text{H}_{36}\text{O}_6$: C, 68.54; H, 8.63. Found: C, 68.35; H, 8.97.

2,3-Epoxy-3-deoxy G-III (=R-III) (12)—Air containing ozone was passed through an ice cooled solution of 50 mg of **11** in 4 ml of AcOH-EtOAc (1:1). After the ozonization, the solution was treated with 3 ml of water and left overnight. The mixture was neutralized with 10% NaHCO_3 , concentrated *in vacuo* and extracted 3 times with EtOAc. The usual working up of the extract gave an amorphous material. To a solution of the product in 2 ml of MeOH, 400 mg of anion exchange resin (IRA 400) was added and the mixture was refluxed for 1 hr. The resin was filtered off and the filtrate was evaporated under reduced

pressure to give a crystalline solid. Recrystallization from EtOAc gave 20 mg of **12** as colorless needles, mp 285–287°, IR $\nu_{\text{max}}^{\text{KBr}}$: 3200–3600 (OH), 855 cm^{-1} , NMR $\delta_{\text{ppm}}^{\text{CDCl}_3}$: 1.21, 1.47, 1.53 and 1.84 (each 3H, s), 3.18 and 4.15 (each 1H, d, $J=3.0$ Hz, C-2H, C-3H), 4.42 (1H, m, C-6H) and 4.83 (1H, s, C-14H). *Anal.* Calcd. for $\text{C}_{20}\text{H}_{32}\text{O}_6$: C, 65.19; H, 8.75. Found: C, 64.49; H, 9.00.

The IR spectrum and NMR data of **12** were identical with those of natural R-III.

[Chem. Pharm. Bull.]
26(5)1619–1622 (1978)

UDC 547.918.02.04 : 581.192

Chemische Untersuchungen der Inhaltsstoffe von *Pteris vittata* L.¹⁾

TOSHIKO SATAKE, TAKAO MURAKAMI,^{2a)} YASUHISA SAIKI,^{2b)}
und CHIU-MING CHEN^{2c)}

Pharmazeutisches Institut, Naturwissenschaftliche Universität Tokyo,^{2a)}
Department of Pharmaceutical Sciences, Kobe Gakuin University^{2b)}
und Department of Chemistry, National Tsing Hua University^{2c)}

(Eingegangen am 20. Oktober 1977)

Aus den oberirdischen Teilen von *Pteris vittata* L. wurden zwei neue Lignanglykoside, d.h., *cis*-Dihydro-dehydro-diconiferylalkohol-9-O- β -D-Glukosid (I) und Lariciresinol-9-O- β -D-Glukosid (III) isoliert und identifiziert.

Keywords—*Pteris vittata*; Pteridaceae; lignans; glucosides; dihydro-dehydro-diconiferylalcohol; lariciresinol; DCCC; NMR; chemotaxonomy

In Fortsetzung der chemischen und chemotaxonomischen Untersuchungen der Gattung *Pteris* und der verwandten Gattungen wurde *Pteris vittata* L. (jap. Name: Moejimashida, Fundort: Okinoerabu/Kagoshima-Präfektur, Sammelzeit: August, 1975) auf die Inhaltsstoffe untersucht und aus den oberirdischen Teilen zwei neue Lignan-Glykoside isoliert.

Glykosid A (I) wurde als amorphes Pulver vom $[\alpha]_{\text{D}}^{25} -23.6^\circ$ ($c=0.98$, MeOH) erhalten. Enzymatische Hydrolyse ergab D-Glukose und ein amorphes Aglykon (II), $\text{C}_{20}\text{H}_{24}\text{O}_6$ (M^+ 360.1596) vom $[\alpha]_{\text{D}}^{25} -8.5^\circ$ ($c=0.96$, Aceton). Die spektroskopischen Daten des Aglykons (II) sind nur mit der Struktur eines Dihydro-dehydro-diconiferylalkohols vereinbar. *rac*-Dehydro-diconiferylalkohol ist als Modellverbindung³⁾ bei der Lignin-Biosynthese bekannt.

Für die optisch aktiven Dihydro-dehydro-diconiferylalkoholen sind zwei Stereoisomeren beschrieben, d.h., eins ist derjenige vom Schmp. 102–103° und $[\alpha]_{\text{D}}^{25} +5.4^\circ$ ($c=2.0$, Aceton), der Weinges und Mitarbb.⁴⁾ durch Hydrierung des natürlichen Dehydro-diconiferylalkohols erhielten und ein anderer derjenige vom $[\alpha]_{\text{D}}^{18} -4.9^\circ$ ($c=0.91$, absol. Aceton), der Takemoto und Mitarbb.⁵⁾ durch alkalische Hydrolyse des natürlichen Diferulasäureester des Dihydro-dehydro-diconiferylalkohols erhielten. II erwies sich als identisch mit dem letzteren Isomeren durch Vergleich von Infrarot (IR)- und Kernmagnetische Resonanz (NMR-) Spektren.

Im NMR-Spektrum des Aglykon-Acetats (II') erscheinen die Signale von Protonen der Acetoxy-tragenden Methylengruppen bei 4.07 (2H am C(9'), t, $J=7$ Hz) und 4.31 (2H am C(9),

- 1) Chemische und chemotaxonomische Untersuchungen der Gattung *Pteris* und der verwandten Gattungen (Pteridaceae) IXX. Mitteil., XVIII. Mitteil.: N. Tanaka, M. Kudo, T. Taniguchi, T. Murakami, Y. Saiki, und C.-M. Chen, *Chem. Pharm. Bull.* (Tokyo), **26**, 1339 (1978).
- 2) Adresse: a) Funakawara-Machi, Shinjuku-Ku, Tokyo, 162, Japan; b) Arise, Igawatani-Machi, Tarumi-Ku, Kobe, 673, Japan; c) Kuang Fu Road, Hsinchu, Taiwan, China.
- 3) K. Freudenberg, *Nature*, **183**, 1152 (1959).
- 4) K. Weinges, R. Müller, P. Kloss, und H. Jaggy, *Ann. Chem.*, **736**, 170 (1970).
- 5) T. Takemoto, T. Momose, und G. Kusano, *Phytochemistry*, **14**, 1890 (1975).