



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

A CONVENIENT AND EFFECTIVE METHOD FOR THE SYNTHESIS OF TETRAOXAQUATERENES

Robert Pajewski ^a, Ryszard Ostaszewski ^a & Janusz Jurczak ^b

^a Institute of Organic Chemistry, Polish Academy of Sciences, Kasprzaka, 44, 01-224, Warsaw

^b Department of Chemistry, Warsaw University, Pasteura 1, 02-093, Warsaw, Poland

Published online: 11 Feb 2009.

To cite this article: Robert Pajewski, Ryszard Ostaszewski & Janusz Jurczak (2000) A CONVENIENT AND EFFECTIVE METHOD FOR THE SYNTHESIS OF TETRAOXAQUATERENES, *Organic Preparations and Procedures International: The New Journal for Organic Synthesis*, 32:4, 394-397, DOI: [10.1080/00304940009355943](https://doi.org/10.1080/00304940009355943)

To link to this article: <http://dx.doi.org/10.1080/00304940009355943>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

3. G. W. Kabalka, R. D. Pace and P. P. Wadgaonkar, *Synth. Commun.*, **20**, 2453 (1990).
4. T. W. Green and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Wiley, New York, 214 (1991) and references cited therein.
5. (a) D. P. Curran, J. F. Brill and D. M. Rakiewicz, *J. Org. Chem.*, **49**, 1654 (1984); (b) P. Vankar, R. Rathore, and S. Chandrasekaran, *J. Org. Chem.*, **51**, 3063 (1986); (c) B. C. Ranu and D. C. Sarkar, *J. Org. Chem.*, **53**, 878 (1988); (d) A. Kamal, M. V. Rao, and H. M. Meshram, *J. Chem. Soc. Perkin Trans. 1*, 2056 (1991); (e) H. Kumar, Kaur, B. and B. Kumar, *Indian. J. Chem.*, **30B**, 869 (1991); (f) M. Curini, M. Constantino O. Rosati and E. Pisani, *Synlett*, 333 (1996); (g) R. S. Varma and H. M. Meshram, *Tetrahedron Lett.*, **38**, 5427 (1997); (h) A. S. Demir, C. Tanyeli and E. Altinel, *Ibid.*, **41**, 7267 (1997).
6. (a) G. A. Olah, Q. Liao, C. S. Lee and G. K. Surya Prakash, *Synlett* 427 (1993); (b) R. S. Varma, R. Dahiya and R. K. Saini, *Tetrahedron Lett.*, **38**, 8819 (1997); (c) J. Drabowicz, *Synthesis*, 1980 125; (d) Y. Yang, T. Li, Y. Li, *Synth. Commun.*, **23**, 1121 (1993); (e) N. B. Barhate, A. S. Gajare, R. D. Wakharkar, and A. Sudalai, *Tetrahedron Lett.*, **38**, 653 (1997).
7. I. Yavari and A. Shaabani, *J. Chem. Res. (S)*, 274 (1994).
8. J. March, *Advanced Organic Chemistry*, 3rd Ed., John Wiley and Sons (1985).
9. (a) A. I. Vogel, *A Text Book of Practical Organic Chemistry*, 3rd Ed., Longman Group (1975).
(b) *Dictionary of Organic Compounds*, 6th Ed., Chapman and Hall, London, (1995).

A CONVENIENT AND EFFECTIVE METHOD FOR THE SYNTHESIS OF TETRAOXAQUATERENES

Submitted by Robert Pajewski^a, Ryszard Ostaszewski,^a and Janusz Jurczak^{a,b*}
(04/25/00)

^a*Institute of Organic Chemistry, Polish Academy of Sciences,
Kasprzaka 44, 01-224 Warsaw*

^b*Department of Chemistry, Warsaw University,
Pasteura 1, 02-093 Warsaw, Poland*

There is a great interest in the synthesis of new types of ionophores; among others calixarenes^{1,2} and cyclodextrins³⁻⁵ are well known. Recent papers dealing with the synthesis of tetraoxaquaterenes show that this may be a very interesting scaffold in supramolecular chemistry. Among all quaterenes, the synthesis of octamethyltetraoxaquaterene (**3**) is documented very well. One of the best methods, already published by Chastrette *et al.*,^{6,7} based on the one-pot reaction of furan (**1**) with acetone (**2**), carried out in the presence of hydrochloric acid and lithium perchlorate, afforded the desired product **3** in 25% yield. Several attempts at modification of the reaction conditions were unsuccessful.^{8,9}

In this paper we present the effect of concentration of sulfuric acid on the yield of the cyclocondensation reaction of furan (1) with acetone (2) and with cyclohexanone (4), performed in dioxane (Scheme, Table).

We found that upon addition of concentrated sulfuric acid (Entry 9) to the mixture of furan (1) and acetone(2), the reaction mixture turned black (temperature of the reaction was kept below 10°C) and the crude product contained a large amount of polymeric material. After purification, the desired product 3 was obtained in a 20% yield.¹⁰ When the concentration of sulfuric acid was decreased to 93.3% (Entry 8), the product 3 was obtained in a 51% yield, and suppression of the polymerization reaction was observed. At a lower concentration of acid (Entry 7, 90.5%) the reaction yield was much higher (71% after recrystallization from toluene). Further experiments showed that the yield drops with decreased acid concentration (Entries 2-6), and below the acid concentration of 83%, no reaction occurred (Entry 1).




TABLE. Influence of the Concentration of Sulfuric Acid on the Yield of 3 and 5

Product 3			Product 5		
Entry	Conc. of H ₂ SO ₄	Yield	Entry	Conc. of H ₂ SO ₄	Yield
[%]	[%]	[%]		[%]	[%]
1	81.3	0	10	70.5	40
2	83.2	57	11	74.3	40
3	84.5	53	12	77.8	46
4	85.7	50	13	81.3	44
5	87.6	51	14	84.5	46
6	88.8	62	15	87.6	49
7	90.5	71	16	90.5	45
8	93.3	51	17	93.3	44
9	96.0	20	18	96	40

When acetone (**2**) was replaced with cyclohexanone (**4**), tetracyclohexyltetraoxaquaternene (**5**) was obtained under above-mentioned conditions (Scheme 1, Table 1). Previous synthetic routes for this compound were not very successful, and yields of 9%¹¹ and 16%⁹ were reported.

The highest yield was obtained for concentration of sulfuric acid 87.6% (Entry 15) similarly as in the case of the synthesis of tetraoxaquaterene **3** (Entry 7). Further experiments showed that the yield drops with decreased acid concentration (Entries 10-14) as well as with increased acid concentration (Entries 16-18).

The compound **5** precipitated from the reaction mixture, and after few attempts we managed to obtain the well-shaped monocrystals suitable for X-ray analysis, which finally confirmed the assumed structure.¹²

In summary, there is presented a one-step procedure for the synthesis of tetraoxaquaterenes **3** and **5**, characterized by relatively high yield and selectivity. A substantial effect of sulfuric acid concentration on the reaction yield is observed. An extension of this procedure to more complex systems which may be of interest as new scaffolds in supramolecular chemistry is now in progress.

EXPERIMENTAL SECTION

Melting points were determined using a Kofler hot-stage apparatus and are uncorrected. ¹H NMR spectra were recorded using a Varian Gemini 200BB (200 MHz) spectrometer. All chemical shifts are quoted in parts per million relative to tetramethylsilane (d, 0.00 ppm). IR spectra were obtained on Perkin-Elmer 1640 FTIR spectrophotometer as KBr pellets. Mass spectra were recorded on an AMD 604 Intecra instrument using the electron impact (EI) technique. Single-crystal X-ray diffraction analysis was performed on an Enraf-Nonius MACH 3 diffractometer.

Synthesis of Octamethyltetraoxaquaterene (3). Typical Procedure.- A precooled 90.5% sulfuric acid (5 mL) was added dropwise to a stirred precooled (0-5°) mixture of acetone (1.46 mL, 20 mmol), furan (1.44 mL, 20 mmol), and dioxane (30 mL). Upon addition of sulfuric acid, the reaction mixture turned red and some precipitation of product was observed. The reaction mixture was stirred overnight at room temperature, then water (50 mL) was added and the acid was neutralized by a sodium hydroxide aqueous solution (20%). Crude product was filtered off, washed with water (2 x 10 mL), methanol (2 x 10 mL) and recrystallized from toluene to give 1.54 g (71% yield) of colorless needles, m.p. 241.5-242° (*lit.*¹³ 243°); *R*_f=0.51 (hexane-ethyl acetate, 95:5 v/v). IR (KBr pellets): 2983, 2871, 1651, 1602, 1559, 1449, 1365, 1262, 1207, 1156, 1112, 1027, 955, 775, 732, 671, 562 cm⁻¹. ¹H NMR (200 MHz, CDCl₃): δ 1.47 (s, 24H), 5.88 (s, 8H). ¹³C NMR: δ 25.5, 36.6, 103.1, 158.6. EI-MS: *m/z* 433[M+H]⁺, 432[M]⁺, 419, 418, 417, 402, 387, 372, 201, 186.

Synthesis of Tetracyclohexyltetraoxaquaterene (5).- For this preparation the above procedure was applied to afford compound **5** in 45% yield, colorless needles, m.p. 268-269° (*lit.*¹³ 268°). *R*_f=0.43 (hexane-ethyl acetate 95:5 v/v). IR (KBr pellets): 2944, 2855, 2662, 1649, 1602, 1554, 1446, 1240, 1124, 1020, 1014, 969, 903, 874, 768, 709, 666, 573 cm⁻¹. ¹H NMR (200 MHz, CDCl₃): δ 1.48 (bs, 24H), 2.01 (bs, 16H), 5.86 (s, 8H). ¹³C NMR: δ 22.31, 26.12, 33.43, 40.75, 103.91, 157.45. EI-MS: *m/z* 594, 593[M+H]⁺, 592[M]⁺, 564, 549, 445, 377, 297, 229.

Anal. Calc. for C₁₀H₄₈O₄: C, 81.08; H, 8.11. Found C, 80.95; H, 8.18

Acknowledgments.- This work was supported by the State Committee for Scientific Research (project 3TO9A 126 15).

REFERENCES

1. C. D. Gutsche, *Calixarenes*, Royal Society of Chemistry, Cambridge, 1989.
2. V. Bohmer and M. A. McKervey, *Chemie Unserer Zeit*, 195 (1991)
3. J. Szejtli, *Cyclodextrins and Their Inclusion Complexes*, Akademiai Kiado, Budapest, 1982.
4. J. F. Stoddart, *Carbohydr. Res.*, 192 (1989).
5. G. Wenz, *Angew. Chem., Int. Ed. Engl.*, **33**, 803, (1994).
6. M. Chastrette and F. Chastrette, *Chem. Commun.*, 534 (1973).
7. M. Chastrette and F. Chastrette, Sabadie, *Org. Synth.*, **57**, 74 (1977).
8. M. de S. Healy and J. A. Rest, *Chem. Commun.*, 149 (1981).
9. M. de S. Healy and J. A. Rest, *J. Chem. Soc., Perkin Trans 1*, 973 (1985).
10. For X-ray analysis of compound **3** see: A. Hazell, *Acta Cryst.*, **C45**, 137 (1989).
11. W. H. Brown and B. J. Hutchinson, *Can. J. Chem.*, **56**, 617 (1978).
12. R. Pajewski, A. Pecak, R. Ostaszewski and J. Jurczak, *Acta Cryst.*, **C55**, 1862 (1999).
13. P. D. Williams and E. J. LeGoff, *J. Org. Chem.*, **46**, 4143 (1981).
