

Enantioselective Total Synthesis of (+)-Isoboonein

Masahiro TADA,* Sayako INOUE, Takao MIKI, Shin-ichi ONOGI, Jun-ichi KAMINAGA, Junko HIRAOKA, Yoshikazu KITANO, and Kazuhiro CHIBA

Laboratory of Bio-organic Chemistry, Tokyo University of Agriculture and Technology, Fuchu, Tokyo 183–8509, Japan.

Received March 9, 1998; accepted June 1, 1998

Enantioselective total synthesis of (+)-isoboonein, an iridoid lactone, was accomplished from (–)-dimenthyl bicyclo[2.2.1]hept-5-ene-2,3-dicarboxylate, which was synthesized by diastereoselective Diels–Alder reaction of dimethyl fumarate with cyclopentadiene.

Key words asymmetric; synthesis; isoboonein; iridoid

More than 600 iridoids have been isolated from medicinal, toxic, edible or other plants, however the biological activities of most iridoids are unreliable.¹⁾ (+)-Isoboonein (**1**) was isolated from *Rauwolfia grandiflora* by A. Bianco *et al.* in 1994.²⁾ The same compound, named as abelialactone, had been obtained earlier by hydrolysis of abelioside A from *Abelia grandiflora*,³⁾ and the structure was correlated with loganin.⁴⁾ Although the biological activity of **1** has not been reported, analogous cyclopentano- δ -lactone iridoids have been reported with unique biological activities, *e.g.* nepetalactone (**2**) with excitative activity toward cats,⁵⁾ and iridomyrmecin (**3**) with strong insecticide activity against preying insects, from ants (*Iridomyrmex humilis* MAYR.).⁶⁾ As the absolute stereochemistry of (+)-isoboonein (**1**) is similar to typical iridoids, enantioselective synthesis of **1** may provide not only a possible synthetic route to these iridoid lactones, but also useful synthetic intermediates to many other iridoids. In this paper, we describe an enantioselective total synthesis⁷⁾ of (+)-isoboonein from the diastereoselective Diels–Alder adduct (**7**) of dimethyl fumarate (**5**) with cyclopentadiene.⁸⁾

Results and Discussion

There are several reports of iridoid synthesis starting from bicyclo[2.2.1]heptanes.^{9–13)} Facile and highly diastereoselective reaction should be chosen to provide the first asymmetric intermediate for our enantioselective synthesis of iridoids. Diels–Alder reaction of dimethyl fumarate (**5**) with cyclopentadiene has been reported to give (–)-dimenthyl bicyclo[2.2.1]hept-5-ene-2,3-dicarboxylate (**7**) diastereoselectively.⁸⁾ Since the stereochemistry at C1, C2 and C3 of (–)-**7** is comparable to the stereochemistry at C4a, C7a and C7 of (+)-isoboonein (**1**), respectively, we planned to create the δ -lactone ring of **1** via the cleavage of the cyclopentene ring of **7**.

Synthesis of racemic isoboonein was examined first from racemic bicyclo[2.2.1]heptane derivative **6** and the same synthetic route was then applied for the synthesis of optically active (+)-isoboonein (**1**). The optically active diol **8** was synthesized via Diels–Alder reaction (92%) of **5**,⁸⁾ followed by reduction of **7** (78%) in 97% ee.^{14,15)} The optical yield was calculated from the $[\alpha]_D$ value of our synthetic **8** and the reported value.¹⁴⁾ The diol **8** was then converted to ketone **12** [optically active **8**→**9** (85%)→**10** (89%)→**11** (89%)→**12** (86%) in Fig. 2] according to the literature for the synthesis of (±)-**12**.¹⁶⁾ Oxidation of **12** with *m*-chloroperbenzoic acid (MCPBA) in dichloromethane provided lactone **13** [(±)-**13**,

97%; optically active **13**, 82%] which could be formed via Baeyer–Villiger oxidation of **12** followed by successive reactions of opening of the lactone ring in the oxidation product (**A**) and recyclization to lactone **13**. Since **13** has a carbon skeleton similar to isoboonein (**1**), with the same stereochemistry at C4a and C7a, the functional groups on the cyclopentane ring of **13** were modified to give **1** as follows. Since the stereochemistry at C6 of **13** is opposite to that of **1**, the hydroxy group was oxidized with pyridinium chlorochromate on alumina (PCC–Al₂O₃) in CH₂Cl₂¹⁷⁾ to afford ketone **14** [(±)-**14**, 80%; optically active **14**, 80%] which was then treated with trifluoroacetic acid–water–CH₂Cl₂ to give enone **15** [(±)-**15**, 55%; optically active **15**, 65%]. Hydrogenation of **15** with H₂ and 5% Pd–C in ethyl acetate (EtOAc) gave a mixture (5 : 1) of ketones **16** and **17** [(±)-**16** and **17**, 79%; optically active compounds **16** and **17**, 97%]. As the stereochemistry at C-7 of the major ketone **16** is opposite to **1**, the mixture was treated with pyridinium *p*-toluenesulfonate in refluxing benzene to give a mixture (1 : 8) of **16** and **17** [(±)-**16** and **17**, 75%; optically active **16** and **17**, 65%]. The major isomer **17** has the same stereochemistry as **1**. The mixture of **16** and **17** was then reduced with NaBH₄ in isopropanol and the product purified to give **18** [(±)-**18**, 68%; optically active **18**, 78%], an epimeric alcohol of isoboonein (**1**) at C6. The alcohol **18** was epimerized by sequential Mitsunobu reaction with azodicarboxylic acid diethyl ester, triphenylphosphine, and acetic acid in tetrahydrofuran (THF)¹⁸⁾ and hydrolysis with K₂CO₃ in methanol, and the resulting product was heated in refluxing benzene to recyclize to the lactone ring, to give isoboonein (**1**) [(±)-**1**, 22% from **18**; optically active **1**, 13% from **18**]. The synthetic (+)-**1**, $[\alpha]_D +65.0^\circ$ (*c*=0.08, methanol) was identical with natural isoboonein (**1**), $[\alpha]_D +65.0^\circ$ (*c*=0.2, methanol).

Since (+)-isoboonein (**1**) has similar absolute stereochemistry at C5, C8 and C9 to those of typical iridoids, this synthetic route should be useful for enantioselective synthesis of various iridoids.




Fig. 1.

* To whom correspondence should be addressed.




Fig. 2

a, cyclopentadiene, Et_2AlCl ; b, LiAlH_4 ; c, MCPBA; d, TBSO; e, oxalyl chloride, DMSO, Et_3N in CH_2Cl_2 ; f, HgCl_2 , Al, THF-EtOH; g, MCPBA.




Fig. 3

a, PCC- Al_2O_3 ; b, TFA- H_2O ; c, H_2 -5% Pd/C; d, reflux, PPTS in PhH; e, NaBH_4 in isopropanol; f, Ph_3P , $(\text{NCO}_2\text{C}_2\text{H}_5)_2$, $\text{CH}_3\text{CO}_2\text{H}$ in THF; g, K_2CO_3 in CH_3OH , HCl.

Experimental

All melting points were measured on a MEL-TEMP (Laboratory Device) without correction. IR and UV spectra were measured on a JEOL JIR-WIN-SPEC50 Fourier transform (FT)-infrared spectrometer and a JASCO UVDEC-460 spectrometer, respectively. Mass spectra were recorded on a JEOL JMS-SX-102A spectrometer. Nuclear magnetic resonance (^1H - and ^{13}C -NMR) spectra were measured on a JEOL JNM-EX270 spectrometer in CDCl_3 containing tetramethylsilane as internal standard, and J values are reported in Hz. TLC was carried out on Kiesel-gel GF₂₅₄ (0.25 mm thickness). Silica gel 60 (70–230 mesh ASTM) was used for column chromatography.

5,6-Di(hydroxymethyl)bicyclo[2.2.1]hept-2-ene (8) Optically active **8** was synthesized from **5** by similar procedures to those described in the literature (78%, 97% ee). Optically active **8**, white crystals, mp 40°C, $[\alpha]_D^{25} -22.2^\circ$ ($c=2.1$, CHCl_3 , at 23°C).

4-Hydroxymethyl-7-oxatricyclo[3.2.1.1^{3,8}]nonan-2-ol (9) Optically active **9** (85%), white crystals, mp 87°C, $[\alpha]_D^{25} -52.5^\circ$ ($c=2.1$, CH_3OH).

4-tert-Butyldimethylsilyloxymethyl-7-oxatricyclo[3.2.1.1^{3,8}]nonan-2-ol (10) Optically active **10** (89%), oil, $[\alpha]_D^{25} -20.0^\circ$ ($c=0.1$, CHCl_3).

4-tert-Butyldimethylsilyloxymethyl-7-oxatricyclo[3.2.1.1^{3,8}]nonan-2-one (11) Optically active **11** (89%), oil, $[\alpha]_D^{25} -63.0^\circ$ ($c=0.1$, CHCl_3).

5-Hydroxymethyl-6-(tert-butyldimethylsilyloxymethyl)bicyclo[2.2.1]heptan-2-one (12) Optically active **12** (86%), oil, $[\alpha]_D^{25} -25.0^\circ$ ($c=0.1$, CHCl_3).

4aR,6R,7S,7aS-Tetrahydro-6-hydroxy-7-(tert-butyldimethylsilyloxymethyl)cyclopenta[c]pyran-3(1H)-one (13) A solution of **12** (420 mg, 1.48 mmol) and MCPBA (1.28 g, 5.94 mmol) in 40 ml of CH_2Cl_2 was stirred for 3 d at ambient temperature. An excess amount of a solution of diazomethane in ether was added to the reaction mixture and the products were separated by silica gel column chromatography to give 433 mg (1.44 mmol) of **13** (97%). ($\pm$)-**13**, oil. MS m/z : 300 (M^+ , 1%), 285 (5), 243 (100), 195 (25), 183 (43), 151 (50), 105 (86). IR (neat) cm^{-1} : 3450, 2952, 2931, 2850, 1764, 1473, 1250, 1089. ^1H -NMR δ : 4.28 (1H, dd, $J=5.0$, 11.5 Hz), 4.20 (1H, dd, $J=5.5$, 11.5 Hz), 3.96 (1H, m), 3.80–3.65 (2H, m), 2.87 (1H, br), 2.62 (1H, dd, $J=6.5$, 13.8 Hz), 2.44 (1H, dd, $J=5.5$, 13.8 Hz), 2.52 (1H, m), 2.29 (1H, m), 2.13 (1H, m), 1.88 (1H, m), 1.39 (1H, m), 0.89 (9H, s), 0.07 (6H, s). ^{13}C -NMR δ : 173.4, 74.7, 69.3, 64.5, 50.6, 40.8, 37.2, 34.6, 30.5, 25.8, 18.1, -5.57.

Optically active **13** (82%), oil, $[\alpha]_D^{25} +38.0^\circ$ ($c=0.1$, CHCl_3).

4aR,7S,7aS-Tetrahydro-6-oxo-7-(tert-butyldimethylsilyloxymethyl)cyclopenta[c]pyran-3(1H)-one (14) To a solution of hydroxy lactone **13** (574 mg, 1.91 mmol) in dry CH_2Cl_2 (50 ml), PCC-alumina (10.3 g, 9.55 mmol) was added and the mixture was stirred for 6 h. After filtration and washing of the alumina with CH_2Cl_2 , the combined CH_2Cl_2 solution was evaporated and the residue was chromatographed on silica gel eluting with hexane-ethyl acetate to give 467 mg (1.56 mmol) of ketolactone **14** (80%). ($\pm$)-**14**: white crystals, mp 88°C. MS m/z : 298 (M^+ , 3%), 283 (5), 257 (25),

241 (100), 181 (43). IR (KBr) cm^{-1} : 2973, 1749, 1727, 1245, 1037. $^1\text{H-NMR}$ δ : 4.40 (1H, m), 4.18 (1H, dd, $J=5.1, 11.8$ Hz), 3.79 (1H, m), 3.71 (1H, m), 2.87–2.65 (3H, m), 2.51 (1H, m), 2.33 (1H, m), 2.26–2.03 (2H, m), 0.77 (9H, s), –0.05 (3H, s), –0.06 (3H, s). $^{13}\text{C-NMR}$ δ : 215.5, 171.7, 76.4, 61.7, 51.8, 44.5, 36.5, 34.6, 28.5, 25.4, 17.7, –6.00, –6.08.

Optically active **14** (79%), oil, $[\alpha]_{\text{D}} -10.0^\circ$ ($c=0.1$, CHCl_3).

4aR,7aS-Tetrahydro-6-oxo-7-methylenecyclopenta[c]pyran-3(1H)-one (15) A solution of **14** (21.0 mg, 0.07 mmol) in 6 ml of CH_2Cl_2 and 4 ml of trifluoroacetic acid– H_2O (9:1) was stirred for 12 h at ambient temperature. The solution was neutralized with saturated aqueous NaHCO_3 and the CH_2Cl_2 layer was washed with brine, dried over MgSO_4 , and evaporated. After purification by column chromatography (silica gel, hexane–ethyl acetate), enone **15** (6.40 mg, 0.039 mmol, 55%) was obtained. ($\pm$)-**15**: white crystals, mp 86°C . MS m/z : 166 (M^+ , 36%), 136 (95), 108 (33), 94 (100). IR (neat) cm^{-1} : 2973, 1749, 1727, 1635, 1245, 1037. $^1\text{H-NMR}$ δ : 6.23 (1H, d, $J=2.3$ Hz), 5.52 (1H, d, $J=2.3$ Hz), 4.52 (1H, dd, $J=4.6, 10.9$ Hz), 4.21 (1H, dd, $J=6.9, 10.9$ Hz), 3.44 (1H, m), 3.00–2.80 (3H, m), 2.42 (1H, dd, $J=6.3, 14.4$ Hz), 2.24 (1H, dd, $J=5.3, 16.4$ Hz). $^{13}\text{C-NMR}$ δ : 204.0, 171.6, 143.4, 121.6, 69.3, 43.6, 38.2, 34.8, 27.5.

Optically active **15** (65%), mp 101°C , $[\alpha]_{\text{D}} +100.0^\circ$ ($c=0.1$, CHCl_3).

4aR,7aS-Tetrahydro-6-oxo-7-methylcyclopenta[c]pyran-3(1H)-one (16 and 17) A mixture of the enone **15** (58.5 mg, 0.352 mmol) and 5% Pd–C (19.8 mg) in ethyl acetate (8 ml) was stirred under H_2 for 72 h. After filtration through celite, the solution was evaporated and the residue was chromatographed over silica gel eluting with hexane–ethyl acetate to afford a mixture of **16** and **17** (5:1, 46.5 mg, 0.276 mmol, 79%). This mixture was then refluxed with 4.3 mg (0.017 mmol) of pyridinium *p*-toluenesulfonate in benzene (10 ml) for 5 h. After evaporation, the residue was chromatographed over silica gel to give a mixture of **16** and **17** (1:8, 34.9 mg, 0.208 mmol, 75%). ($\pm$)-**16**: oil. MS m/z : 168 (M^+ , 25%), 152 (40), 139 (40), 126 (100), 111 (71), 97 (74). IR (neat) cm^{-1} : 2967, 1733, 1255, 1178. $^1\text{H-NMR}$ δ : 4.32 (1H, dd, $J=4.5, 12.0$ Hz), 4.08 (1H, dd, $J=9.3, 12.0$ Hz), 3.00–2.23 (6H, m), 2.04 (1H, m), 1.10 (3H, d, $J=6.5$ Hz). $^{13}\text{C-NMR}$ δ : 208.3, 175.6, 66.4, 45.3, 42.4, 37.3, 34.8, 28.0, 9.16. ($\pm$)-**17**: white crystals, mp 79°C . MS m/z : 168 (M^+ , 100%), 153 (8), 140 (20), 127 (55), 96 (10). IR (neat): 2973, 2931, 1743, 1286, 1245, 1089 cm^{-1} . $^1\text{H-NMR}$ δ : 4.53 (1H, dd, $J=4.0, 11.6$ Hz), 4.29 (1H, dd, $J=4.3, 11.6$ Hz), 3.00–2.78 (3H, m), 2.63 (1H, dd, $J=9.2, 18.9$ Hz), 2.45 (1H, dd, $J=5.7, 15.4$ Hz), 2.38–2.18 (2H, m), 1.14 (3H, d, $J=6.5$ Hz). $^{13}\text{C-NMR}$ δ : 216.0, 171.8, 68.8, 43.7, 43.2, 40.6, 34.8, 27.9, 13.2.

Optically active **16** and **17** (1:8, 65%).

4aR,6R,7aS-Tetrahydro-6-hydroxy-7-methylcyclopenta[c]pyran-3(1H)-one (18) To a solution of **17** (35.3 mg, 0.21 mmol) in isopropanol (6 ml), NaBH_4 (48.1 mg, 1.26 mmol) was added at 0°C and the solution was stirred for 10 min. After addition of saturated aqueous NH_4Cl , the products were extracted with CHCl_3 repeatedly and the solution was washed with brine, dried over MgSO_4 , and evaporated to give alcohol **18** (24.4 mg, 0.143 mmol, 68%). ($\pm$)-**18**: oil. MS m/z : 170 (M^+ , 28%), 152 (69), 139 (28), 126 (100), 111 (71). IR (neat) cm^{-1} : 3460, 2962, 2925, 1737, 1255, 1162, 1062. $^1\text{H-NMR}$ δ : 4.30 (1H, dd, $J=4.8, 11.5$ Hz), 4.18 (1H, dd, $J=5.6, 11.5$ Hz), 3.70 (1H, m), 2.68–2.40 (3H, m), 2.27 (1H, m), 1.99 (1H, m), 1.72 (1H, m), 1.58 (1H, brs), 1.34 (1H, ddd, $J=9.3, 12.9, 12.9$ Hz), 1.09 (3H, d, $J=6.6$ Hz). $^{13}\text{C-NMR}$ δ : 173.9, 78.0, 68.7, 43.9, 42.1, 40.9, 34.8, 30.3, 16.5.

Optically active **18** (78%), $[\alpha]_{\text{D}} +64.0^\circ$ ($c=0.1$, CHCl_3).

Mitsunobu Inversion of Alcohol 18 to Isoboonein (1) To a solution of alcohol **18** (32.2 mg) in THF (4 ml), Ph_3P (71.2 mg, 0.271 mmol), $\text{CH}_3\text{CO}_2\text{H}$ (0.022 ml, 0.384 mmol), and $(\text{NCO}_2\text{C}_2\text{H}_5)_2$ (0.036 ml, 0.228 mmol) were added successively at 0°C , and the solution was stirred at ambient temperature for 24 h. After evaporation, the residue was chromatographed over silica

gel eluting with hexane–ethyl acetate to give crude acetate **19** (16.9 mg). **19**: oil. MS m/z : 212 (M^+ , 7%), 170 (65), 152 (100), 126 (64), 110 (53), 93 (61). IR (neat) cm^{-1} : 2967, 2931, 1737, 1245. $^1\text{H-NMR}$ δ : 5.19 (1H, t, $J=2.8$ Hz), 4.33 (1H, dd, $J=4.0, 12.1$ Hz), 4.17 (1H, dd, $J=3.0, 12.1$ Hz), 2.86 (1H, m), 2.64 (1H, dd, $J=6.8, 14.9$ Hz), 2.38 (1H, dd, $J=4.1, 14.9$ Hz), 2.06 (3H, s), 2.22–2.00 (3H, br), 1.49 (1H, ddd, $J=3.4, 11.5, 14.2$ Hz), 1.02 (3H, d, $J=5.4$ Hz). $^{13}\text{C-NMR}$ δ : 173.0, 170.6, 78.4, 68.3, 42.4, 40.3, 39.2, 34.4, 32.6, 21.1, 12.9. The crude acetate **19** was hydrolyzed without further purification. To a solution of **19** in methanol (5 ml), K_2CO_3 (25 mg, 0.255 mmol) was added and the mixture was stirred for 12 h at ambient temperature. After neutralization of the mixture with 1 M HCl, the solvent was evaporated and the residue was extracted with benzene. The benzene solution was refluxed for 2 h to complete the lactonization and then evaporated. The residue was chromatographed over silica gel to give isoboonein (**1**) (6.6 mg, 0.039 mmol, 22% from **18**). ($\pm$)-**1**: oil. MS m/z : 170 (M^+ , 38%), 152 (33), 139 (100), 124 (60), 111 (55), 97 (45). IR (neat) cm^{-1} : 3466, 3309, 2962, 2931, 1727, 1234, 1068. $^1\text{H-NMR}$ δ : 4.32 (1H, dd, $J=4.0, 12.1$ Hz), 4.16 (1H, dd, $J=3.0, 12.0$ Hz), 4.15 (1H, m), 2.95 (1H, m), 2.65 (1H, dd, $J=7.3, 15.1$ Hz), 2.38 (1H, dd, $J=3.9, 15.1$ Hz), 2.16 (1H, m), 2.06 (1H, ddd, $J=1.0, 8.2, 13.6$ Hz), 1.94 (1H, ddq, $J=4.0, 7.0, 10.2$ Hz), 1.43 (1H, ddd, $J=4.0, 9.6, 13.3$ Hz), 1.08 (3H, d, $J=7.0$ Hz). $^{13}\text{C-NMR}$ δ : 173.5, 75.6, 68.6, 41.7, 41.6, 41.5, 34.5, 32.7, 12.7.

Optically active **1** (13% from **18**), oil, $[\alpha]_{\text{D}} +65.0^\circ$ ($c=0.08$, CH_3OH).

References and Notes

- For reviews on iridoids: Grayson D. H., *Natural Product Reports*, **13**, 214–217 (1997); El-Naggar L. J., Beal J. L., *J. Nat. Prod.*, **43**, 649–707 (1980).
- Bianco A., de Luca A., Mazzei R. A., Nicoletti M., Passacantilli P., de Lima R. A., *Phytochemistry*, **35**, 1485–1487 (1994).
- Murai F., Tagawa M., Matsuda S., Kikuchi T., Uesato S., Inoue H., *Phytochemistry*, **24**, 2329–2335 (1985).
- Brechbuhler-Bader S., Coscia C. J., Loew P., von Szczepanski C., Arigoni D., *Chem. Commun.*, **1968**, 136–137; Battersby A. R., Hall E. S., Southgate R., *J. Chem. Soc., (C)*, **1969**, 721–728; Buchi G., Carlson J. A., Powell Jr. J. E., Tietze L. F., *J. Am. Chem. Soc.*, **95**, 540–545 (1973).
- Sakan T., Murai F., Isoe S., Hyeon S. B., Hayashi Y., *Nippon Kagaku Zasshi*, **90**, 507–528 (1969).
- Fuson R., Trave R., Vercellon A., *Chim. Ind. (Milan)*, **37**, 251–258 (1955).
- For the total synthesis of ($\pm$)-isoboonein: Ohba M., Haneishi T., Fujii T., *Chem. Pharm. Bull.*, **44**, 525–529 (1996).
- Furuta K., Iwanaga K., Yamamoto H., *Tetrahedron Lett.*, **27**, 4507–4510 (1986).
- Kelly R. C., Schletter I., *J. Am. Chem. Soc.*, **95**, 7156–7157 (1973).
- Furuchi K., Abe K., Miwa T., *Tetrahedron Lett.*, **1974**, 3685–3688.
- Arai Y., Kawanami S., Koizumi T., *Chem. Lett.*, **1990**, 1585–1586.
- Fleming I., Terrett N. K., *Tetrahedron Lett.*, **25**, 5103–5104 (1984).
- Callant P., Storme P., Van der Eycken E., Vandewalle M., *Tetrahedron Lett.*, **24**, 5797–5800 (1983).
- Horton D., Machinami T., Takagi Y., *Carbohydr. Res.*, **121**, 135–161 (1983).
- Takano S., Kurotaki A., Ogasawara K., *Synthesis*, **1987**, 1075–1078.
- Van der Eycken E., De Bruyn A., Van der Eycken J., Callant P., Vandewalle M., *Tetrahedron*, **42**, 5385–5396 (1986).
- Cheng Y. S., Liu W. L., Chen S. H., *Synthesis*, **1980**, 223–224.
- Mitsunobu O., *Synthesis*, **1981**, 1–28.