


Accepted Manuscript

Comparative study between 1-Propyl-3-methylimidazolium bromide and trimethylene bis-methylimidazolium bromide ionic liquids by FTIR/ATR and FT-RAMAN spectroscopies



Mohamed Kadari, El Habib Belarbi, Taqiyeddine Moumene, Serge Bresson, Boumediene Haddad, Ouissam Abbas, Brahim Khelifa

PII: S0022-2860(17)30513-6
DOI: 10.1016/j.molstruc.2017.04.076
Reference: MOLSTR 23699
To appear in: *Journal of Molecular Structure*
Received Date: 14 November 2016
Revised Date: 15 April 2017
Accepted Date: 17 April 2017

Please cite this article as: Mohamed Kadari, El Habib Belarbi, Taqiyeddine Moumene, Serge Bresson, Boumediene Haddad, Ouissam Abbas, Brahim Khelifa, Comparative study between 1-Propyl-3-methylimidazolium bromide and trimethylene bis-methylimidazolium bromide ionic liquids by FTIR/ATR and FT-RAMAN spectroscopies, *Journal of Molecular Structure* (2017), doi: 10.1016/j.molstruc.2017.04.076

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Synthesis of ionic liquids ($[\text{PrMIM}^+][\text{Br}^-]$), ($[\text{M}(\text{CH}_2)_3\text{IM}^{2+}][2\text{Br}^-]$).
- FTIR/ATR and FT-RAMAN spectroscopy study of previous ionic liquids.
- Comparison between dicationic ionic liquid and monocationic ionic liquid.
- Each ionic liquid has a specific vibrational behavior.
- The passage from mono to dicationic is responsible of important vibrational changes.

1 **Comparative study between 1-Propyl-3-methylimidazolium bromide and**
2 **trimethylene bis-methylimidazolium bromide ionic liquids by FTIR/ATR and**
3 **FT-RAMAN spectroscopies**

4 Mohamed KADARI¹, El Habib BELARBI¹, Taqiyeddine MOUMENE¹, Serge BRESSON³,
5 Boumediene HADDAD¹, Ouissam ABBAS², Brahim KHELIFA³

6
7 ¹ Laboratoire de Synthèse et Catalyse Tiaret LSCT, Univ Tiaret, BP 78 RP, 14000 Tiaret, Algeria

8 ²Centre Wallon de Recherche Agronomique, CRA-W, Bâtiment Maurice Henseval, chaussée de
9 Namur, 24, 50030 Gembloux, Belgium.

10 ³Laboratoire de Physique des Systèmes Complexes, Université Picardie Jules Verne, 33 rue S^t Leu
11 80039 Amiens cedex, France

12
13 **Abstract:**

14 In this study, we synthesized two ionic liquids based on imidazolium: one is a monocationic and the
15 other is a dicationic. They are respectively 1-Propyl-3-methylimidazolium bromide ([PrMIM⁺][Br⁻]) and
16 trimethylene bis-methylimidazolium bromide ([M(CH₂)₃IM²⁺][2Br⁻]). The structures of these two ionic
17 liquids which are composed of ions with atoms of the same nature were first identified by ¹H, ¹³C NMR,
18 and then compared in a study by FT-RAMAN and FTIR/ATR spectroscopies.

19 FT-RAMAN spectras of the dicationic ionic liquid are richer in modes in the different spectral regions.
20 Hence this richness seems to be a consequence of the passage from one to two rings in the imidazolium
21 cation. In particular, the vibrational modes in the spectral ranges 700 – 600 cm⁻¹, 1700 – 1500 cm⁻¹ and 3200
22 – 2700 cm⁻¹ by FTIR/ATR seem to be sensitive to the change from mono to dicationic than in FT-RAMAN.
23 The spectral range in which the intermolecular interactions are present (200-50 cm⁻¹) is a marker of
24 differentiation between the mono and the dicationic. The spectral ranges on 1700 – 1200 cm⁻¹ and 3200 –
25 2700 cm⁻¹ also show signs of upheaval between our two samples. We can also notice that there are much
26 more active modes in FT-RAMAN spectroscopy than in FTIR / ATR spectroscopy.

27
28 **Keywords:** monocationic and dicationic ionic liquids, imidazolium, RAMAN spectroscopy, FTIR/ATR
29 spectroscopy.

1 **1. Introduction:**

2 Ionic liquids are liquids that consist almost exclusively of ions (an organic cation associated
3 with an organic or inorganic anion). They are a class of salts with melting point temperature below
4 a hundred degrees. J. M. Crosthwaite et al. noticed that the ionic liquids based on pyridinium
5 become liquid around the 80°C [1]. J. D. Holbrey and K. R. Seddon obtained the same results for
6 ionic liquids based on imidazole [2]. In addition to their low melting points, ionic liquids are
7 characterized by exceptional thermal stability, enabling their use for applications at high
8 temperatures. They have a good ionic conductivity, generally of the order of 10^{-1} .S.m⁻¹ [3]. Their
9 electrochemical window is wide as they can provide tension between 5-6 V [4], which has been
10 exploited in the field of fuel cells [5, 6]. The Ionic liquids have also remarkable solvation capacities
11 of organic and inorganic products comparable to polar solvents. Notwithstanding the fact that they
12 are used as solvents in the biocatalytic reactions [7], it is most interesting to note that their
13 physicochemical properties are adjustable by changing the nature of the anion and/or the cation
14 (alkyl chain).

15 The applications of ionic liquids are dependent on their properties, while the latter dependent
16 on the nature of the cation and anion combination and the length of the alkyl chains incorporated on
17 the cation. The possible combinations of cations/anions are very numerous and continuously
18 evolving, such that in general the properties of ionic liquids are governed by Coulomb forces,
19 hydrogen bonding and Van der Waals interactions. In this paper, we will study two imidazolium-
20 based ionic liquids .

21 In our study of both ionic liquids: a monocationic and dicationic which are respectively 1-
22 Propyl-3-methylimidazolium bromide ([PrMIM⁺][Br⁻]) and trimethylene bis-methyl imidazolium
23 bromide ([M(CH₂)₃IM²⁺][2Br⁻]), the nature of atoms constituting the anions and cations of both
24 (ILs) are similar. However the difference is that the cation of the monocationic (IL) is constituted of
25 a single imidazolium ring while the cation of the dicationic consists of two imidazolium rings. So
26 two singly charged cations linked by an alkyl chain and paired with two singly charged anions. In
27 this study, we wanted to compare the vibrational behavior of these two ILs. Research conducted on
28 the properties of ionic liquids and which is based on the understanding of the relationship between
29 the structure of the cation and that of the anion aims at obtaining (ILs) with better properties
30 whilst improving the theoretical model.

31 The study of the vibrational behavior of a material can be achieved by means of the following

1 characterization techniques: FTIR spectroscopy, and RAMAN spectroscopy. These two non-
2 destructive techniques are complementary and represent the best performing and most widely used
3 tools. For monocationic ILs many studies have been carried out using these techniques [8, 9]. This
4 is not the case for dicationic ILs.

5 **2. Experimental**

6 **2.1. Reagents and materials**

7 The reagents used in this study are 1-methylimidazole (98 wt.%), 1-bromopropane (99.5
8 wt.%). They were purchased from Fluka and used as received.


9 **2.2. NMR spectroscopy analysis**

10 ^1H NMR (400 MHz), ^{13}C NMR, spectra were recorded on a NMR 400 MHz spectrometer. The
11 chemical shifts (δ) are given in ppm and referenced to the internal solvent signal, TMS
12 (Tetramethylsilane) and CFCl_3 , respectively.

13 **2.3. Synthesis and characterization of ionic liquids:**


14 **2.3.1. Synthesis of $[\text{PrMIM}^+][\text{Br}^-]$:**

15 The synthesis was prepared according to procedures describe in the literature. Briefly, 1-
16 methylimidazole (6.79 ml, 85.26 mmol), and 1-bromopropane (9.87 ml, 89.03 mmol) were
17 combined at a 1:1 ratio before being stired at 70°C for 6 hours. The $[\text{PrMIM}^+][\text{Br}^-]$ was then washed
18 with ethyl acetate to remove any excess reactants or contaminants. Sebsequently, about 20 ml of
19 ethyl acetate was added to the $[\text{PrMIM}^+][\text{Br}^-]$ and the mix shaken together in a separatory funnel.
20 After a few minutes, the mixture separated into two distinct layers; the top layer being ethyl acetate
21 and all of the contaminants and the bottom layer was $[\text{PrMIM}^+][\text{Br}^-]$. The top layer, which was
22 discarded later, was poured into a clean beaker. The bottom layer was saved in the original flask .
23 This washing procedure was repeated four times.



1 2.3.2. Synthesis of ([M(CH₂)₃IM²⁺][2Br⁻):

2 In a flask of 100 ml, we prepared a mixture of 1-methyl imidazole (9.07 ml, 100 mmol) and
3 1,3-dibromopropane (5.21 ml, 49.98 mmol) and 23 ml of DMF. This reaction mixture is heated
4 under reflux with magnetic stirring for 4 hours while maintaining the temperature at 75°C. Bromide
5 trimethylene bis-methyl imidazolium was obtained as a slightly yellow solid (6.60g, 17.12 mmol)
6 with a yield of 63%.



8 2.4. NMR results:

9 The spectra details are given below,

10 ¹H-NMR (CDCl₃) δ_H (ppm) [PrMIM⁺][Br⁻]: 10.27 (1H, s, N⁺CHN), 7.58 (1H, s, N⁺CHCHNCH₃),
11 7.49 (1H, s, N⁺CHCH), 4.26 (3H, s, NCH₃), 1.95-1.93 (2H, m, N⁺CH₂CH₂), 1.87-1.85 (2H, m,
12 CH₂CH₂CH₃), 0.91 (3H, t, J=6.8 Hz, N⁺(CH₂)₂CH₃).

13

14 ¹³C-NMR (CDCl₃) δ_C (ppm) [M(CH₂)₃IM²⁺][2Br⁻]: 10.67 (CH₂CH₂CH₃), 32.62 (CH₂CH₂CH₃),
15 36.62 (NCH₃), 51.35 (N⁺CH₂CH₂), 122.32 (NCHCHN⁺), 123.75 (NCHCHN⁺), 137.01 (N⁺CHN).

16 2.5. FTIR/ATR and FT-RAMAN measurements

17 The measurements were realised in the Walloon Agricultural Research Center (CRA-W)
18 Belgium.

19 2.5.1. FTIR/ATR measurements

20 All attenuated total reflectance Fourier transform mid-infrared (ATR/FTIR) spectra were
21 obtained on a Vertex 70-RAM II Bruker spectrometer (Bruker Analytical, Madison, WI) operating
22 with a Golden Gate TM diamond ATR accessory (Specac Ltd, Slough, UK).

23 FTIR/ATR spectra [4000-600 cm⁻¹] were collected with a resolution of 1 cm⁻¹ by co-adding 64
24 scans for each spectrum. The OPUS 6.0 software for windows of Bruker Instruments was used for
25 instrument management.

26 2.5.2. FT-RAMAN measurements

27 FT-RAMAN spectra were obtained on a Vertex 70-RAM II Bruker FT-RAMAN spectrometer.
28 This instrument is equipped with a Nd: YAG laser (yttrium aluminium garnet crystal doped with
29 triply ionized neodymium) with a wavelength for incident laser at 1064 nm (9398,5 cm⁻¹). The

1 maximum laser power was 1.5 W. The measurement accessory was pre-aligned, only the Z-axis of
 2 the scattered light was adjusted to set the sample in the appropriate position relation to the local
 3 point. The RAM II spectrometer is equipped with a liquid-nitrogen cooled Ge detector. FT-
 4 RAMAN spectra [4000-45cm⁻¹] were collected with a resolution of 1 cm⁻¹ by co-adding 256 scans
 5 for each spectrum. The OPUS 6.0 software was used for the spectral acquisition, manipulation and
 6 transformation.

7 3. Results and Discussion

8 3.1. Comparison between the FTIR/ATR spectra of ionic liquids: ([PrMIM⁺][Br⁻]) and 9 ([M(CH₂)₃IM²⁺][2Br⁻):

10 Table 1 below shows the observed FTIR/ATR bands and their assignments in the spectral
 11 region of 600 cm⁻¹ to 3420 cm⁻¹ of two ionic liquids mono and di-cationic [PrMIM⁺] [Br⁻] and
 12 ([M(CH₂)₃IM²⁺][2Br⁻]). Figures 1, 2 and 3 show the infrared spectra of the two samples in the
 13 spectral regions 1200 cm⁻¹ to 600 cm⁻¹, 1800 cm⁻¹ to 1200 cm⁻¹ and 3700 cm⁻¹ to 2700 cm⁻¹
 14 respectively. There are 30 peaks in this zone for the dicationic and 34 peaks for the monocationic.

15 **Table 1**

16 Observed FTIR/ATR bands of ([PrMIM⁺][Br⁻] and ([M(CH₂)₃IM²⁺][2Br⁻]) and their assignments
 17 (vw = very weak; w = weak; m = medium, s = strong; sh = shoulder; v=st=stretching; δ=def=
 18 deformation; bend = bending deformation; ω=wagging; ρ= rocking; γ=bending out-of-plane;
 19 t=twist=twisting, s = symmetric; as = antisymmetric, * = fitted value).


[M(CH ₂) ₃ IM ²⁺][2Br ⁻]	[PrMIM ⁺][Br ⁻]	Vibrational assignment	Refs.
620 (vs)	618 (s)	CH ₃ (N)CN str	[12]
639 (m)		γ(N-H)	[17]
647 (m)	649 (m)	CH ₃ NCN str	[9]
660 (m)	660 sh	CH def-vib, NH ₂ wagging vib	[15]
694 (w)	696 (w)	Out-of-plane C-H def vib	[15]
714 (w)		CH ₂ (N) and CH ₃ (N)CN str	[10]
756 (m)	752 (m)	δHCCH/Ring HCCH asym bend/ω(C-H)	[8]/[13]
	800 (m)	Ring HCCH asym bend/CH ₂ rocking def- vib	[8]/[15]
826+836 (m)	827 (m)	NC(H)N bend/CCH bend/ω(CH ₂) and ω(CH)	[12]/[11]
	864 (m)	ω(C-H)/CH ₂ out-of-plane def vib	[11]/[15]
	900 (w)	C-H out-of-plane vib/CH ₂ rocking vib	[15]
	1010 (w)	v(C-C)/ρ(C-H) [ip]/CH ₃ rocking vib	[11]/[15]
1022 (w)	1019 (w)	ρ(CH ₂), t(CH ₂)/CH ₃ rocking vib and CN str	[11]/[15]
1062 (w)	1049 (w)	CC str, NCH ₃ twist/CH ₃ rocking vib	[8]/[15]
1097 (m)	1089 (w)	CC str/ρ(N-C)	[12]/[11]
	1116 (w)	ρ(C-H)	[11]
1165 (vs)	1168 (vs)	CH ₂ (N) and CH ₃ (N)CN str, CC str	[8]
	1230 (vw)	CH ₂ twisting vib/v(CC)	[15]/[17]
1253 (vw)		t(CH ₂)/ CH ₂ wagging vib.	[17]/[15]

	1272 (w)	CH ₂ wagging vib/ ρ (C-H)	[15]/[11]
	1297 (w)	CH ₂ twisting vib/ γ (C-H)	[17]
1339 (w)	1336 (w)	CH ₂ (N) and CH ₃ (N) CN str	[8]
	1345 (w)	CH ₂ (N), CH ₃ (N) CN str	[8]
1389 (m)	1385 (w)	CH ₂ (N) CN bend, CH ₂ (N) and CH ₃ (N) CN str/ ρ (N-H)	[8]/[13]
	1428 (w)	CH ₃ (N) str, CH ₃ (N) HCH sym bend or NC(CH ₃)N HCH bend	[8]
1458 (m)	1458 (m)	δ (CH ₂)/CCH HCH asym bend, CH ₃ (N)HCH sym bend, CH ₃ (N) str	[18]/[12]
1562 (s)	1564 (s)	ν (N=C)/ ρ (C-H)	[11]
1574 (s)	1570 (s)	ring ip sym/asym str, CH ₃ (N) str, CH ₂ (N) CN str	[12]
1655 (s)		ν (C=C), ν (C-N), δ (N-H)	[17]
2858 (vw)	2859*	ν_s (CH ₂)	[17]
	2876 (w)	ν_{as} (CH ₂)	[18]
2928* sh	2933* sh (m)	ν_s (CH ₃)	[18]
2952 (w)	2964 (m)	ν_{as} (CH ₂ HCH)	[8]/[12]
3030* sh	3014* sh	ν (C-H)	[11]
3085 (m)	3062 (m)	ν (C-H)	[11]
3145 (m)	3140 (w)	ν (C-H)	[8]
3245* sh		Free N-H str	[15]
3362* sh	3385* (w)	Free N-H str	[15]
3430 (s)	3450* (w)	Free N-H str	[15]
3495* sh		Free N-H str	[15]

1 3.1.1. The region 1200-600 cm⁻¹:

2 Almost the same number of peaks were observed for the two samples in this spectral zone: 15
3 peaks for the monocationic and 13 peaks for the dicationic. Although the spectra of these two
4 samples are similar, there are peaks in the spectrum of the mono that do not exist in the spectrum of
5 dicationic, and vice versa. With regard to the modes present in the two spectra, some peaks which
6 have undergone wavenumber shifts of the order of 4 cm⁻¹ were observed.. The Peaks at 620, 694,
7 756, 1022 and 1165 cm⁻¹ for the dicationic corresponded respectively to the peaks at 618, 696, 752,
8 1019 and 1168 cm⁻¹ for the monocationic. These peaks are assigned to the stretching vibration of
9 CH₃(N)CN and CH₂(N) bonds [8, 9, 12] and C-N stretching [15]. It is also noted that in the region
10 675-635 cm⁻¹ the peak at 649 cm⁻¹ with a shoulder at 660 cm⁻¹ became a triplet for the case of
11 dicationic ionic liquid: 639, 647 and 660 cm⁻¹. Therefore the passage from one to two rings
12 imidazolium has an influence on the vibrational behavior of ring bonds. Furthermore the following
13 observations were made: deformation (HCCH) and wagging (C-H) for 752 cm⁻¹ [13], stretching (C-
14 C) for 1168 cm⁻¹ [11], rocking (CH₂ and CH₃) and twisting (CH₂) for 1019 cm⁻¹ [11] and out-of-
15 plane C-H def-vib for 696 cm⁻¹ [15]. Therefore, this passage also has an effect on the vibrational
16 modes of other bonds of the cation. Other peaks underwent more important wavenumber shifts of
17 the order of 12 cm⁻¹, for example the peaks at 1062 and 1097 cm⁻¹ for the dicationic corresponded
18 to the ones at 1049 and 1089 cm⁻¹ for the monocationic. In the literature, these peaks are attributed

1 to the stretching vibration of the (C-C) bond [8,12], the rocking of the (N-C) [11] and the twisting
2 of NCH₃ [8] indicating that this zone is very sensitive to the passage of the mono to dicationic ionic
3 liquid.



4
5 *Fig. 1. FTIR/ATR spectra of: [PrMIM⁺][Br⁻] and ([M(CH₂)₃IM²⁺][2Br⁻]) in the region*
6 *1200-600 cm⁻¹.*

7 3.1.2. The region 1800-1200 cm⁻¹:

8 In this spectral region, there were more peaks in the case of monocationic than in the case of
9 dicationic, which can be explained by a congestion masking certain vibrational modes for the
10 dicationic. The monocationic spectrum peaks are a little broader than the dicationic ones,
11 suggesting more order in the dicationic structure. We observed a very intense new peak at 1655 cm⁻¹
12 in the case of dicationic. In the literature, this peak is attributed to stretching vibration of (C=C)
13 and (C-N) bonds, and deformation of (N-H) bond [17]. Its appearance and its intensity can be
14 explained by the increase in the number of these bonds ((C=C), (C-N), (N-H)) that represent a
15 signature of the passage of the mono to dicationic ionic liquid.





Fig.2. FTIR/ATR spectra of: $([PrMIM^+][Br^-])$ and $([M(CH_2)_3IM^{2+}][2Br^-])$ in the region from 1800 to 1200 cm^{-1} .

3.1.3. The region 3700-2700 cm^{-1} :

In this region, almost the same number of peaks for both samples was seen, 9 peaks for the monocationic and 10 peaks for the dicationic. The spectral band between 3500 - 3200 cm^{-1} was much higher in intensity for the dicationic compared to the monocationic. An actual rollover was observed in intensity in the spectral regions 3200-2800 cm^{-1} and 3500-3200 cm^{-1} between the monocationic and dicationic. This phenomenon was explained by computing different intensity ratios shown in Table 2 by referring to the peak at 2955 cm^{-1} . The increase in intensity is obvious, especially on the modes corresponding to $\nu(CH)$ at 3145 cm^{-1} [8] and $\nu(NH)$ at 3430 cm^{-1} . The number of vibrational modes in the spectral region 3500-3200 cm^{-1} also increased, 2 modes for monocationic, 4 modes for dicationic assigned to $\nu(NH)$. All these variations seem to be explained by the increased number of N-H bonds.


Table 2. Reports of intensity of many peaks of $[PrMIM^+][Br^-]$ and $[M(CH_2)_3IM^{2+}][2Br^-]$ in the spectral range 3500-2800 cm^{-1} .

Mono or dicationic ionic liquid	I_{2850}/I_{2955}	I_{3075}/I_{2955}	I_{3145}/I_{2955}	I_{3430}/I_{2955}
$[PrMIM^+][Br^-]$	0.48 ± 0.02	0.92 ± 0.02	0.36 ± 0.02	0.20 ± 0.02
$[M(CH_2)_3IM^{2+}][2Br^-]$	0.68 ± 0.02	1.68 ± 0.02	1.16 ± 0.02	2.27 ± 0.02



1
2
3

Fig.3. FTIR/ATR spectra of $:[PrMIM^+][Br^-]$ and $:[M(CH_2)_3IM^2+][2Br^-]$ in the region 3700-2700 cm^{-1} .



4
5
6
7
8

Fig.4. The Results of Lorentzian curve fitting of FTIR / ATR spectra for: (a) $:[PrMIM^+][Br^-]$ and (b) $:[M(CH_2)_3IM^2+][2Br^-]$ in the region 3700-2700 cm^{-1} .

1 **3.2. Comparison between FT-RAMAN spectra of ionic liquids ([PrMIM⁺][Br⁻]) and**
 2 **([M(CH₂)₃IM²⁺][2Br⁻]):**

3 Table 3 below represents the observed FT-RAMAN bands and their assignments for both ionic
 4 liquids ([PrMIM⁺][Br⁻]) and ([M(CH₂)₃IM²⁺][2Br⁻]). Figures 5, 6, 7 and 8 show the FT-RAMAN
 5 spectra of the two samples in the spectral regions 1000 cm⁻¹ to 45 cm⁻¹, 200 cm⁻¹ to 45 cm⁻¹, 1700
 6 cm⁻¹ to 1000 cm⁻¹ and 3200 cm⁻¹ to 2700 cm⁻¹ respectively. We noticed that the FT-RAMAN
 7 spectrum of dicationic was richer in vibrational modes than that of monocationic: 75 peaks for the
 8 dicationic and 38 picks for the monocationic.

9 **Table 3:**

10 Observed FT-RAMAN bands and their assignments (vw = very weak; w = weak; m = medium, s =
 11 strong; sh = shoulder; v = str = stretching; δ = def = deformation; bend = bending deformation; ω =
 12 wagging, ρ = rocking, τ = torsion; t = twist = twisting; β = bending in-plane, s = symmetric; as =
 13 antisymmetric, [ip] = in plane, [op] = out of plane).

[M(CH ₂) ₃ IM ²⁺][2Br ⁻]	[PrMIM ⁺][Br ⁻]	Vibrational assignment	Refs.
57 (s)	57 (vs)	Intermolecular vibration	[14]
68 (s)		Intermolecular vibration	[14]
80 (s)		Intermolecular vibration	[14]
86 (vs)		Intermolecular vibration	[14]
100 (s)		Intramolecular vibration	[14]
127 (m)		Intramolecular vibration	[14]
146 (m)		Intermolecular vibration	[14]
189 (m)		Intermolecular vibration / τ (N-C)/ ρ (CH ₂)	[14]/[11]
	228 (vw)	β (CCC)	[11]
256 sh+267 (w)		ρ (N-C)	[11]
	282 (vw)	CH ₃ torsional vib	[15]
299 (vw)		CH ₂ (N) and CH ₃ (N)CH bend	[8]
317 (vw)	316 (vw)	CH ₂ (N) and CH ₃ (N)CH bend	[8]
	365 (vw)	ν (C-C)	[11]
418 (m)	418 (vw)	t(CH ₂)	[11]
429 (vw)		CC def vib [op]	[15]
475 (w)	461 (vw)	ν (C-H)	[15]
595 (vw)	601 (w)	ω (N-C)	[11]
613 (m)		ω (N-H)/CH ₂ str, CH ₃ (N)CN str	[10]/[12]
621sh+624 (w)	621 (w)	ν (N-CH ₃) [ip]	[11]
641 (vw)		ρ (CH ₂), ρ (CH)	[11]
663 (m)	661 (vw)	ρ (CH ₂)	[11]
695 (w)	697 (vw)	ω (C-H)	[11]
718 (vw)		CH ₂ rocking	[15]
	738 (vw)	CH def-vib	[15]
781 (w)		ν_s (CC), ring HCCH sym bend, NC(H)N sym [op]	[8]
	803 (vw)	Ring HCCH sym bend, NC(H)N bending [op]	[8]
829+841 (w)		CH def vib [op]	[15]
865 sh+872+884 (w,m)	868 (w)	NC(H)N bend/CCH bend	[13]


892 (vw)		CH def vib [op]/CH ₃ rocking vib	[15]
	905 (vw)	CH def vib [op]	[15]
976 (w)		v(CC), ring asym bend [ip]	[12]
1010+1018 (s)	1012 sh+1021 (m)	v(C-C)/ρ(C-H)	[11]
1029 (m)		CH ₃ (N) str, CH ₂ (N) str/ρ(C-H)	[12]/[11]
1039 (m)	1045 (w)	CH ₃ (N) str, CH ₂ (N) str/v(N-C)	[12]/[11]
1056 (vw)		v(N-C)/v(CC), ring CN asym ip str [ip]	[11]/[8]
1094 (w)	1094 sh (w)	v(CC), ring HCCH sym bend, ring sym bend [ip]	[12]
1107 (m)	1116 (w)	v(CC), ring HCCH sym bend, ring sym str [ip]	[12]
1123 (w)		v(CC)	[13]
1163 (vw)	1173 (vw)	ρ(CH)	[11]
1189 (vw)		ρ(CH)	[11]
1207 (vw)		γ(CH) [op], ρ(CH ₂), v(CC) [ip]	[17]
	1228 (vw)	v(C-N), ρ(N-H)	[11]
1277 (m)	1272 (vw)	ρ(C-H) [ip], t(CH ₂) [ip]	[17]
1291 (vw)	1295 (vw)	t(CH ₂) [op], γ(CH)	[17]
1307 (w)		t(CH ₂) [op], γ(CH)	[17]
1333 (m)	1336 (m)	Ring sym str [ip], CC str, CH ₂ (N) and CH ₃ (N) CN str	[8]
1346 (m)		Ring sym str [ip], CC str, CH ₂ (N) and CH ₃ (N) CN str	[8]
1358 (w)		ρ(CH ₂), ρ(CH)/CH ₂ (N) str, v(CC), CH ₂ (N)CN str	[11]/[12]
1365 (w)		ρ(CH ₂), ρ(CH)	[11]
1382+ 1386 (m)	1385 (m)	ρ(CH ₂), ρ(CH)/δ(CH ₂) [op], ω(CH), ω(NH)	[11]/[17]
1410 sh+1415+1428 (m,s,vs)	1416 (s)	CH ₂ (N)/CH ₃ (N)CN str	[13]
1439 (m)		δ(CH ₂)	[13]
1443 sh (m)		CH ₃ (N) HCH sym bend/δ(CH)	[8]/[11]
1460 (m)	1454 (m)	CCH HCC sym bend, CH ₃ (N)HCH sym bend	[12]
1467 (m)	1469 (m)	v(N-C), δ(CH ₂)	[11]
1557 sh+1565+1575 sh (vw,w,vw)	1564 (w)	v(N-C)/CH ₂ (N) and CH ₃ (N) CN str	[11]/[8]
2708 (vw)		CH def-vib [ip]	[15]
2755 (vw)	2748 (vw)	v _s (CH ₃)	[15]
2811 (vw)		v _s (CH ₃)/v _{as} (CH ₃)	[15]
2837 (vw)	2828 (w)	v _s (CH ₂)/v _s (CH ₃)	[13]/[15]
	2876 (s)	v _s (CH ₂)/CH ₃ HCH sym str/v _{as} (CH ₂)	[17]/[12]/[13]
2896 (m)	2898 (m)	v _s (CH ₃)/v _{as} (CH ₂), v _s (CH ₂)	[15]/[17]
2928 (m)		Ring CH ₃ HCH sym str	[12]
2948 (s)	2943 (vs)	v _{as} (CH ₂), v(CH)	[17]
	2972 sh (s)	CH ₂ HCH asym str, ethyl HCH asym str	[12]
2982+3000 sh (m)		v _{as} (CH ₃)/N-H str	[13]/[15]
3045 (w)		v(C-H)	[13]
	3063 (w)	v(C-H)	[13]
3087+3098 (w)		v(C-H)/CH ₃ (N)HCH asym Str	[11]/[13]
3136 (vw)	3138 (vw)	ring (HCCH and N-(C-H)-N) C-H str/v(C-H)	[16]/[13]
3156 (vw)		ring (HCCH and N-(C-H)-N) C-H str/v(C-H)/v(N-H)	[16]/[13]/[15]

1 3.2.1. The region 200-45 cm⁻¹:

2 In this region an important difference between the two spectra was observed. The spectrum of
3 the monocationic ionic liquid had a wideband with a single peak that characterizes the amorphous
4 phase, while the spectrum of dicationic ionic liquid showed several peaks, indicating in this case the

1 presence of a crystalline phase.

2 The slight swinging motion of the imidazolium ring, an absent phenomenon in the non-
3 aromatic ionic liquids must be expressed in the RAMAN spectra of both ionic liquids: a broad peak
4 at $\approx 57\text{ cm}^{-1}$ in the case of monocationic ionic liquid, which becomes relatively narrow at 86 cm^{-1} in
5 the case of dicationic, due here to the presence of the crystalline phase. Furthermore the results of
6 C. Penna et al. [14] indicate that the RAMAN spectrum must present other peaks attributed to the
7 mode linked to the rocking motion: 146 cm^{-1} and 189 cm^{-1} (the wavenumber 189 cm^{-1} from the
8 literature can also be attributed to twisting of the bond (N-C) or to rocking (CH_2) [11]). The number
9 and intensities of the peaks show the influence of the passage of a monocationic ionic liquid to a
10 dicationic ionic liquid. The peaks at 100 cm^{-1} and 127 cm^{-1} are attributed to intramolecular
11 interactions [14].




12

13 *Fig.5. The FT-RAMAN spectra of: [PrMIM⁺][Br] and ([M(CH₂)₃IM²⁺][2Br]) in the region 1000-*
14 *1045 cm⁻¹.*

15 The frequencies $\approx 57\text{ cm}^{-1}$, $\approx 68\text{ cm}^{-1}$ and $\approx 80\text{ cm}^{-1}$ are attributed to partial character of acoustic
16 excitations for a crystalline physical state (intermolecular vibrational modes) [14]. This character is
17 compatible with the fact that we found these vibrational modes regardless of the specific molecular
18 structure. Therefore, these modes are also independent of the alkyl chain length of the cation, either
19 be it an aromatic or a non-aromatic ionic liquid. On the other hand, these modes depend on the
20 strength of the cation-anion interactions. Penna et al. [14] studied two monocationic ionic liquids
21 $[\text{C}_6\text{C}_1\text{im}][\text{Tf}_2\text{N}]$ and $[\text{C}_6\text{C}_1\text{im}]\text{Br}$ in crystalline phase. When passing from $[\text{C}_6\text{C}_1\text{im}][\text{Tf}_2\text{N}]$ to

1 [C₆C₁im]Br, they noticed in this zone of low frequencies a wavenumber shift of about 10 cm⁻¹,
 2 which shows the influence of the nature of the anion. Moreover, for our dicationic ionic liquid
 3 which represents the crystalline phase and at the same time their cations, they contain two
 4 imidazolium rings. We also registered a wavenumber shift with a higher order than the previous,
 5 which indicates that the number of cycles in the imidazolium cation also has an influence on the
 6 modes of intermolecular vibrations.



7
 8 *Fig.6. The FT-RAMAN spectra of [PrMIM⁺][Br⁻] and ([M(CH₂)₃IM²⁺][2Br⁻]) in the region*
 9 *200-45 cm⁻¹.*

10 **3.2.2. The region 1000-200 cm⁻¹ :**

11 In this region, there were 14 peaks for the monocationic ionic liquid and 23 peaks for the
 12 dicationic, indicating, for the latter, a richer RAMAN spectrum in vibration modes.

13 While passing from the monocationic ionic liquid to the dicationic ionic liquid, we observed
 14 the appearance of several new peaks: at 256 and 267 cm⁻¹, attributed to rocking mode of the (N-C)
 15 bond [11], and at 299 cm⁻¹ attributed to modes CH₂(N) and CH₃(N)CH bend [8]. In addition several
 16 other more or less intense peaks: 429, 641, 718, 781, 829, 841, 892 and 976 cm⁻¹ were observed. At
 17 613 cm⁻¹ a peak was present with medium intensity assigned to ω(NH) mode, the stretching of
 18 (CH₂) or (CH₃(N)CN) [12].

19 We also noticed deconvolutions of peaks: 621 cm⁻¹ in the case of the monocationic which
 20 corresponded to a doublet 621 and 624 cm⁻¹ in the case of dicationic, is attributed to the vibration
 21 mode in the plane (stretching of the (N-CH₃) bond) [11]. The peak at 868 cm⁻¹ in the case of
 22 monocationic became a doublet for the dicationic: 872 and 884 cm⁻¹ with a shoulder at 865 cm⁻¹, all


1 assigned to vibrations NC(H)N bend and CCH bend [13].

2 We observed for the other peaks minor wavenumber shifts of the order of 2 cm^{-1} (from 316 cm^{-1}
3 1 for the mono to 317 cm^{-1} for the dicationic, from 661 cm^{-1} to 663 cm^{-1} and 697 to 695 cm^{-1}), the
4 peak at 418 cm^{-1} remained unchanged for mono and dicationic. Still there were two peaks with
5 remarkable wavenumber shifts (from 461 cm^{-1} to 475 cm^{-1} assigned to (C-H) stretching [15] and
6 from 601 cm^{-1} to 595 cm^{-1} assigned to $\omega(\text{N-C})$ [11]).

7 3.2.3. The region $1700\text{-}1000\text{ cm}^{-1}$:

8 In this region, we observed 30 peaks for the dicationic versus 15 peaks for the monocationic.
9 So also in this region, the spectrum of dicationic is richer in modes than the spectrum of
10 monocationic. New peaks appeared in the case of dicationic: two less intense peaks at 1056 cm^{-1}
11 and 1358 cm^{-1} , other peaks slightly more intense at 1029 , 1123 , 1189 , 1207 , 1307 , 1346 , 1365 , 1439
12 cm^{-1} and a shoulder at 1443 cm^{-1} , the assignments of these peaks are in table (3). Among these
13 peaks the most remarkable one is at 1346 cm^{-1} assigned to Ring sym str [ip], CC str, $\text{CH}_2(\text{N})$ and
14 $\text{CH}_3(\text{N})$ CN str [8].

15



16

17 *Fig.7. FT-RAMAN spectra of: $[\text{PrMIM}^+][\text{Br}]$ and $([\text{M}(\text{CH}_2)_3\text{IM}^2+][2\text{Br}])$*

18

in the region $1700\text{ -}1000\text{ cm}^{-1}$.

19 Deconvolutions of the peaks was also observed in this region. The peak at 1385 cm^{-1} for mono
20 became a doublet for the dicationic: 1382 and 1386 cm^{-1} . These peaks are attributed in the literature


1 to the CH bond vibration or the NH bond [8, 17]. The peak at 1416 cm^{-1} changed into a triplet: 1410
2 cm^{-1} , 1415 cm^{-1} and 1428 cm^{-1} . These peaks are assigned to the stretching vibration of $\text{CH}_2(\text{N})$ and
3 $\text{CH}_3(\text{N})\text{CN}$ [13]. Also a peak at 1564 cm^{-1} changed into a triplet: 1557 cm^{-1} , 1565 cm^{-1} and 1575
4 cm^{-1} . These frequencies are also attributed to stretching vibration of $\text{CH}_2(\text{N})$ and $\text{CH}_3(\text{N})\text{CN}$ [8]
5 and $\nu(\text{N}-\text{C})$ [11]. For the rest of the peaks, we observed more or less significant shifts of the order
6 of 2 cm^{-1} to 10 cm^{-1} .

7 It is also envisaged that the peaks of the RAMAN spectrum of the dicationic are narrower than
8 for the monocationic, which means that it has more order in the dicationic structure than the
9 monocationic.

10 **3.2.4. The region $3200\text{-}2700\text{ cm}^{-1}$:**

11 As in previous regions, in region $3200\text{-}2700\text{ cm}^{-1}$, the dicationic spectrum was richer in
12 vibration modes than the monocationic one, with 14 peaks versus 8 peaks observed. New peaks
13 were present at 2708 cm^{-1} (assigned to CH in-plane def-vib), 2811 cm^{-1} (assigned to sym CH_3 str,
14 asym CH_3 str), 2928 cm^{-1} (assigned to ring CH_3HCH sym str), 2982 cm^{-1} and 3000 cm^{-1} (assigned
15 to $\nu_{\text{as}}(\text{CH}_3)$), 3045 cm^{-1} , 3087 cm^{-1} and 3098 cm^{-1} (assigned to $\nu(\text{CH})$), and 3156 cm^{-1} (assigned to
16 the ring (HCCH and $\text{N}(\text{CH})\text{N}(\text{CH})\text{CH}$ str/ $\nu(\text{CH})$). In the literature 2982 cm^{-1} , 3000 cm^{-1} and 3156 cm^{-1}
17 are attributed to the stretching vibration of NH bond. For modes present in both spectra, we had a
18 frequency shift of the order of 2 cm^{-1} to 9 cm^{-1} .

19 Therefore, all spectral regions of the FT-RAMAN spectrum that we saw above show sensitivity
20 towards the passage of mono to di-cationic by more peaks in the case of dicationic and wavenumber
21 shifts.



1
2 *Fig.8. The FT-RAMAN spectra of: [PrMIM⁺] [Br⁻] (monocationic) and*
3 *([M(CH₂)₃IM²⁺][2Br⁻]) (dicationic), in the region 3200-2700 cm⁻¹.*

4 **4. Conclusion:**

5 This study showed that in passing of an imidazolium-based monocationic (IL) where the cation
6 is composed of a one imidazolium ring to a dicationic (IL) where the cation is composed of a two
7 imidazolium rings, there is a change of environment for ions, hence more or less freedom of
8 movement. This was expressed on the FT-RAMAN spectrum in a wide spectral region by the
9 appearance of new vibrational modes (37 new peaks), deconvolutions and wavenumber shifts,
10 which can serve as a signing of differentiation between these (ILs).

11 This change in the number of imidazolium rings in the cation also influences the intermolecular
12 (anion-cation interaction) and intramolecular vibration modes. Indeed in the spectral zone 200-45
13 cm⁻¹, representative of these modes, we could see: a broadband with only a single peak for the
14 monocationic, characteristic of an amorphous phase, versus several peaks for the dicationic,
15 characteristic of a crystalline phase. So the self-organization known to ionic liquids was manifested
16 in the case of dicationic. Therefore, this spectral zone is a signature of the physical state of the (IL),
17 than the change of the number of imidazolium rings in the cation.

18 We note that for the FTIR/ATR spectroscopy the clutter of ions masked certain vibration
19 modes for the dicationic. Yet, we had more or less important wavenumber shifts, peaks with
20 pronounced intensities in the case of dicationic such as the new peak at 1655 cm⁻¹, and a failover in
21 intensity from the monocationic and dicationic between the spectral regions 3200-2800 cm⁻¹ and
22 3500-3200 cm⁻¹. The number of vibrational modes in the spectral region 3500-3200 cm⁻¹ also

1 increased, which can be explained by an increase in the number of N-H bonds. Therefore, these
2 spectral regions are sensitive to the change of the number of imidazolium rings.

3 Such as the FT-RAMAN spectroscopy, we also have for FTIR/ATR spectroscopy an indicator
4 of the existence of the order for the dicationic. This indicator is represented by the peaks in the
5 spectrum of the dicationic being slightly narrower than the peaks in the spectrum of the
6 monocationic.

7 **Acknowledgment**

8 The authors thank the ATRST-DGRSDT for the financial support. We would like to thank Quentin
9 Arnould, technician of Walloon Agricultural Research Centre (CRA-W), who participated to FT-
10 RAMAN and FTIR/ATR measurements.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

1 **References:**

- 2 [1] J. M. Crosthwaite, M. J. Muldoon, J. K. Dixon, J. L. Anderson, and J. F. Brennecke, "Phase
3 transition and decomposition temperatures, heat capacities and viscosities of pyridinium ionic
4 liquids," *The Journal of Chemical Thermodynamics*, vol. 37, no. 6, pp. 559–568, 2005.
- 5 [2] J. D. Holbrey and K. R. Seddon, "The phase behaviour of 1-alkyl-3-methylimidazolium
6 tetrafluoroborates; ionic liquids and ionic liquid crystals," *Journal of the Chemical Society,
7 Dalton Transactions*, no. 13, pp. 2133–2140, 1999.
- 8 [3] R. Hagiwara and Y. Ito, "Room temperature ionic liquids of alkylimidazolium cations and
9 fluoroanions," *Journal of Fluorine Chemistry*, vol. 105, no. 2, pp. 221–227, 2000.
- 10 [4] F. Endres, M. Bukowski, R. Hempelmann, and H. Natter, "Electrodeposition of
11 nanocrystalline metals and alloys from ionic liquids," *Angewandte Chemie International
12 Edition*, vol. 42, no. 29, pp. 3428–3430, 2003.
- 13 [5] H. Ohno, *Electrochemical aspects of ionic liquids*. John Wiley & Sons, 2011.
- 14 [6] M. Freemantle, *An introduction to ionic liquids*. Royal Society of chemistry, 2010.
- 15 [7] A. J. Walker and N. C. Bruce, "Cofactor-dependent enzyme catalysis in functionalized ionic
16 solvents," *Chemical communications*, no. 22, pp. 2570–2571, 2004.
- 17 [8] K. Noack, P. S. Schulz, N. Paape, J. Kiefer, P. Wasserscheid, and A. Leipertz, "The role of the
18 C2 position in interionic interactions of imidazolium based ionic liquids: a vibrational and
19 NMR spectroscopic study," *Physical Chemistry Chemical Physics*, vol. 12, no. 42, pp. 14153–
20 14161, 2010.
- 21 [9] A. M. Moschovi, S. Ntais, V. Dracopoulos, and V. Nikolakis, "Vibrational spectroscopic study
22 of the protic ionic liquid 1-H-3-methylimidazolium bis (trifluoromethanesulfonyl) imide,"
23 *Vibrational Spectroscopy*, vol. 63, pp. 350–359, 2012.
- 24 [10] T. Moumene, E. H. Belarbi, B. Haddad, D. Villemin, O. Abbas, B. Khelifa, and S. Bresson,
25 "Vibrational spectroscopic study of ionic liquids: Comparison between monocationic and
26 dicationic imidazolium ionic liquids," *Journal of Molecular Structure*, vol. 1065, pp. 86–92,
27 2014.
- 28 [11] K. Malek, A. Puc, G. Schroeder, V. I. Rybachenko, and L. M. Proniewicz, "FT-IR and FT-
29 Raman spectroscopies and DFT modelling of benzimidazolium salts," *Chemical physics*, vol.
30 327, no. 2, pp. 439–451, 2006.
- 31 [12] J. Kiefer, J. Fries, and A. Leipertz, "Experimental vibrational study of imidazolium-based
32 ionic liquids: Raman and infrared spectra of 1-ethyl-3-methylimidazolium bis
33 (trifluoromethylsulfonyl) imide and 1-ethyl-3-methylimidazolium ethylsulfate," *Applied
34 spectroscopy*, vol. 61, no. 12, pp. 1306–1311, 2007.
- 35 [13] T. Moumene, E. H. Belarbi, B. Haddad, D. Villemin, O. Abbas, B. Khelifa, and S. Bresson,
36 "Study of imidazolium dicationic ionic liquids by Raman and FTIR spectroscopies: The effect
37 of the nature of the anion," *Journal of Molecular Structure*, vol. 1083, pp. 179–186, 2015.
- 38 [14] T. C. Penna, L. F. Faria, J. R. Matos, and M. C. Ribeiro, "Pressure and temperature effects on
39 intermolecular vibrational dynamics of ionic liquids," *The Journal of chemical physics*, vol.
40 138, no. 10, p. 104503, 2013.
- 41 [15] G. Socrates, *Infrared and Raman characteristic group frequencies: tables and charts*. John
42 Wiley & Sons, 2004.
- 43 [16] M. Koel, *Ionic liquids in chemical analysis*. CRC Press, 2008.
- 44 [17] K. Malek, E. Podstawka, J. Milecki, G. Schroeder, and L. M. Proniewicz, "Structural features
45 of the adenosine conjugate in means of vibrational spectroscopy and DFT," *Biophysical
46 chemistry*, vol. 142, no. 1, pp. 17–26, 2009.

1 [18]S. Bresson, D. Rousseau, S. Ghosh, M. E. Marssi, and V. Faivre, “Raman spectroscopy of the
2 polymorphic forms and liquid state of cocoa butter,” *European Journal of Lipid Science and*
3 *Technology*, vol. 113, no. 8, pp. 992–1004, 2011.

4

5

6