

242 m μ , log ϵ 4.23). Mild alkaline hydrolysis of V, VI and VII afforded 6 α -fluoro compound "S" (VII) (m.p. 203–205°, [α]_D +135°, λ_{\max} 236 m μ , log ϵ 4.21), 6 β -fluoro compound "S" (IX) (m.p. 222–224°, [α]_D +14°, λ_{\max} 234 m μ , log ϵ 4.09) and Δ^1 -dehydro-6 α -fluoro compound "S" (X) (m.p. 210–212°, [α]_D +66°, λ_{\max} 241 m μ , log ϵ 4.22), respectively.

Adrenal incubation⁸ of both VIII and X followed by monoacetylation at C-21 gave, in good yield, 6 α -fluorohydrocortisone acetate (XI) (m.p. 215–217°, [α]_D +149° (diox.), λ_{\max} 237 m μ , log ϵ 4.23) and 6 α -fluoroprednisolone acetate (XII) (m.p. 235–237°, [α]_D +114° (diox.), λ_{\max} 242 m μ , log ϵ 4.25), respectively. Selenium dioxide oxidation⁹ of XI gave XII. Chromic acid oxidation of XI and XII provided 6 α -fluorocortisone acetate (XIII) (m.p. 215–216°, [α]_D +190°, λ_{\max} 233 m μ , log ϵ 4.21) and 6 α -fluoroprednisone acetate (XIV) (m.p. 228–230°, [α]_D +142° (diox.), λ_{\max} 237 m μ , log ϵ 4.23).

An alternative completely chemical route to the 6-fluoro corticoids proceeded from cortisone acetate 3-monoketal (XV).⁷ Acetylation of XV at C-17 and peracid epoxidation of the resulting 17,21-diacetate (XVI) (m.p. 180–182°, [α]_D –20°) afforded the 5 α ,6 α -epoxide (XVII) (m.p. 233–234°, [α]_D –18°). Boron trifluoride opening of XVII gave the corresponding 5 α -hydroxy-6 β -fluoro-3-ketal (XVIII) (m.p. 231–232°, [α]_D –9°) which was directly converted by acid treatment to 6 α -fluorocortisone 17,21-diacetate (XIX) (m.p. 270–272°, [α]_D +108°, λ_{\max} 233 m μ , log ϵ 4.20). Alkaline hydrolysis of XIX and subsequent monoacetylation gave XIII. Milder acid treatment of XVIII afforded 6 β -fluorocortisone 17,21-diacetate (XX) (m.p. 212–213°, [α]_D +62°, λ_{\max} 230 m μ , log ϵ 4.09).

In the anti-inflammatory and thymolytic assays,⁸ 6 α -fluorocortisone acetate (XIII), 6 α -fluorohydrocortisone acetate (XI), 6 α -fluoroprednisolone acetate (XII) and 6 α -fluoroprednisone acetate (XIV) had respective activities of (10,6), (10,8), (20,62), (20,23) times that of hydrocortisone acetate. In the adrenalectomized nonsalt loaded rat XI, XII and XIV promoted sodium excretion.⁹

The details of this work together with the syntheses of other 6-fluoro steroid hormones^{4b,10} will be described in future publications.

RESEARCH LABORATORIES
SYNTEX, S. A.
APDO. POSTAL 2679
MEXICO, D. F.

A. BOWERS
H. J. RINGOLD

RECEIVED JUNE 23, 1958

(6) A. Zaffaroni, H. J. Ringold, G. Rosenkranz, F. Sondheimer, G. H. Thomas and C. Djerassi, *THIS JOURNAL*, **80**, October (1958).

(7) R. Antonucci, S. Bernstein, M. Heller, R. Lenhard, R. Littell and J. H. Williams, *J. Org. Chem.*, **18**, 70 (1953).

(8) Anti-inflammatory (cotton pellet implant) and thymolytic activity in adrenalectomized rat, oral route, hydrocortisone acetate = 1. We wish to thank Dr. R. Dorfman of the Worcester Foundation for these assays.

(9) Mineral assays by Dr. R. Dorfman and the Endocrine Labs., Madison, Wisconsin.

(10) By a similar series of reactions there was prepared 6 α -fluoro-17 α -acetoxy-progesterone (m.p. 249–250°, [α]_D +56°, λ_{\max} 236 m μ , log ϵ 4.21) and 1-dehydro-6 α -fluoro-17 α -acetoxyprogesterone (m.p. 258–261°, [α]_D +24°, λ_{\max} 241 m μ , log ϵ 4.18) which possess high progestational activity. Thus the former in the Clauberg assay exhibits oral activity equal to 17 α -ethynyl-19-nortestosterone (Norlutin).

TRANSPARENT BaFe₁₂O₁₉ AND SrFe₁₂O₁₉

Sir:

By a solid state reaction of barium fluoride and ferric oxide in the ratio 1:6, under oxygen, a red, transparent, single crystalline form of BaFe₁₂O₁₉ was obtained. The hexagonal platelets, approximately 50 microns thick and up to 2 mm. in diameter, are perfectly suitable for the observation and study of Weiss domains, which in the virgin state are about 1 micron wide. As an alternate technique, the reaction of ferric oxide in molten barium chloride or strontium chloride under oxygen, also yielded transparent platelets of BaFe₁₂O₁₉ and SrFe₁₂O₁₉. The compositions of the reaction products were established by chemical analysis and single crystal X-ray studies. An apparently similar form of BaFe₁₂O₁₉ was described recently by C. Kooy,¹ without indicating the method of preparation.

(1) C. Kooy, *Philips Technical Review*, **19**, 286 (1958).

RESEARCH DIVISION
PIGMENTS DEPARTMENT
E. I. DU PONT DE NEMOURS AND COMPANY, INC.
WILMINGTON 98, DELAWARE

LOTHAR H. BRINER

RECEIVED JULY 16, 1958

THE RADIOLYTIC SYNTHESIS OF THE *cis*- AND *trans*-ISOMERS OF 1,2-DICHLOROETHYLENE OXIDE

Sir:

In the course of some investigations on the radiation chemistry of the symmetrical dichloroethylenes, the *cis* and *trans* forms of 1,2-dichloroethylene oxide have been produced. These compounds have not been reported previously, and in view of the current interest in radiation utilization, we are presenting preliminary data on the radiolytic synthesis and physical properties of these compounds.

The *cis* and *trans* forms of 1,2-dichloroethylene were irradiated in glass cells *in vacuo* with 40-Mev. helium ions impinging on the liquid. After irradiation, the low-boiling products plus the bulk of the 1,2-dichloroethylene were stored in glass-stoppered bottles in the presence of air after preliminary analysis for high-boiling products. In samples that had stood several months, two peaks were found in GLP chromatograms (nonyl phthalate column) that were not present in samples chromatographed immediately after irradiation. These peaks were concentrated in the pot liquid by distillation through a small Vigreux column and were further concentrated and purified by running repeat GLP chromatograms of the pot liquid, collecting the respective peaks each time. By this means samples of 1 g. and 0.2 g. of the two respective peak materials were isolated in relatively pure form. These are referred to as dichloroethylene oxide I and dichloroethylene oxide II according to their respective GLP chromatographic emergence times.

The relative yield of oxide I to oxide II from *trans*-1,2-dichloroethylene was about 4.6:1. The ratio from irradiated *cis*-1,2-dichloroethylene was smaller but has not been well determined because of the much lower yield from this isomer.

The compounds isolated have been characterized as the *cis* and *trans* isomers of 1,2-dichloroethylene

oxide by mass-spectrometer spectral patterns and by comparison of the infrared spectra with the spectra of the *cis* and *trans* forms of 2,3-epoxybutane. The mass spectra, with parent masses at 112 to 117 mass units, give the empirical formula $C_2H_2Cl_2O$ from isotope ratio arguments. Chemical analysis for C, H, and Cl yielded 21.2%, 1.97%, and 63.5% for oxide I; and 21.2%, 1.92%, and 63.6% for oxide II. The calculated values for $C_2H_2Cl_2O$ are 21.27%, 1.78%, and 62.78%.

Known compounds of this composition, dichloroacetaldehyde and chloroacetyl chloride, were shown to differ from oxide I and oxide II in both mass spectrometer ionization patterns and GLP chromatographic retention times.

The properties, b.p., m.p., d^{25}_4 and n^{25}_D found for these compounds are oxide I: 80–81°, –43°, 1.405 and 1.438; oxide II, 111–112°, –63°, 1.470 and 1.452. From present information it has not been possible to decide unequivocally which compound is the *cis*-isomer and which is the *trans*-isomer.

At present one can only speculate as to the origin of these compounds. Yields of oxide I as high as $G = 9$ were found in old samples of irradiated *trans* 1,2-dichloroethylene. As this is as large as the total yield of "polymer" determined immediately after irradiation, a chain reaction for the formation of the oxides is indicated. An attractive possibility is an auto-oxidation reaction using a long-lived free radical present in very low concentration as the intermediate. Preliminary tests with other sources of radicals (benzoyl peroxide) in the presence of oxygen showed that small yields of both the dichloroethylene oxides are formed from both *cis*- and *trans*-1,2-dichloroethylenes.

This work was performed under the auspices of the United States Atomic Energy Commission.

RADIATION LABORATORY AND
DEPARTMENT OF CHEMISTRY
RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA

JEAN H. FUTRELL
AMOS S. NEWTON

RECEIVED JUNE 30, 1958

SIMPLE METAL ORBITAL TREATMENT OF THE BINDING OF THE HYDROGEN ATOM IN COBALT CARBONYL HYDRIDE

Sir:

On the basis of nuclear resonance and acid dissociation constants, Cotton and Wilkinson¹ have suggested that protons in $HCo(CO)_4$ and $H_2Fe(CO)_4$ are physically close to and primarily bound to the metal atoms. Liehr² has discussed which metal orbitals might be used in bonding. Edgell and Gallup^{3,4} consider the proton in $HCo(CO)_4$ to be placed so far from the Co atom (~ 2.0 Å.) that it is primarily bound to the carbon $p\pi$ orbitals. This view seems to be based on observed splitting of CO stretching frequencies while corresponding Co–C modes appear unsplit by the proton (according to their assignments). Since such reasoning seems

(1) F. A. Cotton and G. Wilkinson, *Chemistry and Industry (London)*, 1305 (1956).

(2) A. D. Liehr, *Z. Naturforsch.*, **12b**, 95 (1957).

(3) W. F. Edgell, C. Magee and G. Gallup, *THIS JOURNAL*, **78**, 4185 (1956).

(4) W. F. Edgell and G. Gallup, *ibid.*, **78**, 4188 (1956).




Fig. 1.

in principle valid, we note that Stammreich⁵ has recently observed distinct splittings of *all* triply degenerate vibrations in going from $Fe(CO)_4^{-2}$ to $HFe(CO)_4^{-}$.

In an effort to substantiate their view, Edgell and Gallup⁴ reported that a calculation of overlaps shows "that the hydrogen overlap integrals all have a maximum at a Co–H distance just short of 2 Å. . . ." and they call attention to "the relatively large value of the $\phi_{H\phi C}$ overlap and the relatively small value of the $\phi_{H\phi Co}$ overlap." We question whether this type of calculation is as meaningful as they suggest, but granting that it is, there appear to be large numerical errors in their results. Repeating this calculation [(1) placing the hydrogen atom on a C_3 axis of the $Co(CO)_4$ tetrahedron and computing overlaps over the d_{Co-H} range 0.2 to 3.0 Å.; (2) taking $S(Co-H)$ as $S(1s-3d_{z^2})$; (3) computing for $S(C-H)$ the overlap of the H 1s orbital with a simple LCAO-MO of A_1 symmetry, *viz.*, $1/\sqrt{3}[p\pi_{C1} + p\pi_{C2} + p\pi_{C3}]$; (4) deriving the master formula for $S(1s-3d_{z^2})$ and using published tables of integrals for Slater orbitals⁷] the results shown in the figure were obtained.

Features of principal interest are: (1) Both the Co–H and C–H overlaps are of comparable magnitude. (2) The C–H overlap maximizes at ~ 1.4 Å. while the Co–H overlap maximizes at ~ 0.9 Å. (3) The greatest total overlap ($\sim 40\%$ Co–H and $\sim 60\%$ C–H) occurs at $d_{Co-H} \approx 1.2$ Å. The relative magnitudes of $S(Co-H)$ and $S(C-H)$ could be further altered in favor of $S(Co-H)$ by including overlap in calculating the normalizing factor for the carbon $p\pi$ MO and by calculating $S(Co-H)$ using for Co a $3d_{z^2}-4p_z$ hybrid orbital. However, taking the results as they are, they suggest that Co–H bonding is quite important and that the hydrogen atom lies within 1.2 Å. of the Co atom. Since the sum of the Co and H radii is 1.2–1.4 Å., it appears not unjustified to describe the proton as somewhat

(5) H. Stammreich, private communication, to be published shortly.

(6) Note that when $d_{Co-H} = 2.0$ Å., $d_{C-H} = \sim 2.2$ Å., *i.e.*, about twice the normal C–H bond distance; yet a total C–H overlap of ~ 0.64 was reported.⁴ It was the improbability of this result which led to the present recalculation.

(7) (a) M. Kotani, A. Amemiya, E. Ishiguro and T. Kimura, *Tables of Molecular Integrals*, Maruzen Co., Tokyo, Japan (1955); (b) R. S. Mulliken, C. A. Rieke, D. Orloff and H. Orloff, *J. Chem. Phys.*, **17**, 1248 (1949).