CHEMICAL COMMUNICATIONS, 1971

Ring Contraction of 1-Alkyl-1,2,4-triazinium Iodides to 1,2,4-Triazoles


By JAEKEUN LEE and WILLIAM W. PAUDLER* (Department of Chemistry, Ohio University, Athens, Ohio 45701)

Summary 1-Alkyl-1,2,4-triazinium iodides readily undergo ring contraction to 1,2,4-triazoles on treatment with base.

It has been shown that 1,2,4-triazines undergo the Diels-Alder reaction,^{1,2} readily form covalent hydrates across the N(4)-C(5) bond,³ and easily dimerize.⁴

If polar substituents such as a 3-oxo-group are present, a 5-alkyl substituent, with a proton on the carbon atom fused to the 1,2,4-triazine ring, readily tautomerizes to the 5-alkylidene isomer.⁵ 1,2,4-Triazin-3(2H)-ones, when treated with hydroxylamine-O-sulphonic acid in base, ring contract to imidazolin-2-ones,⁶ and to 1,2,3-triazoles when treated with chloramine.⁷

Other reported ring-contractions of 1,2,4-triazines involve the conversion of 2-amino-5-phenyl-1,2,4-triazine into 3acetamido-5-methyl-1,2,4-triazole on treatment with acetic anhydride, the base-catalysed generation of 4-phenyl-1,2,3triazole from 3-amino-5-phenyl-1,2,4-triazine 2-oxide, and



SCHEME 1


the acid-catalysed conversion of 3-acetamido-5-phenyl-1,2,4-triazine 2-oxide into 4-phenylimidazol-2(3H)-one.8

We now describe another unusual reaction of the 1,2,4triazine ring system.

Hydrolysis of N^1 -alkylated 1,2,4-triazines (1) under basic conditions might be expected to give the carbinolamine (2). In fact a minor product is the 'oxidized carbinolamine' (3)when C(6) is unsubstituted, but the major products (ca. 65%) are the ring-contracted 1,2,4-triazole(4a-f). Structural proof for these 1,2,4-triazoles is based upon correct molecular weights (mass spectroscopy), ¹H n.m.r. spectra, elemental analyses, and comparison with an authentic sample for compound (4c).9

- ¹ H. Neunhoeffer and N. H. Frauhauf, *Tetrahedron Letters*, 1969, 3151.
 ² J. A. Elix, W. S. Wilson, and R. N. Warrener, *Tetrahedron Letters*, 1970, 1387.
 ⁸ W. W. Paudler and T. K. Chen, *J. Heterocyclic Chem.*, 1970, 767.
 ⁴ T. K. Chen, Ph.D. thesis, Ohio University, 1971.
 ⁵ W. W. Paudler and I. K. Lorg, *Chem.*, in the press.

- ⁶ W. W. Paudler and J. Lee, J. Org. Chem., in the press.
 ⁶ C. W. Rees and A. A. Sale, Chem. Comm., 1971, 531.
 ⁸ C. W. Rees and A. A. Sale, Chem. Comm., 1971, 532.
 ⁸ T. Sasaki and K. Minamoto, J. Org. Chem., 1966, 31, 3917.
 ⁹ M. R. Aikinson and J. B. Polya, J. Chem. Soc., 1954, 3319.



This new ring contraction may occur via the carbinolamine (2) as in Scheme 2.

(Received, September 27th, 1971; Com. 1685.)