

Benzoheterocycles *via* Aryne C-C Cyclisation: 2-Methylisindole

By BRIAN JAKES* and RAYMOND G. WALLACE
(School of Pharmacy, Portsmouth Polytechnic, Portsmouth PO1 2DZ)

Summary 2-Methylisindole is obtained in high yield by the reaction of *N*-methyl-2-chlorobenzylaminoacetonitrile with potassamide in liquid ammonia.

METHODS used for the synthesis of isindoles have recently been reviewed.¹ Most of those which involve simple stages rely upon prior formation and isolation of an isindoline or a related cyclic compound, elimination from an isindoline with the leaving group on nitrogen being an example.^{1,2}

Treatment of *N*-methyl-2-chlorobenzylaminoacetonitrile† in liquid ammonia with potassamide (4 equiv.) over 0.5 h and quenching with ammonium nitrate gave 2-methylisindole (89%). The m.p.,³⁻⁵ u.v.,⁶ and n.m.r. spectra, and the 1,3,5-trinitrobenzene adduct were consistent with 2-methylisindole.

Its identity was further confirmed by its reduction (NaBH₄-EtOH) to 2-methylisindoline (identified by i.r.,⁴ n.m.r., and analysis). Reduction was only efficient with an acid (HCl) work-up, suggesting that the reduction is actually of a protonated species.‡

In the formation of 2-methylisindole it is considered that cyclisation to 1-cyano-2-methylisindoline occurs first followed by hydrogen cyanide elimination to yield the isindole. It is not known whether cyanide is eliminated from an anion following proton abstraction from C-1 and proton shift (or direct abstraction from C-3) or is eliminated first to give a cationic isindolenine followed by proton loss and rearrangement.

(Received, 12th January 1972; Com. 043.)

† Prepared by treating 2-chlorobenzyl chloride with methylaminoacetonitrile.

‡ Although acetic acid work-up of borohydride mixtures has been shown to cause generation of diborane (J. A. Marshall and W. S. Johnson, *J. Amer. Chem. Soc.*, 1963, **28**, 421, 595), because of the presence of ethanol and water in the reaction mixture we favour the suggestion made in the text.

¹ J. D. White and M. E. Mann, *Adv. Heterocyclic Chem.*, 1969, **10**, 113.

² J. E. Shields and J. Bornstein, *Chem. and Ind.*, 1967, 1404; J. E. Shields and J. Bornstein, *J. Amer. Chem. Soc.*, 1969, **91**, 5192; J. Bornstein, D. A. McGowan, A. L. Di Salvo, J. E. Shields, and J. Kopecky, *Chem. Comm.*, 1971, 1503.

³ J. Thesing, W. Schäfer, and L. D. Melchior, *Annalen*, 1964, **671**, 119.

⁴ G. Wittig and H. Streib, *Annalen*, 1953, **584**, 1.

⁵ G. Wittig, H. Tenhaeff, W. Schoch, and G. Koenig, *Annalen*, 1951, **572**, 1.

⁶ J. Schmutz and F. Kunzle, *Helv. Chim. Acta*, 1956, **39**, 1144.