

Triclinic

$P\bar{1}$
 $a = 9.3876(1) \text{ \AA}$
 $b = 12.4039(2) \text{ \AA}$
 $c = 12.7550(3) \text{ \AA}$
 $\alpha = 104.807(1)^\circ$
 $\beta = 99.431(1)^\circ$
 $\gamma = 110.927(1)^\circ$
 $V = 1286.54(4) \text{ \AA}^3$
 $Z = 2$
 $D_x = 1.289 \text{ Mg m}^{-3}$
 D_m not measured

Data collection

Siemens SMART CCD area-detector diffractometer
 ω scans
 Absorption correction: empirical (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.915$, $T_{\max} = 0.972$
 9032 measured reflections

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.075$
 $wR(F^2) = 0.176$
 $S = 1.074$
 5102 reflections
 314 parameters
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0574P)^2 + 0.426P]$
 where $P = (F_o^2 + 2F_c^2)/3$

Cell parameters from 5053 reflections
 $\theta = 1.72\text{--}27.05^\circ$
 $\mu = 0.282 \text{ mm}^{-1}$
 $T = 293(2) \text{ K}$
 Parallelepiped
 $0.32 \times 0.22 \times 0.10 \text{ mm}$
 Yellow

5102 independent reflections
 3148 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$
 $\theta_{\max} = 27.05^\circ$
 $h = -11 \rightarrow 10$
 $k = -15 \rightarrow 15$
 $l = 0 \rightarrow 16$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.376 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.273 \text{ e \AA}^{-3}$
 Extinction correction: SHELXTL
 Extinction coefficient: 0.008(3)
 Scattering factors from International Tables for Crystallography (Vol. C)

Data collection: SMART (Siemens, 1996). Cell refinement: SAINT (Siemens, 1996). Data reduction: SAINT. Program(s) used to solve structure: SHELXTL (Sheldrick, 1997). Program(s) used to refine structure: SHELXTL. Molecular graphics: SHELXTL. Software used to prepare material for publication: SHELXTL and PARST (Nardelli, 1995).

The authors would like to thank the Malaysian Government for research grant R&D No. 190-9609-2801. This work was also supported by a grant for the Key Research Project from the State Science and Technology Commission and the National Nature Science Foundation of China. KC thanks the Universiti Sains Malaysia for a Visiting Postdoctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: HA1233). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Chapman, R. L., Stephens, F. S. & Vagg, R. S. (1981). *Inorg. Chim. Acta*, **52**, 161–168.
 Ko, P. H., Chen, T. Y., Zhu, J., Cheng, K. F., Peng, S. M. & Che, C. M. (1995). *J. Chem. Soc. Dalton Trans.* pp. 2215–2219.
 Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
 Sheldrick, G. M. (1996). *SADABS. Siemens Area Detector Absorption Correction Software*. University of Göttingen, Germany.
 Sheldrick, G. M. (1997). *SHELXTL. Structure Determination Software Programs*. Version 5.10. Bruker Analytical X-ray Systems, Madison, Wisconsin, USA.
 Siemens (1996). *SMART and SAINT. Area Detector Control and Integration Software*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 100–102

1-Isobutyl-2,6,6-trimethyl-4-oxo-4,5,6,7-tetrahydro-3-indoleacetic acid

S. DEEPTHI,^a VASANTHA PATTABHI^a AND K. NAGARAJAN^b

^aDepartment of Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and ^bRecon Limited, Bangalore 560 076, India. E-mail: crystal@giasmd01.vsnl.net.in

(Received 23 February 1998; accepted 30 July 1998)

Abstract

The title compound (C₁₇H₂₅NO₃) is a perhydro-indoleacetic acid which exhibits hypoglycemic activity. The cyclohexyl ring adopts a sofa conformation. The molecule contains an intramolecular O—H...O hydrogen bond.

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

O1—C10	1.213 (4)	N2—C11	1.401 (4)
O2—C17	1.213 (4)	N3—C17	1.357 (4)
N1—C7	1.333 (4)	N3—C16	1.416 (4)
N1—C8	1.341 (4)	N4—C22	1.331 (4)
N2—C10	1.350 (4)	N4—C18	1.337 (4)
C11—N2—C10—C8	177.3 (3)	C17—N3—C16—C15	51.7 (5)
N1—C8—C10—N2	−1.8 (5)	C16—N3—C17—C18	−178.6 (3)
C10—N2—C11—C12	6.0 (6)	N3—C17—C18—N4	−1.6 (4)

Table 2. Hydrogen-bonding geometry ($\text{\AA}$, $^\circ$)


D—H...A	D—H	H...A	D...A	D—H...A
C12—H12A...O1	0.93	2.276	2.880 (4)	122
C25—H25A...O1 ⁱ	0.96	2.501	3.449 (5)	170

Symmetry code: (i) $1 + x, y, 1 + z$.

The data collection covered over a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle (0, 88 and 180°) for the crystal and each exposure of 60 s covered 0.3° in ω . The crystal-to-detector distance was 3 cm and the detector swing angle was -20° . Coverage of the unique set is over 99% complete. Crystal decay was monitored by repeating 30 initial frames at the end of data collection and analysing the duplicate reflections, and was found to be negligible. The structure was solved by direct methods and refined by full-matrix least-squares techniques. All H atoms were geometrically fixed and allowed to ride on their attached atoms.

Comment

Non-insulin-dependent diabetes mellitus (NIDDM) represents a singular endocrinal disorder which is assuming alarming global proportions (Davis & Granner, 1996; Colca & Tannis, 1992). We had reported earlier that a series of 4-oxo-4,5,6,7-tetrahydro-3-indoleacetic acids displayed antidiabetic properties (Nagarajan *et al.*, 1988). One of the most potent compounds in this series was GO 9001, 1-isobutyl-2,6,6-trimethyl-4-oxo-4,5,6,7-tetrahydro-3-indoleacetic acid (Nagarajan *et al.*, 1989). Its structure (I) is reported here.



The five-membered ring and part of the cyclohexyl ring (C17, C6, C5, C12 and O1) form a delocalized system with the bonds having partial double-bond character. The five-membered ring is in the envelope conformation and C6 deviates significantly (4σ) from the plane through C5, C7, C8 and N1. It is clear from the torsion angles that the cyclohexyl ring is in the sofa conformation and the best plane through the atoms in the ring makes angles of $8.44(7)$ and $57.90(2)^\circ$ with the five-membered ring and carboxyl group, respectively. The carboxyl group attached to the five-membered ring is planar and is in a perpendicular conformation [$118.04(7)^\circ$] with respect to the five-membered ring.




Fig. 1. ZORTEP (Zsolnai, 1994) drawing of the title compound with 50% probability displacement ellipsoids and the atom-numbering scheme.

The isobutyl group of the five-membered ring is in *g* + *tg* conformation. There is an intramolecular hydrogen bond between O2—H...O1 [O2...O1 = $2.641(2)$, H...O1 = $1.71(2)$ Å, O2—H...O1 = $173(2)^\circ$] stabilizing the conformation of the molecule.

Experimental

The title compound was synthesized from dimedone in three steps: addition to maleic acid, treatment of the adduct with acetic anhydride and reaction of the resultant triketo acid with isobutylamine in acetic acid. Single crystals were obtained when the compound was dissolved in benzene and slowly evaporated. The compound melts at $427\text{--}429$ K.

Crystal data

$\text{C}_{17}\text{H}_{25}\text{NO}_3$
 $M_r = 291.38$
 Monoclinic
 $P2_1/c$
 $a = 9.233(5)$ Å
 $b = 12.952(3)$ Å
 $c = 13.943(5)$ Å
 $\beta = 105.93(1)^\circ$
 $V = 1603.3(11)$ Å³
 $Z = 4$
 $D_x = 1.207$ Mg m⁻³
 D_m not measured

Cu K α radiation
 $\lambda = 1.54180$ Å
 Cell parameters from 23 reflections
 $\theta = 11\text{--}30^\circ$
 $\mu = 0.656$ mm⁻¹
 $T = 293(2)$ K
 Rectangular
 $0.45 \times 0.2 \times 0.2$ mm
 Pale white

Data collection

Enraf-Nonius CAD-4 diffractometer
 $\omega/2\theta$ scans
 Absorption correction: none
 2351 measured reflections
 2101 independent reflections
 1812 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$
 $\theta_{\text{max}} = 71.86^\circ$
 $h = 0 \rightarrow 11$
 $k = 0 \rightarrow 15$
 $l = -17 \rightarrow 16$
 3 standard reflections
 frequency: 120 min
 intensity decay: <1%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.108$
 $S = 1.102$
 2101 reflections
 200 parameters
 H atoms: see below
 $w = 1/[\sigma^2(F_o^2) + (0.0546P)^2 + 0.3299P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.009$
 $\Delta\rho_{\text{max}} = 0.174$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.120$ e Å⁻³
 Extinction correction: SHELXL
 Extinction coefficient: 0.0188 (10)
 Scattering factors from International Tables for Crystallography (Vol. C)

Table 1. Selected geometric parameters (Å)

C5—N1	1.361 (2)	C6—C7	1.436 (2)
C5—C6	1.382 (3)	C7—C8	1.365 (2)
C5—C12	1.489 (3)	C8—N1	1.398 (3)
C6—C17	1.435 (2)	C17—O1	1.238 (3)

The H atom attached to O2 has been refined isotropically. The other H atoms have been treated as riding.

Data collection: CAD-4 Software (Enraf-Nonius, 1989). Cell refinement: SDP (Frenz, 1978). Data reduction: CAD-4

Software. Program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ZORTEP* (Zsolnai, 1994). Software used to prepare material for publication: *SHELXL93* and *PARST* (Nardelli, 1983).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: VJ1085). Services for accessing these data are described at the back of the journal.

References

- Colca, J. R. & Tannis, S. P. (1992). *Annu. Rep. Med. Chem.* **27**, 219–226.
- Davis, S. N. & Granner, D. K. (1996). In *The Pharmacological Basis of Therapeutics*, pp. 1493–1517. New York: McGraw-Hill.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Frenz, B. A. (1978). *The Enraf–Nonius CAD-4 SDP – a Real-Time System for Concurrent X-ray Data Collection and Crystal Structure Solution. Computing in Crystallography*, edited by H. Schenk, R. Olthof-Hazekamp, H. van Koningsveld & G. C. Bassi, pp. 64–71. Delft University Press.
- Nagarajan, K., Talwalker, P. K., Nagane Goud, A., Shah, R. K. & Shenoy, S. J. (1989). *Arzneim. Forsch.* **39**, 548–550.
- Nagarajan, K., Talwalker, P. K., Nagane Goud, A., Shah, R. K., Shenoy, S. J. & Desai, N. D. (1988). *Indian J. Chem.* **27B**, 1113–1123.
- Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXS97. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
- Zsolnai, L. (1994). *ZORTEP. An Interactive Graphics Program*. University of Heidelberg, Germany.

Acta Cryst. (1999). **C55**, 102–104

1,3-Bis(3-nitrophenyl)triazene

DE-CHUN ZHANG,^a ZHENG-HAO FEI,^b TIAN-ZHU ZHANG,^c YAN-QIU ZHANG^a AND KAI-BEI YU^d

^aDepartment of Chemistry, Suzhou University, Suzhou 215006, People's Republic of China, ^bDepartment of Chemistry, Yancheng Teacher's College, Yancheng 220004, People's Republic of China, ^cDepartment of Radiation Medicine, Suzhou Medicine College, Suzhou 215007, People's Republic of China, and ^dChendu Institute of Organic Chemistry, Chinese Academy of Sciences, Chendu 610041, People's Republic of China. E-mail: dc:hang@suda.edu.cn

(Received 1 May 1998; accepted 31 July 1998)


Abstract

The title molecule, C₁₂H₉N₃O₄, has *trans* geometry about the azo linkage. The dihedral angle between the two phenyl rings is 6.2(2)°. The whole molecule is

almost planar and the maximum deviation from the mean plane is 0.166(3) Å. The resonance effect in the triazene group which might explain the antitumour activity of some triazene derivatives is discussed. In the crystal structure, while the twofold screw-related molecules are packed in a simple herringbone pattern, the inversion-centre-related molecules form hydrogen-bonded dimers which are held together through $\pi \cdots \pi$ interactions along the [010] direction.

Comment

It has been reported that 1,3-diphenyltriazene shows antitumour activity comparable to that of 3,3-dimethyl-1-aryltriazenes. 1,3-Diaryltriazenes can act as pro-drugs, or 'carriers' of diazonium compounds, since they undergo hydrolysis to form aryl diazonium compounds (Nifontov *et al.*, 1988, 1994). Surprisingly few X-ray structural studies of these compounds have been reported. As part of the X-ray structural studies of substituted triazene compounds, we have synthesized the title compound, (I), and confirmed its structure by X-ray analysis.



Within the limits of accuracy, the bond lengths and angles of the two nitrophenyl rings are in agreement with each other, but the whole molecule is asymmetric. N3=N4 [1.261(2) Å] is longer than expected for a double bond, N2=N3 [1.326(2) Å] is shorter than expected for a single bond, and C5=N2 and C11=N4 are both short for single C_{ar}—N bonds; there is clear evidence of a resonance effect in this part of the molecule. In view of the carcinogenic activity of some azobenzene derivatives (Kurosaki *et al.*, 1976) closely related to their conjugation structure (Arcos & Arcos, 1962), we think that the resonance effect in the triazene group in triazene derivatives might be related to their antitumour activity. While the nitro groups O3=N5—O4 and O1=N1—O2 are twisted by 3.3(2) and 6.7(2)° from their benzene planes, the dihedral angle between the two phenyl rings is 6.2(2)°, and the torsion angles C5=N2=N3=N4 and N2=N3=N4—C11 are 179.4(1) and 179.6(1)°, respectively, showing the near coplanarity of the whole molecule. The shortest distance between the intermolecular mean planes is 3.27(3) Å.

In the crystal structure, the twofold screw-related molecules are packed in a simple herringbone pattern; interactions between them include the close contact O3 $\cdots$ N5 (Table 2). The molecules related by inversion centres are linked together to form dimers by O4 $\cdots$ H—