

A Novel Synthesis of 4-Propyl-2*H*-1-benzopyran-2-ones

Vinod K. Ahluwalia*, Rishi P. Singh, and Rama P. Tripathi

Department of Chemistry, University of Delhi, Delhi-110 007, India

(Received 10 October 1983. Accepted 24 November 1983)

Reaction of 2-hydroxybutyrophenones **1–5** with ethoxycarbonylmethylenetriphenylphosphorane furnishes the 4-propyl-2*H*-1-benzopyran-2-ones **6–10**.

(Keywords: Ethoxycarbonylmethylenetriphenylphosphorane; 2-Hydroxybutyrophenones; Insecticidal properties; 4-Propyl-2*H*-1-benzopyran-2-ones; Wittig reaction)

Eine neue Synthese von 4-Propyl-2H-1-benzopyran-2-onen


Die Reaktion von 2-Hydroxybutyrophenonen **1–5** ergab mit Ethoxycarbonylmethylenetriphenylphosphoran die 4-Propyl-2*H*-1-benzopyran-2-one **6–10**.

Introduction

A number of 4-propylcoumarins (4-propyl-2*H*-1-benzopyran-2-ones) have been isolated¹ from plants of various *Mammea species*. These coumarins are found to possess insecticidal properties^{2–4}. The structures of the natural 4-propylcoumarins have been established by spectroscopic methods and degradative studies. No considerable efforts have been made so far to synthesise them. The methods available for the synthesis of 4-propylcoumarins are (i) *Pechmann* condensation^{5–7} of phenols with ethyl butyryl acetate and (ii) reaction⁸ of β -diketones with $\text{HBr} - \text{Ac}_2\text{O}$. In general the yields reported by these procedures are not satisfactory. We report herein a general and efficient method for the synthesis of 4-propyl-2*H*-1-benzopyran-2-ones. It involves the *Wittig* reaction⁹ of 2-hydroxybutyrophenones **1–5** with ethoxycarbonylmethylenetriphenylphosphorane $[(\text{C}_6\text{H}_5)_3\text{P} = \text{CH} \cdot \text{COOC}_2\text{H}_5]$.

Results and Discussion

Initially the reaction of 2-hydroxy-4-methoxybutyrophenone¹⁰ (**1**) with ethoxycarbonylmethylenetriphenylphosphorane in anhyd. benzene gave only one product in excellent yield (80%). It was assigned the structure 7-methoxy-4-propyl-2*H*-1-benzopyran-2-one (**6**) on the basis of its elemental analysis, IR and ¹H-NMR spectral data. Using the same procedure different 4-propyl-2*H*-1-benzopyran-2-ones **6–10** have been synthesised starting from the appropriate 2-hydroxybutyrophenones **2–5**. The ketones 2-hydroxy-4-methoxy-3-methylbutyrophenone (**3**) and 2-hydroxy-4-methoxy-6-methylbutyrophenone (**5**) were obtained by *Friedel-Crafts* reaction of *n*-butyryl chloride with 1,3-dimethoxy-2-methylbenzene and 1,3-dimethoxy-5-methylbenzene, respectively, while **2** and **4** are known earlier¹⁰. The compounds 7-methoxy-8-methyl-4-propyl-2*H*-1-benzopyran-2-one (**8**) and 7-methoxy-5-methyl-4-propyl-2*H*-1-benzopyran-2-one (**10**) have been synthesised for the first time.



All these coumarins exhibited a C=O band in their IR spectra between 1710 and 1700 cm⁻¹. The ¹H-NMR spectral data confirmed the proposed structures.

Neutral conditions, superior yields and ready availability of the stable phosphorane render the present method attractive and enhance its utility.

Acknowledgement

The financial assistance from University Grants Commission, New Delhi, is acknowledged.

Experimental

All melting points are uncorrected. ^1H -NMR spectra (CDCl_3 ; *TMS*; δ/ppm) were recorded on a Perkin-Elmer R-32 (90 MHz) spectrometer. IR spectra (KBr ; cm^{-1}) were measured on a Perkin-Elmer Grating Infrared spectrophotometer, model 621. The experimental C, H values for compounds **3**, **5** and **6–10** were in full agreement with the given structures.

Preparation of 2-Hydroxy-4-methoxy-3-methylbutyrophenone (**3**)

General Procedure

To a cooled and stirred solution of 1,3-dimethoxy-3-methylbenzene (5 g, 0.0142 mol) in absolute diethyl ether (15 ml) and anhydrous aluminium chloride (11.2 g, 0.084 mol) was added butyryl chloride (4.5 g, 0.042 mol) over 1 h and stirring continued for a further 4 h. Working up of the reaction mixture gave **3** (4.8 g) which crystallised from benzene–petroleum ether as colourless needles, m.p. 65–66°.

Preparation of 7-methoxy-4-propyl-2*H*-1-benzopyran-2-one (**6**)

General Procedure

A mixture of **1**¹⁰ (1.95 g, 0.01 mol) and ethoxycarbonylmethylene-triphenylphosphorane (5.2 g, 0.015 mol) was refluxed in dry benzene (25 ml), according to the time given below. After evaporation of the solvent the residue was chromatographed over silica gel. Elution of the column with benzene–petroleum ether (3:1) gave **6** (1.7 g). It crystallised from benzene–petroleum ether as colourless prisms, m.p. 76–78°.

3: Yield 55.5%; m.p. 65–66°C; NMR: 0.91 (t, $J = 7$ Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.60–1.99 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 2.10 (s, 3 H, 3- CH_3); 2.88 (t, $J = 7$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 3.84 (s, 3 H, 4- OCH_3); 6.29 (d, $J = 9.5$ Hz, 1 H, H-5); 7.45 (d, $J = 9.5$ Hz, H-6); 13.02 (s, 1 H, 2-OH, D_2O exchangeable).

5: Yield 50%; m.p. 68–69°C; NMR: 0.95 (t, 7 Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.49–1.90 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 2.22 (s, 3 H, 6- CH_3); 2.90 (t, $J = 7$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 3.77 (s, 3 H, 4- OCH_3); 6.10 (d, $J = 2.5$ Hz, 1 H, H-3); 6.20 (d, $J = 2.5$ Hz, 1 H, H-5); 13.55 (s, 1 H, 2-OH, D_2O exchangeable).

6: React. time 35 h; yield 80%; m.p. 76–78°C (Lit.⁵ 145°); IR: 1710 ($\text{C}=\text{O}$), 1610 ($\text{C}=\text{C}$); NMR: 1.02 (t, $J = 7$ Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.39–1.78 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 2.60 (t, $J = 7$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 3.75 (s, 3 H, 7- OCH_3); 5.95 (s, 1 H, H-3); 6.62 (d, $J = 2.5$ Hz, 1 H, H-8); 6.69 (dd, $J = 9.5$ Hz, 2.5 Hz, 1 H, H-6); 7.34 (d, $J = 9.5$ Hz, 1 H, H-5).

7: React. time 34 h; yield 76%; m.p. 79–80°C (Lit.⁵ 95°); IR: 1705 ($\text{C}=\text{O}$), 1605 ($\text{C}=\text{C}$); NMR: 1.00 (t, $J = 7$ Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.33–1.77 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 2.60 (t, $J = 7$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 3.83 and 3.88 (each s, each 3 H, $2 \times \text{OCH}_3$); 5.95 (s, 1 H, H-3); 6.72 (d, $J = 9.5$ Hz, 1 H, H-6); 7.14 (d, $J = 9.5$ Hz, 1 H, H-5).

8: React. time 20 h; yield 80%; m.p. 115–116°C; IR: 1700 ($\text{C}=\text{O}$), 1595 ($\text{C}=\text{C}$); NMR: 1.00 (t, $J = 7$ Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.47–1.85 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 2.20 (s, 3 H, 8- CH_3); 2.61 (t, $J = 7$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 3.81 (s, 3 H, 7- OCH_3); 5.95 (s, 1 H, H-3); 6.68 (d, $J = 9.5$ Hz, 1 H, H-6); 7.29 (d, $J = 9.5$ Hz, 1 H, H-5).

9: React. time 35 h; yield 75%; m.p. 118–119°C (Lit.⁸ 117–118°); IR: 1700 ($\text{C}=\text{O}$), 1600 ($\text{C}=\text{C}$); NMR: 1.10 (t, $J = 7$ Hz, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_3$); 1.46–1.83 (m,

2 H, CH₂CH₂CH₃); 2.92 (t, $J = 7$ Hz, 2 H, CH₂CH₂CH₃); 3.88 and 3.92 (each s, each 3 H, $2 \times$ OCH₃); 5.98 (s, 1 H, H-3); 6.31 and 6.46 (each d, $J = 2.5$ Hz, each 1 H, H-6 and H-8).

10: React. time 30 h; yield 65%; m.p. 97–98 °C; IR: 1710 (C=O), 1600 (C=C); NMR: 1.00 (t, $J = 7$ Hz, 3 H, CH₂CH₂CH₃); 1.37–1.70 (m, 2 H, CH₂CH₂CH₃); 2.32 (s, 3 H, 5-CH₃); 2.72 (t, $J = 7$ Hz, 2 H, CH₂CH₂CH₃); 3.82 (s, 3 H, 7-OCH₃); 5.95 (s, 1 H, H-3); 6.44 and 6.64 (each d, $J = 2.5$ Hz, each 1 H, H-6 and H-8).

References

- ¹ Murray R. D. H., Naturally Occurring Plant Coumarins. Fortschr. Chem. Org. Naturst. **35**, 199 (1978).
- ² Djerassi C., Eisenbraun E. J., Finnegan R. A., Gilbert B., J. Org. Chem. **25**, 2164 (1960).
- ³ Joshi B. S., Kamat V. N., Govindachari T. R., Ganguly A. K., Tetrahedron **25**, 1453 (1969).
- ⁴ Crombie L., Games D. E., Haskins N. J., Reed G. F., Tetrahedron Lett. **1970**, 251.
- ⁵ Kotwani N. G., Sethna S. M., Advani C. D., Proc. Indian Acad. Sci. [A] **15**, 441 (1942).
- ⁶ Djerassi C., Eisenbraun E. J., Finnegan R. A., Gilbert B., J. Amer. Chem. Soc. **80**, 3686 (1958).
- ⁷ Finnegan R. A., J. Amer. Chem. Soc. **25**, 2169 (1960).
- ⁸ Ahluwalia V. K., Rani N., Indian J. Chem. [B] **15**, 1003 (1977).
- ⁹ Mali R. S., Yadav V. J., Synthesis **1977**, 464.
- ¹⁰ Ahluwalia V. K., Prakash C., Jolly R. S., Gazz. Chim. Ital. **109**, 641 (1979).