This article was downloaded by: [University of Arizona] On: 22 January 2013, At: 02:00 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

http://www.tandfonline.com/loi/uopp20

IMPROVED SYNTHESIS OF 3-AMINOFURAZAN-4-CARBOXYLIC ACID

Kevin G. Meyer^a

^a DowAgroSciences, LLC, 9330 Zionsville Road, Indianapolis, IN, 46268

Version of record first published: 21 Feb 2009.

To cite this article: Kevin G. Meyer (2004): IMPROVED SYNTHESIS OF 3-AMINOFURAZAN-4-CARBOXYLIC ACID, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 36:4, 361-362

To link to this article: <u>http://dx.doi.org/10.1080/00304940409458679</u>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

OPPI BRIEFS

IMPROVED SYNTHESIS OF 3-AMINOFURAZAN-4-CARBOXYLIC ACID

Submitted by Kevin G. Meyer

(03/30/04)

DowAgroSciences, LLC 9330 Zionsville Road Indianapolis, IN 46268

In the course of the synthesis of a variety of heterocyclic carboxylic acids for our studies on bioactive amides, a new route for 3-aminofurazan-4-carboxylic acid (2) was discovered. The advent of combinatorial chemistry and high-throughput screening has increased the need for a diverse pool of molecular building blocks that are used to construct libraries of biologically active compounds. Rare and unnatural β -amino acids such as 2 are useful molecules for these purposes.

 $\underbrace{\begin{array}{c} 0\\ MeO \end{array}}_{MeO} \underbrace{\begin{array}{c} CN\\ H_2O, 88\% \end{array}}_{I} \underbrace{\begin{array}{c} 0\\ MeO \end{array}}_{N} \underbrace{\begin{array}{c} 0\\ MeO \end{array}}_{N} \underbrace{\begin{array}{c} CN\\ N\\ H_2O, 39\% \end{array}}_{I} \underbrace{\begin{array}{c} NH_2OH \bullet HCI, KOH \\ H_2O, 39\% \end{array}}_{N} \underbrace{\begin{array}{c} 0\\ HO \end{array}}_{N} \underbrace{\begin{array}{c} 0\\ HO \end{array}}_{N} \underbrace{\begin{array}{c} 0\\ NH_2 \end{array}}_{N} \underbrace{\begin{array}{c} 0\\$

Amino acid 2 was obtained in two convenient steps from readily accessible materials and, although the overall yield is 34%, the chemistry provides a safer, more direct route than previous accounts in which 2 was isolated as an unexpected side-product in the basic hydrolysis of a poly-oxime nitrile,¹ of the methyl iminoester of 2,² and of 7-amino-4*H*-furazano[3,4c][1,2,6] thiadiazine 5,5-dioxide.³ Amino acid 2 was isolated as a fine white powder and has been prepared on a multi-gram scale.

EXPERIMENTAL SECTION

Melting points are uncorrected. All reagents were purchased and used without further purification. Proton and ¹³C NMR spectra were obtained on a Varian Gemini 300 spectrometer. Infrared spectra were obtained on a Bio-Rad spectrophotometer. Mass spectra were obtained on a Hewlett-Packard Model 5989A mass spectrometer using the electron impact-direct insertion probe (EI-DIP) and are reported as m/z. Microanalyses were performed by Midwest Microlab of Indianapolis, Indiana.

^{© 2004} by Organic Preparations and Procedures Inc.

Methyl 2-Cyano-2-(hydroxyimino)acetate (1): Acetic acid (11.5 mL, 0.200 mol) was added drop-wise to a 0°C suspension of methyl cyanoacetate (13.35 mL, 0.151 mol) and NaNO₂ (12.5 g, 0.181 mol) in H₂O (60 mL). The reaction was allowed to warm to ambient temperature and after 4 hr the mixture was cooled to 0°C and adjusted to pH = 1 with conc. HCl. The solution was extracted with EtOAc and the organic phase was dried (MgSO₄) and conc. *in vacuo* to give 17.0 g (88%) of a white solid, mp 118-119°C, *lit.*⁴ mp 120-122°C, which was used without further purification; [M]⁺ = 128. IR (KBr): 3388, 3238, 2848, 2231, 1740, 1450, 1310, 1069, 817, 762 cm⁻¹; ¹H NMR (300MHz, DMSO-*d*₆): δ 3.63 (s 3H) ppm; ¹³C NMR (300MHz, DMSO-*d*₆): δ 161.0, 126.5, 109.0, 53.9 ppm.

3-Aminofurazan-4-carboxylic Acid (2): To a solution of hydroxylamine hydrochloride (8.13 g, 0.117 mol) in H₂O (120 mL) was added solid KOH (13.1 g, 0.234 mol) portionwise, followed by a solution of **1** (15.0 g, 0.117 mol) in H₂O (30 mL). After 1.5 hr, additional KOH (13.1 g, 0.234 mol) was added to the orange-red solution portion-wise and the reaction heated to 80°C. After 2 hr, the mixture was cooled to 0°C and acidified to pH < 1 with conc. HCl. The resulting white precipitate was collected and dried by azeotropic removal of water with toluene (2x) to yield 5.90 g (39%) of a white solid, mp 220°C (dec.)⁵, *lit.*³ mp 200°C; [M]⁺ = 129. IR (KBr): 3444, 3332, 1746, 1620, 1510 cm⁻¹; ¹H NMR (300MHz, DMSO-*d*₆): δ 10.3 (1H, br s), 6.3 (3H, br s) ppm; ¹³C NMR (300MHz, DMSO-*d*₆): δ 161.3, 157.1, 143.3 ppm.

Anal. Calcd for C₃H₃N₃O₃: C, 27.92, H, 2.34, N, 32.55. Found: C, 28.17, H, 2.06, N, 32.42

REFERENCES

- 1. H. Wieland, Z. Kitasato, and S. Utzino, Ann., 478, 43, (1930).
- V. Tselinskii, S. F. Mel'mikova, S. V. Pirogov, and A. V. Sergievskii, Zh. Org. Khim., 35, 315 (1999).
- P. Fernández-Resa, P. Goya, R. Nieves, C. Ochoa, and M. Stud, *Heterocycles*, 20, 2351 (1983).
- 4. L. Cheng and D. A. Lightner, Synthesis, 46, (1999).
- 5. The compound begins to decompose at 200°C, but does not melt until 220°C.
