Trifluoromethyl-substituted Dehydrodiazepines and Cyanopyrroles from Azido-/Tetrazolo-pyridines

Richard A. Evans and Curt Wentrup*

Chemistry Department, The University of Queensland, Brisbane, Queensland 4072, Australia

1,2-Didehydro-1,3-diazepines **4D**, **5D**, **4,6D** and **5,6D** are identified as unique products of both photolysis and thermolysis of azido-/tetrazolo-pyridines; the ultimate thermolysis products are trifluoromethylpyrrolecarbonitriles (7–9, 11–13 and 16–18).

The thermal and photochemical ring expansion of 2-pyridylnitrenes^{1,2} can in principle be used to synthesise 1,3-diazepine derivatives,³ but the reaction has found little use, mainly because the starting materials, tetrazolo[1,5-*a*]pyridines, do not photolyse well under ordinary reaction conditions.

We report here a study of trifluoromethyl-substituted 2-pyridylnitrene precursors that photolyse very efficiently to 1,2-didehydro-1,3-diazepines (1,3-diazacyclohepta-1,2,4,6-tetraenes) (**4D**, **5D**, **4,6D** and **5,6D**).[†] We demonstrate the formation of distinct dehydrodiazepines linking different sets of 2-pyridylnitrenes (Schemes 1–3) under both photochemical and thermal conditions. The CF₃ group was chosen because of its proven pharmaceutical importance⁴ and because of its electron-withdrawing effect which, as shown below, facilitates tetrazole–azide isomerism and increases nitrene reactivity. Furthermore, the reluctance of the CF₃ group to undergo migration guarantees the uniqueness of the novel cyano(trifluoromethyl)pyrroles obtained as final thermal reaction products (Schemes 1–3).

The tetrazole or azide starting materials were synthesised from the appropriate 2-chloropyridines by two methods, (*i*) direct nucleophilic substitution by NaN₃ in dimethylformamide (DMF) at 70 °C (**4T**, **5T**, **3T**, **6A**, **3,5T** and **4,5T**) and (*ii*) two-step reaction by diazotization of 2-hydrazinopyridines, obtained from the 2-chloropyridines with hydrazine hydrate in ethanol (reflux, 2–3 h) (**4T**, **5T**, **3T**, **6A** and **4,6A**). The yields

[†] Formulae are numbered according to the following system: **4A** is 4-trifluoromethyl-2-azidopyridine, **4T** is the corresponding tetrazolo[1,5-*a*]pyridine, and **4N** is the corresponding 2-pyridylnitrene; **5D** is 5-trifluoromethyl-1,2-didehydro-1,3-diazepine, *etc.*



1063



Scheme 2 Relative yields of products are given

were 30–60%.‡ The tetrazole–azide valence tautomerisms were investigated by ¹H and ¹³C NMR spectroscopy in CDCl₃, (CD₃)₂SO and CF₃CO₂D solution, and by IR spectroscopy in KBr, Me₂SO and CHCl₃. Compounds **6A** and **4,6A** exist as azides in the pure liquid state at room temperature. However, all these azide-to-tetrazole isomerisations are exothermic ($\Delta H^{\circ}_{isom} = -13$ to -30 kJ mol⁻¹) as determined by variable temperature ¹H NMR spectroscopy, thus implying that the existence of azides is an entropic effect; that this is indeed the case is revealed in the magnitude of ΔS°_{isom} (-50 to -59 ± 8 J mol⁻¹ K⁻¹). Simple sublimation, usually at 40–50 °C caused significant ring opening of tetrazoles to azides in most cases.

Photolysis (unfiltered; high-presure Xe–Hg lamp; 12–18 K; 2–20 min) of the Ar matrix isolated azide-tetrazole mixtures in each case caused the rapid disappearance of the azides and tetrazoles and formation of new products absorbing strongly in the carbodiimide region in the IR spectra and therefore assigned as the dehydrodiazepines **4D**, **5D**, **4,6D** and **5,6D** (Schemes 1–3).‡

It is particularly noteworthy that *the same* dehydrodiazepine (**5D**) is formed from the isomeric tetrazoles **5T** and **4T**



Scheme 3 Relative yields of products are given

(Scheme 1); likewise, one, and only one dehydrodiazepine **4D** is formed from **3T** and **6A** (Scheme 2), and these two dehydrodiazepines (**5D** and **4D**) are different from each other, as demonstrated by their FTIR spectra. Similarly, two different bis(trifluoromethyl)dehydrodiazepines were formed from the three different precursors shown in Scheme $3.\ddagger$

The ring opening and ring expansion reactions also occurred under flash vacuum pyrolysis (FVP) conditions as monitored by FTIR spectroscopy of the products isolated neat (*ca.* -196 °C) or in Ar matrix (*ca.* 20 K). Complete ring opening of tetrazoles to azides occurred at 200–250 °C/10⁻⁴ mbar. Formation of dehydrodiazepines occurred at 360–410 °C/10⁻⁴ mbar and the identities of **4D** and **5D** with those obtained photochemically were confirmed by Ar matrix isolation. In separate experiments it was shown by Ar matrix ESR spectroscopy of the products of FVP at 400 °C that nitrenes were formed and survived the pyrolysis conditions, giving rise to strong signals near 730 mT.⁵

At FVP temperatures above 450 °C the dehydrodiazepines disappeared (gradually) and nitriles were formed in their place. The products of the preparative pyrolyses at 520 °C/ 10^{-3} mbar were analysed by ¹H and ¹³C NMR, GC-mass, high resolution mass and IR spectroscopy.‡ The results are given in Schemes 1–3 (relative yields of 7–18 by ¹H NMR and/or GC).

[‡] Correct microanalytical data (C,H,N) were obtained for **3T**, **4T**, **5T**, **3,5T** and **4,5T** and correct high resolution mass spectra for **6A** and **7–18**. ¹H and ¹³C NMR spectra are available for all tetrazoles, azides, pyrroles and pentenedinitriles. FTIR spectra of dehydrodiazepines (Ar matrix, 12–18 K) **4D** v/cm⁻¹: 1992s, 1319m, 1294m, 1192s, 1164vs (v_{CF}), 1040m and 1006w; **5D** v/cm⁻¹: 1997s, 1311m, 1184m, 1143s (v_{CF}) and 985m; **4,6D** v/cm⁻¹: 2000s, 1279s, 1277s, 1200s, 1190w, 1186m, 1171s, 1151m (this and four preceding due to v_{CF}), 1041m and 985w; **5,6D** v/cm⁻¹: 2010m, 1336s, 1307s, 1183s, 1167s (v_{CF}), 991m, 970w and 875w.

References

- 2 O. L. Chapman, Pure Appl. Chem., 1979, 51, 331.
- 3 H. Sawanishi and T. Tsuchiya, J. Chem. Soc., Chem. Commun., 1990, 723.
- 4 J. B. Patrick, J. Chem. Educ., 1979, 56, 228; J. T. Welch and S. Eswarakrishnan, Fluorine in Bioorganic Chemistry, Wiley, New York, 1991; Selective Fluorination in Organic and Bioorganic Chemistry, ed. J. T. Welch, ACS Symposium Series No 456, Washington, 1991; J. T. Welch, Tetrahedron, 1987, 43, 3123; R. Bohlmann, Nachr. Chem. Tech. Lab., 1990, 38, 40; H. Nachtsheim, Nachr. Chem. Tech. Lab., 1992, 40, 56.
- 5 Nitrene **4,6N** was observed by both IR and ESR spectroscopy: P. Kambouris, T. Mosandl, R. A. Evans and C. Wentrup, unpublished results.
- 6 C. Wentrup, in *Reactive Intermediates*, ed. R. A. Abramovitch, Plenum Press, New York, 1980, vol. 1, p. 263.

The mechanism of ring contraction^{1,6} will be discussed elsewhere, but it is noteworthy that unique sets of products are formed from each set of starting materials, and there is little or no cross-over in products between reactions involving different nitrenes or dehydrodiazepines. This is due to the fact that almost no products of CF₃ group migration are formed; only the minor compound **18** from **4,6A** and **3,5T** and compounds **16** and **17** from **4,5T** (Scheme 3) necessitate such migration. To our knowledge, no other routes to cyano(trifluoromethyl)pyrroles are known. The present FVP reactions offer relatively easy access, particularly to the bis(trifluo-

romethyl)cyanopyrroles. The isolated yield of 18 from 4,5T was 72%. This research was supported by the Australian Research

Council. We thank Ishihara Sangyo Kaisha, Ltd., Tokyo, for generous gifts of chloro(trifluoromethyl)pyridines.

Received, 3rd March 1992; Com. 2/01145J

J. CHEM. SOC., CHEM. COMMUN., 1992

View Article Online

1064