

- (1972); D. R. Coulson and N. C. Yang, *ibid.*, **88**, 4511 (1966); A. Padwa and W. Bergmark, *Tetrahedron Lett.*, 5795 (1968); F. D. Lewis, *J. Am. Chem. Soc.*, **92**, 5602 (1970); F. D. Lewis, R. W. Johnson, and D. R. Kory, *ibid.*, **96**, 6101 (1974).
- (21) (a) G. Tsuchihashi, M. Yamauchi, and A. Ohno, *Bull. Chem. Soc. Jpn.*, **43**, 968 (1970); (b) J. C. Scalano and K. U. Ingold, *J. Am. Chem. Soc.*, **98**, 4727 (1976).
- (22) Bicyclohexyl appeared to be absent.¹⁰ Cyclohexene in small amounts could have escaped detection, but since disproportionation and combination are about equal²³ it seems unlikely that any significant amount was formed.
- (23) W. A. Cramer, *J. Phys. Chem.*, **71**, 1171 (1967); see M. J. Gibian and R. C. Corley, *Chem. Rev.*, **73**, 441 (1973).
- (24) J. R. Bolton, K. S. Chen, A. H. Lawrence, and P. de Mayo, *J. Am. Chem. Soc.*, **97**, 1832 (1975).
- (25) Sought, but unfortunately missed, previously.¹⁰
- (26) See, for instance, W. D. Ollis, M. Rey, I. O. Sutherland, and G. L. Closs, *J. Chem. Soc., Chem. Commun.*, 543 (1975); W. Braun, L. Rajkerback, and F. R. Einick, *J. Phys. Chem.*, **66**, 1591 (1962); H. Kiefer and T. A. Traylor, *J. Am. Chem. Soc.*, **89**, 6667 (1967).
- (27) We have not yet identified any termination products. The disulfide may be one (in small part) or other, more complex molecule which was nonvolatile and in small quantity. Others have observed this phenomenon with thiones: see ref 21b, and also J. C. Scalano, J. P. A. Tremblay, and K. U. Ingold, *Can. J. Chem.*, **54**, 3407 (1976).
- (28) Random walk in the absence of interactive forces. σ is the sum of the collision radii: S. W. Benson, "The Foundations of Chemical Kinetics", McGraw-Hill Co., New York, N.Y., 1960, p 495.
- (29) W. R. Ware and J. S. Novros, *J. Phys. Chem.*, **70**, 3246 (1966).
- (30) The structures of the products were established mass spectroscopically and by ¹H NMR and will be discussed in detail in our full publication.
- (31) B. Maillard, D. Forrest, and K. U. Ingold, *J. Am. Chem. Soc.*, **98**, 7024 (1976); see also J. K. Kochi, P. J. Krusic, and D. R. Eaton, *ibid.*, **91**, 1877, 1879 (1969).
- (32) G. A. Russell, *J. Am. Chem. Soc.*, **80**, 4987, 4997 (1958).
- (33) C. Walling and W. Thaler, *J. Am. Chem. Soc.*, **83**, 3877 (1961).

K. Y. Law, P. de Mayo,* S. K. Wong


Photochemistry Unit, Department of Chemistry
University of Western Ontario
London N6A 5B7, Ontario, Canada

Received April 5, 1977

Interconversion of [6]Paracyclophane and 1,4-Hexamethylene(Dewar benzene)¹


Sir:

Although many valence isomerizations of substituted benzenes are known,² photochemical isomerizations to a single Dewar form are rare, and there seems to be no example of a quantitative isomerization to a Dewar form followed by an equally clean thermal rearomatization. Hexafluorobenzene is photochemically isomerized in modest yield to a Dewar form which is reconverted to the aromatic form on heating.³ Similarly, photolysis of *peri*-di-*tert*-butylnaphthalenes produces a photostationary state in which the Dewar form greatly predominates. Here too, thermal rearomatization has been achieved.⁴ We present here our results on the light-induced closure of [6]paracyclophane (**1**) to 1,4-hexamethylene(Dewar benzene) (**2**) and the subsequent thermal rearomatization.



Irradiation of a solution of [6]paracyclophane⁵ in cyclohexane-*d*₁₂ with an unfiltered 450-W Hanovia medium-pressure mercury arc led to slow and apparently quantitative conversion to a single new product (33% in 90 min) which was identified as 1,4-hexamethylene(Dewar benzene) by a comparison of spectra with those of a sample prepared by the silver-catalyzed⁶ rearrangement of the bicyclopropenyl **3**.⁷ Treatment of **3** [NMR (CDCl₃): δ 7.10 (s, 4 H), 1.60 (m, 12

H)] with silver perchlorate in acetonitrile at 0 °C gives **2** and its 1,2 isomer **4** in the ratio 1:9. Separation by gas chromatography on a 3% S.E. 30 column at 55 °C gave **2** [NMR (CDCl₃): δ 6.53 (s, 4 H), 1.80 (m, 4 H), 1.50 (m, 8 H)] contaminated only by a few percent benzocyclooctene.⁷



Heating **2** between 50 and 90 °C results in a rapid and clean reversion to **1**. The activation parameters for the formation of **1** from **2** were determined both by integration of the signals for the aromatic protons in the NMR spectrum of **1** ($E_a = 20.9 \pm 1.5$ kcal/mol; $\log A = 9.8 \pm 0.9$) and by monitoring the band at 253 nm in the ultraviolet spectrum ($E_a = 19.9 \pm 0.9$ kcal/mol; $\log A = 9.3 \pm 0.6$). Thus only 1-chloro- and 1-fluoro-(Dewar benzene) exceed **1** in their measured rates of rearomatization.⁸

This work establishes the "breakpoint" in the series of [*m*]paracyclophanes and their Dewar isomers. In the hexamethylene case (**1**) and doubtless all higher homologues, it is the aromatic partner that is the more stable. For the pentamethylene case it has already been shown that the Dewar form is favored thermodynamically.⁶ Here the Dewar form does not rearrange to the open compound on heating in solution, but instead undergoes a remarkable rearrangement to benzocycloheptene. At higher temperatures [5]paracyclophane may be formed as an intermediate on the way to other compounds.⁶ It is not yet known if [5]paracyclophane is protected by a sufficiently high kinetic barrier to be isolable at lower temperatures.

Under conditions sufficient to completely rearrange **1** to **2**, the higher congener, [7]paracyclophane, undergoes only a slow photochemical polymerization. We are now testing to see if a Dewar benzene is formed, only to revert rapidly at room temperature to the aromatic compound.

References and Notes

- (1) The work at Princeton was generously supported by the National Science Foundation through Grant MPS-74-05690. Grant 1071 from the NATO Research Grants Programme greatly aided this collaborative effort and is warmly acknowledged.
- (2) For a review and references prior to 1972, see L. T. Scott and M. Jones, Jr., *Chem. Rev.*, **72**, 171 (1972), or G. Maier, "Valenzisomerisierungen", Verlag Chemie, Weinheim/Bergstr., 1972; see also (a) J. H. Dopfer, B. Grejdanus, and H. Wynberg, *J. Am. Chem. Soc.*, **97**, 216 (1975); (b) D. M. Lemal and L. H. Dunlap, Jr., *ibid.*, **94**, 6562 (1972); (c) W. Adam and J. C. Chang, *Int. J. Chem. Kinet.*, **1**, 487 (1969).
- (3) (a) I. Haller, *J. Phys. Chem.*, **72**, 2882 (1968); (b) G. Camaggi and F. Gozzo, *J. Chem. Soc. C*, 489 (1969); (c) E. Ratajczak and A. F. Trotman-Dickenson, *J. Chem. Soc. A*, 509 (1968); (d) G. P. Sameluk and R. D. S. Stevens, *Chem. Commun.*, 1720 (1970); for a study of pentafluorobenzene see E. Ratajczak, *Rocz. Chem.*, **44**, 447 (1970).
- (4) W. L. Mandella and R. W. Franck, *J. Am. Chem. Soc.*, **95**, 971 (1973).
- (5) V. V. Kane, A. D. Wolf, and M. Jones, Jr., *J. Am. Chem. Soc.*, **96**, 2643 (1974).
- (6) J. W. van Straten, I. J. Landheer, W. H. de Wolf, and F. Bickelhaupt, *Tetrahedron Lett.*, 4499 (1975), and references therein.
- (7) Compositions of new compounds were established by determinations of precise masses by mass spectral analyses.
- (8) R. Breslow, J. Napierki, and A. H. Schmidt, *J. Am. Chem. Soc.*, **94**, 5906 (1972); for other measured rates see ref 2b, 2c, 3a, 3c, 4, and H. C. Volger and H. Hogeveen, *Recl. Trav. Chim. Pays-Bas*, **86**, 830 (1967); J. F. M. Oth, *ibid.*, **87**, 1185 (1968).

Seetha L. Kammula, Linda D. Iroff, Maitland Jones, Jr.*

Department of Chemistry, Princeton University
Princeton, New Jersey 08540

J. W. van Straten, W. H. de Wolf, F. Bickelhaupt*

Scheikundig Laboratorium der Vrije Universiteit
Amsterdam-Z, The Netherlands

Received April 11, 1977