



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

SIMPLE AND RAPID SYNTHESIS OF BIOLOGICALLY ACTIVE PHTHALEMIDE AND N-HYDROXYPHTHALIMIDE DIMERS USING POLYMER-SUPPORTED ANIONS

D. S. Dalal^a, N. S. Pawar^b & P. P. Mahulikar^c

^a Moreshwar Arts & Science College, Bhokardan Dist., Jalna

^b Pratap College, Amalner Dist, Jalgaon

^c School of Chemical Sciences, North Maharashtra University, Jalgaon, 425 001, M.S., INDIA E-mail:

Published online: 18 Feb 2009.

To cite this article: D. S. Dalal, N. S. Pawar & P. P. Mahulikar (2007) SIMPLE AND RAPID SYNTHESIS OF BIOLOGICALLY ACTIVE PHTHALEMIDE AND N-HYDROXYPHTHALIMIDE DIMERS USING POLYMER-SUPPORTED ANIONS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 39:1, 81-85, DOI: [10.1080/00304940709458584](https://doi.org/10.1080/00304940709458584)

To link to this article: <http://dx.doi.org/10.1080/00304940709458584>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

SIMPLE AND RAPID SYNTHESIS OF BIOLOGICALLY ACTIVE PHTHALIMIDE AND *N*-HYDROXYPHTHALIMIDE DIMERS USING POLYMER-SUPPORTED ANIONS ^{†,††}

Submitted by
(07/14/06)

D. S. Dalal[†], N. S. Pawar^{††} and P. P. Mahulikar^{*}

[†] *Moreshwar Arts & Science College, Bhokardan Dist. Jalna*


^{††} *Pratap College, Amalner Dist. Jalgaon*

^{*} *School of Chemical Sciences, North Maharashtra University
Jalgaon - 425 001 (M.S.), INDIA*


E-mail: mahulikarpp@rediffmail.com

Compounds containing a phthalimide moiety are distinguished by their potent fungicidal action.^{1,2} Phthalimide and *N*-hydroxyphthalimide are important intermediates for the preparation of primary amines, *O*-substituted hydroxylamines, agricultural pesticides and are also used in preservatives, pigments and pharmaceuticals.¹⁻³ The phthaloyl group is a well-established protective group for primary amines⁴ in various types of compounds, particularly peptides,⁵ aminoglycosides⁶ and β -lactam antibiotics.⁷ Derivatives of both phthalimide and *N*-hydroxyphthalimide have been reported to possess good antibacterial efficacy and antifungal potency.^{8,9} However, a literature survey has revealed that the synthesis of phthalimide¹⁰⁻¹⁴ and *N*-hydroxyphthalimide¹⁵⁻¹⁷ dimers requires strictly anhydrous conditions with high temperature and long reaction times. Furthermore, purification of compounds is compulsory and tedious. In continuation of our work on polymer-supported reactions,^{8,9,18,19} we report herein a simple, rapid and environmentally friendly method for synthesis of phthalimide and *N*-hydroxyphthalimide dimers in better yields with higher purity under mild conditions.

Amberlite IRA-400 (chloride form) resin was used to support the phthalimide and *N*-hydroxyphthalimide anions. The α,ω -dibromoalkanes and diacid chlorides were added to the phthalimide and *N*-hydroxyphthalimide anion-supported resin in THF and the mixture was stirred until the reaction was complete, as monitored by silica gel TLC [Schemes 1 and 2]. The



present work appears to be an attractive alternative due to its experimental simplicity, high yields and facile isolation of pure products by simple filtration and evaporation of solvent.



EXPERIMENTAL SECTION

Phthalimide, *N*-hydroxyphthalimide, α,ω -dibromoalkanes and oxalyl chloride were synthetic grade chemicals (s. d. Fine Chem. Ltd., India). THF was freshly distilled prior to use. Malonyl chloride, succinoyl chloride and adipoyl chloride were prepared in the laboratory.²⁰ Commercial Amberlite IRA-400 (chloride form) resin was activated by treatment with 5 N HCl solution (100 mL) before use. The reactions of phthalimide and *N*-hydroxyphthalimide were monitored by silica gel TLC (pet. ether:CHCl₃, 1:1 and pet. ether:methanol, 9:1, respectively). The synthesized products were characterized by their physical constants, ¹H NMR spectroscopy and elemental analysis. Melting points and boiling points are uncorrected.

General Procedure for Supporting Phthalimide and *N*-Hydroxyphthalimide Anions on Amberlite IRA-400. Phthalimide (100 mmoles) was dissolved in 100 mL of 1N aqueous potassium hydroxide (100 mmoles). The activated Amberlite IRA-400 (chloride form) (60 g) was packed into a column (2 cm diameter and 45 cm length) and the above solution of the potassium salt of phthalimide was eluted slowly (1.5 mL/min). The resin was washed with distilled water, chloride ions and excess of phthalimide anion were removed. It was then washed with ethanol followed by acetone and dried *in vacuo* (10 mm Hg) at 50°C over P₂O₅ for 3 hr.

N-hydroxyphthalimide (100 mmoles) was similarly supported on the resin using 1N aqueous sodium hydroxide (100 mmoles). The exchange capacity of phthalimide and *N*-hydroxyphthalimide anion supported resin was determined by passing aqueous 1 N KCl and 1 N NaCl (100 mL), respectively, through the supported resin (1 g) packed in a column (0.5 cm diameter and 10 cm length). The anion in the eluent was titrated with 0.01 N HCl using methyl orange as an indicator. The exchange capacity of the supported resin was found to be 1.5 mmoles of phthalimide and 1 mmole of *N*-hydroxyphthalimide anion per gram of dry resin.

General Procedure for the Synthesis of α,ω -Diphthalimido (1-5) and α,ω -Diphthalimidoxy Alkanes (10-14). A mixture of Amberlite IRA-400 phthalimide or *N*-hydroxyphthal-

imide anion supported resin (15 mmoles), α,ω -dibromoalkane (7.5 mmoles) in THF (40 mL) was stirred and refluxed for 45 to 90 min depending on the reactivity of dibromoalkane. The progress of the reaction was monitored by TLC. After completion of the reaction, the resin was filtered and washed with THF (3 x 5 mL). The filtrate was dried over anhydrous sodium sulfate and removal of solvent gave products (**1-5** and **10-14**) in pure form as listed in *Table 1*.

Synthesis of *N*-Acyl (6-9**) and *N*-Acyloxy (**15-18**) Phthalimide Dimers.** The diacid chloride (7.5 mmoles) was added dropwise to a stirred suspension of Amberlite IRA-400 phthalimide or *N*-hydroxyphthalimide anion supported resin (15 mmoles) in THF (40 mL) at room temperature for 10 to 30 min depending on the reactivity of diacid chloride(s). The progress of the reaction was monitored by TLC. After completion of the reaction, the work-up as described above afforded the products (**6-9** and **15-18**) as listed in *Table 1*.

The compounds were characterized by comparison of their physical constants with literature data.^{21, 22} Newly synthesized compounds were characterized by ¹H NMR and IR spectroscopic methods and elemental analysis.

Table 1. Phthalimide and *N*-Hydroxyphthalimide Dimers

Cmpd No.	n	Yield (%)	mp (°C)	lit. ^{21, 22} (°C)	¹ H NMR ^{a)} (δ)
1	2	93	232	232-234	----
2	3	92	189	189	----
3	4	92	226	226-227	----
4	5	91	188	188	----
5	6	90	178	178-179	----
6	0	94	257 (dec.)	----	7.05-7.96 (m, 8H, Ar-H)
7	1	92	132	----	3.91 (s, 2H, CH ₂); 7.44-7.83 (m, 8H, Ar-H) ^{b)}
8	2	91	99-101	----	2.53 (s, 4H, CH ₂ CH ₂); 7.35-8.13 (m, 8H, Ar-H)
9	4	93	118-119	----	1.63 (m, 4H, middle CH ₂); 2.28 (t, 4H, 2 COCH ₂); 7.54-7.82 (m, 8H, Ar-H) ^{b)}
10	2	96	254	254	----
11	3	95	180	179-181	----
12	4	92	262	260-265	2.06 (m, 4H, middle CH ₂ CH ₂); 3.42 (t, 4H, 2 OCH ₂); 7.29-7.87 (m, 8H, Ar-H)
13	5	91	172	172-174	----
14	6	92	175	175-176	----
15	0	96	233 (dec.)	----	7.48-8.09 (m, 8H, Ar-H) ^{b)}
16	1	93	165	165-167	----
17	2	94	141	140-141	----
18	4	95	172-175 (dec.)	----	1.67 (m, 4H, middle CH ₂ CH ₂); 2.26 (t, 4H, 2 COCH ₂); 7.46-7.85 (m, 8H, Ar-H) ^{b)}

a) CDCl₃ b) CDCl₃ + DMSO-d₆

Table 2. Elemental Analysis Data for Compounds **1-18**

Cmpd No.	Elemental Analysis Data (Found)		
	C	H	N
1	67.50 (67.24)	3.78 (3.68)	8.75 (8.90)
2	68.26 (68.14)	4.22 (4.34)	8.38 (8.46)
3	68.96 (69.14)	4.63 (4.42)	8.04 (8.27)
4	69.60 (69.72)	5.01 (4.88)	7.73 (7.66)
5	70.20 (70.36)	5.36 (5.22)	7.44 (7.62)
6	62.07 (62.32)	2.32 (2.40)	8.04 (8.22)
7	62.99 (62.86)	2.78 (2.87)	7.73 (7.84)
8	63.83 (63.95)	3.21 (3.34)	7.44 (7.30)
9	65.35 (65.21)	3.99 (4.22)	6.93 (7.12)
10	61.36 (61.57)	3.43 (3.56)	7.96 (8.18)
11	62.29 (62.43)	3.85 (4.15)	7.65 (7.78)
12	63.16 (63.34)	4.22 (4.02)	7.36 (7.54)
13	63.95 (63.79)	4.60 (4.73)	7.10 (7.29)
14	64.69 (64.63)	4.93 (5.07)	6.86 (6.92)
15	56.85 (56.72)	2.12 (1.93)	7.37 (7.52)
16	57.88 (57.80)	2.56 (2.82)	7.10 (7.18)
17	58.83 (59.10)	2.96 (3.09)	6.86 (6.79)
18	60.55 (60.72)	3.69 (3.82)	6.42 (6.55)

REFERENCES

- [†] This work was presented at C. V. Raman Memorial Seminar, North Maharashtra University, Jalgaon, INDIA, Feb. 2004, PC 16, p. 16.
- ^{††} This paper is dedicated to Dr. P. P. Wadgaonkar, Division of Polymer Chemistry, National Chemical Laboratory, Pune- 411 008, (M.S.), INDIA.
1. G. S. Gruzdyes, V. A. Zinchenko and R. I. Slovtsov in *The Chemical Protection of Plants*, by G. S. Gruzdyes, Mir Publishers, Moscow, **1983**, 272.
 2. U. S. Sree Ramulu, in *Chemistry of Insecticides and Fungicides*, 2nd Ed., Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, **1995**, 246.
 3. *Ullmann's Encyclopedia of Industrial Chemistry*, Editors B. Elvers, S. Hawkins, G. Schulz, 5th Ed., **1991**, Vol. A 20, 190-191.
 4. T. W. Greene and P. G. M. Wuts *Protective Groups In Organic Synthesis*, 2nd Ed., Wiley: New York, **1991**.

5. M. Bondanszky and A. Bondanszky, *The Practice of Peptide Synthesis*, 2nd Ed., Springer: Heidelberg, **1994**.
6. P. J. Kocienski, *Protective Groups*, Thieme: Stuttgart, **1994**.
7. (a) T. Kamiya, M. Hashimoto, O. Nakaguchi and T. Oku, *Tetrahedron*, **35**, 323 (1979).
(b) C. A. Townsend, L. T. Nguyen, *Tetrahedron Lett.*, **25**, 4859 (1982).
8. N. S. Pawar, S. B. Attarde, D. S. Dalal, G. R. Sonar, U. R. Kapadi, D. G. Hundiware and P. P. Kumbhar (alias Mahulikar), *J. Indian Coun. Chem.*, **18(2)**, 29 (2001), Chem. Abst. 138:237963 (2002).
9. N. S. Pawar, U. R. Kapadi, D. G. Hundiware and P. P. Kumbhar (alias Mahulikar), *J. Sci Ind. Res*, **61**, 454 (2002), Chem. Abst. 137:196937 (2003).
10. D. Bogdal, J. Pielichowski and A. Boron, *Synlett*, 873 (**1996**).
11. D. Landini and F. Rolla, *Synthesis*, 389 (**1976**).
12. S. Wolfe and S. K. Hasan, *Can. J. Chem.*, **48**, 3566 (1970).
13. H. Normant and T. Cavigny, *Bull. Soc. Chim. Fr.*, 1866 (**1965**).
14. D. R. Maulding, *J. Org. Chem.*, **33**, 250 (1968).
15. L. Bauer and K. S. Suresh, *J. Org. Chem.*, **28**, 1604 (1963).
16. E. L. Schumann, *J. Med. Chem.*, **7**, 329 (1964).
17. L. J. Bollyky, *J. Org. Chem.*, **33**, 4266 (1968).
18. P. P. Kumbhar (Mahulikar), U. R. Kapadi, D. G. Hundiware, S. B. Attarde, P. M. Dewang, and N. S. Pawar, *Org. Prep. Proced. Int.*, **32**, 600 (2000).
19. D. S. Dalal, N. S. Pawar and P. P. Mahulikar, *Org. Prep. Proced. Int.*, **37**, 539 (2005)
20. B. S. Furniss, A. J. Hannaford, P. W. G. Smith, and A. R. Tatchell, *Vogel's Textbook of Practical Organic Chemistry*, 5th Ed., ELBS and Longman, London, **1994**.
21. *Beilstein Handbook of Organic Chemistry*, EIII-IV, Vol. 21.
22. M. J. Hearn, M. L. Cambell, M. Hoppe, J. Rosenberg and A. Sinha, *J. Chem. Eng. Data*, **31**, 255 (1986).
