FURTHER NEW REARRANGED LANOSTANOIDS FROM THE SEEDS OF <u>ABLES MARIESII</u> AND <u>A. FIRMA</u>

SHINICHI HASEGAWA^{*}, TADAYOSHI MIURA, NOBUTADA KANEKO, YOSHIYUKI HIROSE and YOICHI IITAKA[†]

Department of Applied Biological Science, Faculty of Science and Technology, Science University of Tokyo, Noda 278, Japan

[†] Faculty of Pharmaceutical Sciences, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113

(Received in Japan 22 December 1986)

<u>Abstract</u> - In addition to reported 17,14-<u>friedo</u>lanostane type (I) of triterpene further triterpenes having new rearranged lanostane skeletons, 17,13-<u>friedo</u>lanostane (II) and $8(14 \rightarrow 13R)$ <u>abeo</u>-17,13-<u>friedo</u>lanostane (III), have been isolated from the seed of <u>Abies</u><u>mariesii</u>. The skeletons I and II were also found to occur in <u>A</u>.<u>firma</u>. Their structures have been elucidated by spectral and X-ray diffraction analyses.

In the preceding papers, we reported the occurrence of an antimicrobial triterpene, mariesiic acid A $(\underline{5a})$,¹⁾ with an unusual skeleton, 17,14-<u>friedo</u>lanostane (I), in the seed of <u>Abies mariesii</u> Mast., and of several normal lanostane triterpenes, firmanoic acid (<u>1a</u>), isofirmanoic acid (<u>2a</u>) and firmanolides (<u>3</u> and <u>4</u>),²⁾ in the seed of <u>A. firma</u> Sieb. <u>et</u> Zucc. Further detailed search for both the seed extracts led to the isolation of five new rearranged lanostanoids, named mariesiic acid B (<u>6a</u>), mariesiic acid C (<u>7a</u>) and isomariesiic acid C (<u>8a</u>) from <u>A. mariesii</u> and 23-oxo-mariesiic acid A (<u>9a</u>) and 23-oxo-mariesiic acid B (<u>10a</u>) in common. In addition, <u>1a</u> and <u>Za</u> were also found to occur in <u>A. mariesii</u>. In the present paper, we describe the structural elucidation and biogenesis of these new triterpene acids along with their antimicrobial activities.

Ether extract (yield <u>ca</u> 30%) from the seed of <u>A. mariesii</u> comprised acidic components (22.8% of the extract), from which new triterpene acids were chromatographycally purified as their methyl esters (<u>6b-10b</u>) after treatment with diazomethane.

Methyl mariesiate B (<u>6b</u>) was analyzed by mass spectrum to have the molecular formula $C_{31}H_{48}O_4$ and its IR and ¹³C NMR (Table 2) spectra showed the presence of an α,β -unsaturated ester carbonyl (1705 cm⁻¹; 168.6 ppm) and two secondary hydroxyl groups [3600 and 3450 cm⁻¹; 66.9 (d) and 76.5 (d) ppm] in common with the case of <u>5b</u>. The ¹³C NMR spectrum showed also six <u>sp</u>² carbon signals due to three double bonds suggestive of tetracarbocyclic nature for this triterpene. The ¹H NMR spectrum (Table 1) revealed signals of two methine protons (δ 3.46 and 4.50) geminating to the hydroxyl groups, three trisubsutituted olefinic protons (δ 5.48, 5.63 and 6.70) and seven C-methyl groups ascribed to five tertiary, one secondary and one vinylic methyls. The above proton signals were very similar to those of <u>5b</u> except for the chemical shifts of two tertiary methyl groups and one olefinic proton [δ 5.48 for <u>6b</u>; δ 5.18 for <u>5b</u>], and thus <u>6b</u> involved 3a-hydroxyl group, C-7



double bond and the same side chain as that of 5b. The absence of absorption maximum above 202 nm in the UV spectra of 6b as well as the triol 11 derived from 6b by reduction with lithium aluminum hydride showed that two double bonds in the ring were unconjugated each other, in contrast to <u>5b</u> [<u>5b</u>: λ_{max} 219 nm; the triol <u>12</u>: λ_{max} 226 nm]. Therefore, two biogenetically conceivable structures, <u>6b</u> and <u>13</u>, were supposed for methyl mariesiate B. In the ¹H shift-correlated 2D-NMR spectrum, the olefinic proton signal at $_{\rm \delta}$ 5.48 (dd, J=8.4 and 2.4 Hz) was observed to correlate with signals at δ 2.22 (ddd, J=14.4, 8.4 and 3.4 Hz) and δ 1.83 (ddd, J=14.4, 12.0 and 2.4 Hz) assigned to allylic methylene protons. In addition, the allylic methylene signals were shown to correlate with a proton signal at δ 1.41 ascribed to 9β -H, shielded by the anisotropic effect of C-12 double bond, supporting the structure 6b for the compound. This was also in consistent with the fact that the largely different coupling constants between the olefinic proton and each of the allylic methylene protons were observed indicating the location of the double bond in the six-membered ring. Consequently, the strucutre of methyl mariesiate B was deduced as methyl $(24\underline{E})-3\alpha,23\underline{R}$ -dihydroxy-17,13-<u>friedo</u>-9 β -lanosta-7,12,24-trien-26-oate, 6b.

	<u>5b</u>	<u>9b</u>	<u>6b</u>	<u>10b</u>	<u>7b</u>	<u>8b</u>
3-н*	3.46 m	3.49 m	3.46 m	3.49 m		
7-H	5.57 br d	5.58 br d	5.63 br d	5.65 br d	1 5.44 dt	5.23 dt
	J=4.2	J=4.9	J=5.4	J=5.9	J=7.3, 2.2	J=7.6, 2.0
12-н			5.48 dd	5.49 dd		
			J=8.4, 2.4	J≠8.3, 2.2	2	
15-Н	5.18 dd	5.20 dd				5.27 m
	J=2.9, 1.2	J=2.9, 1.2				$W_{1/2} = 6.0$
21-н ⁺	0.94 d	0.85 d	0.98 d	0.85 d	0.84 d	0.69 d
23-н†	4.57 ddd		4.50 ddd			
24-н [†]	6.71 dq	7.09 g	6.70 dq	7.06 q	7.02 q	7.07 g
27-н [†]	1.87 d	2.22 d	1.87 d	2.22 d	2.22 d	2.21 d
angular	1.00, 0.97	1.00, 0.98	1.17, 0.98	1.21, 0.99	1.12, 1.10	1.19, 1.11
methyls	0.94, 0.93	0.95, 0.90	0.94, 0.93	0.93, 0.93	1.04, 0.81	1.08, 0.95
	0.86	0.90	0.93	0.93		
30-н				4	4.70 m, 4.56	m 1.43 m
				v	$W_{1/2} = 4 W_{1/2} = 0$	6 W _{1/2} =5
осн3	3.76	3.82	3.75	3.81	3.81	3.81
	<u>т</u>					

Table 1. ¹H NMR spectral data for $\underline{5b}-\underline{10b}$ (400 MHz, CDCl₃, J and W_{1/2}/Hz)

* W_{1/2}=7 Hz; [†]J_{20,21}=6.6 Hz, J_{22,23}=10 Hz, J_{22',23}=3 Hz, J_{23,24}=8 Hz, J_{24,27}=1.4 Hz.

Methyl mariesiate C (<u>7b</u>) had the molecular formula $C_{31}H_{44}O_4$ and its IR spectrum indicated strong absorption bands due to a methoxycarbonyl (1730 and 1255 cm⁻¹), two ketone carbonyl (1710 and 1700 cm⁻¹) and an exocyclic methylene (885 cm⁻¹) groups. In analogy with the case of methyl firmanoate (<u>1b</u>), the presence of a partial structure, $-COCH=C(CH_3)COOCH_3$, in the side chain was revealed by its mass fragment ion at $\underline{m/z}$ 127 ($C_6H_7O_3$), UV absorption maximum at 235 nm and proton signals [δ 7.02 (1H, q, J=1.5 Hz), 2.22 (3H, d, J=1.5 Hz) and 3.81 (3H, s)] and, on biogenetic consideration, another carbonyl group was supposed to be located at C-3. The ¹H NMR spectrum (Table 1) showed proton signals of another trisubstituted double bond (δ 5.44), an exocyclic methylene group (δ 4.56 and 4.70) and four tertiary and one secondary methyl groups. The presence of the secondary methyl

Table 2. ¹³C NMR spectral data (25.1 MHz, CDCl₃)

с	<u>5b</u>	<u>9b</u>	<u>21</u> *	<u>6b</u>	<u>10b</u>	<u>22</u> *	<u>7b</u>	<u>8b</u>
1	29.2	28.7	35.6	29.7	29.4	36.2	36.3	36.6
2	25.3	25.3	35.0	25.2	25.4	35.1	34.9	34.9
3	76.4	76.5	216.2	76.5	76.5	216.4	216.3	216.3
4	37.1	37.1	47.3	37.0	37.0	47.2	47.5	47.4
5	37.9	38.0	44.9	37.9+	38.0	45.2	48.0	46.9
6	23.1	23.1	23.9	23.2	23.2	23.9	24.4	24.0
7	120.4	120.8	120.2	122.0	122.5	121.8	123.8	122.6
8	136.8	136.6	136.7	146.2	146.0	146.2	144.8	145.5
9	53.1	52.9	52.4	51.1	51.0	50.4	58.2	55.6
10	34.7	34.8	34.7	34.8	34.8	34.8	34.9	34.6
11	33.5	33.5	33.3	28.2	28.1	27.7	24.6+	26.1+
12	25.3	25.3	25.0	118.6	118.7	118.4	27.7 ⁺	27.0+
13	51.7	51.7	51.7	156.4	156.0	156.2	61.4	63.1
14	153.3	152.8	152.0	50.0	50.0	49.8	160.8	146.5
15	114.7	115.0	115.7	36.7	36.8	36.9	33.0 ^{††}	118.7
16	45.4	45.0	44.9	38.1+	38.4	38.1	33.5**	38.4
17	50.8	50.5	50.5	46.8	46.4	46.5	49.1	51.2
18	16.7	16.6	16.6	25.2	24.8	24.9	16.8	22.3
19	22.3	22.3	22.6+	22.2	22.2	22.6+	22.4	22.3
20	33.5	34.0	33.9	38.1	38.7	38.6	31.1	35.3
21	19.0	19.2	19.2	15.4	15.8	15.8	17.2	19.7
22	39.5	48.6	48.6	39.3	48.6	48.4	49.0	49.0
23	66.6	202.0	201.7	66.9	202.4	202.1	201.9	202.1
24	145.3	133.0	132.9	144.9	133.0	132.9	133.1	132.9
25	126.4	140.0	140.1	126.8	139.8	139.8	139.9	140.1
26	168.6	168.1	168.0	168.6	168.2	168.0	168.1	168.1
27	12.6	14.3	14.3	12.7	14.3	14.2	14.4	14.3
28	28.3	28.2	25.5	28.2	28.1	25.5	26.0	25.7
29	23.1	23.1	21.6+	23.0	23.0	21.5+	22.8	22.7
30	15.4	17.1	17.1	26.1	25.8	25.9	104.5	13.2
OMe	51.8	52.5	52.4	51.9	52.5	52.4	52.6	52.5

* For assistance of carbon assignment methyl 3,23-dioxo-mariesiates A $(\underline{21})$ and B ($\underline{22}$) were prepared. +, ++ Assignments may be interchanged.

group ascribable to 21-CH₃ and the lack of an angular methyl group for the tetracyclic triterpene suggested that the exocyclic methylene group must come from the angular methyl group. Thus, methyl mariesiate C was considered to have an unusual carbon skeleton which might arise from the lanostane and the establishment of its gross structure was subjected to X-ray analysis.

The crystalline derivative 14 suited for the analysis was prepared together with the epimer 15 at C-23 from 7b by reduction with sodium borohydride. The crystals grown in a solution of chloroform-hexane were thin plates, mp 105-106 °C, solvated with water. The cell dimensions and intensity data were obtained from the measurement on a Philips PW1100 diffractometer using Cu Klpha radiation monochromated with graphite plate.

Crystal data: $C_{31}H_{48}O_4 \cdot H_2O$, FW=502.7, monoclinic, space group P2₁, a=12.208(8), b=17.445(10), c=6.967(4) Å, ß=105.40(5)°, V=1430.5 Å³, Z=2, D_c=1.167 g cm⁻³, μ for Cu K α =5.7 cm⁻¹.

Intensities of 2594 reflections out of 2953 theoretically possible ones were measured as above the $2\sigma(\underline{I})$ level in the 20 range 6° through 156°. The crystal structure was determined by the direct method based on \underline{MULTAN}^{3} procedure and refined by the block-diagonal least-squares method. All the hydrogen atoms except those attached to C-31 were found on the difference electron-density map and their atomic coordinates and isotropic temperature factors were refined. The final <u>R</u> value was 0.052.⁴⁾ The molecular structure was illustrated in Fig. 1 drawn by <u>PLUTO</u>⁵⁾ program. Accordingly, the structure of methyl mariesiate C itself was determined as methyl (24<u>E</u>)-3,23-dioxo-8(14-13<u>R</u>)abeo-17,13-friedo-9\beta-lanosta-7,14 (30),24-trien-26-oate represented by the chemical structure <u>7b</u>.



Figure 1. Molecular structure of the diol $\underline{14}$ derived from methyl mariesiate C ($\underline{7b}$).

Methyl isomariesiate C (8b) having the molecular formula $C_{31}H_{44}O_4$ showed a UV absorption maximum at 235 nm and IR absorption bands at 1720, 1700 and 1695 cm⁻¹ in common with the case of <u>7b</u>, suggesting the presense of 3-oxo group and the same side chain as that of 7b, which was ascertained by their proton and carbon signals in its NMR spectra (see Tables 1 and 2). In addition, the ¹H NMR spectrum showed proton signals for two trisubsituted double bonds (δ 5.27 and 5.23) and four tertiary and one vinylic (δ 1.43) methyl groups, the last of which was shown to be coupled with the olefin signal at δ 5.27 by decoupling experiment. The olefin signal at δ 5.23 appearing as double triplet (J=7.6 and 2.0 Hz) was assigned to that of C-7 double bond because the coupling pattern was almost identical to that of 7b. The above spectral features suggested that methyl isomariesiate C was an isomer (8b) of 7b in which the double bond at C-14 migrated into the ring. This was confirmed by the formations of the compound by acid catalyzed isomerization of 7b and of a mixture of four diastereoisomers at C-14 and C-25, 18, from 15 and 17, derived from <u>8b</u>, upon catalytic hydrogenation.

The structures of the other two triterpenes designated methyl 23-oxomariesiates A $[C_{31}H_{46}O_4, \lambda_{max} 232 \text{ nm} (\epsilon 12800), \nu_{max} 3640, 1730, 1690 \text{ cm}^{-1}]$ and B $[C_{31}H_{46}O_4, \lambda_{max} 235 \text{ nm} (\epsilon 9500), \nu_{max} 3650, 1730, 1690 \text{ cm}^{-1}]$, were supposed to be 23-oxo derivatives corresponding to $\underline{5b}$ and $\underline{6b}$ mainly on the basis of their ¹H and ¹³C NMR spectral comparisons (see Tables 1 and 2). This was chemically confirmed by conversions of $\underline{5b}$ and $\underline{6b}$ with manganese dioxide into $\underline{9b}$ and $\underline{10b}$, respectively, identical in all respects with the natural compounds.

The new skeletons I-III of the above triterpenes were considered most likely to be biosynthesized from the lanostane skeleton by enzymic dehydrogenation of 17-H or dehydroxylation of 17-OH followed by successive 1,2-shifts of methyl group(s) and a ring bond as shown in Scheme 1, because the normal lanostane and/or its 17oxygenated triterpens ($\underline{1-4}$) co-occurred in both seeds.



Scheme 1. Postulated biogenesis of the rearranged lanostanoids.

As summarized in Table 3, all triterpene acids exhibited antimicrobial activity against Gram-positive bacteria and actinomycetes tested. However, their methyl esters and simple model compounds with the side chain moiety, $(2\underline{E})$ -2-methyl-4-oxo-2-pentenoic acid (<u>19</u>) and its 4-hydroxyl derivative (<u>20</u>), were inactive at 30 µg/disc. This result suggested that not only the carboxylic moiety but also the hydrophobic group played an important role in revealing the inhibitory activity.

		<u>5a</u>		-	<u>6a</u>		<u>7a</u>	<u>9a</u>	<u>10a</u>	<u>1a</u>
	Concentration					(µg/	disc)			
Test organism	30	15	7.5	30	15	7.5	 30	30	30	30
<u>Bacillus</u> <u>subtilis</u>	14	12	9	14	12	+	9	9	11	13
Micrococcus luteus	15	13	9	16	12	10	9	9	12	13
<u>Nocardia</u> <u>corallina</u>	11	10	9	9	+	-	+	+	+	+

Table 3. Antimicrobial activity of mariesiic acids and firmanoic acid*

* The inhibitory activity was represented by the diameter of clear zone using paper disc (\$\phi\$ 8 mm\$) after incubation at 30 °C for 21 h. All compounds tested were inactive against Gram-negative bacteria (<u>Escherichia coli</u>, <u>Pseudomonas aeruginosa</u>) and yeast (<u>Saccharomyces cerevisiae</u>, <u>Candida utilis</u>).

EXPERIMENTAL

<u>General</u>. Melting points are uncorrected. Optical rotations were determined on a JASCO DIP-SL automatic polarimeter. UV spectra were measured in EtOH solution with a Hitachi 320 spectrophotometer. IR spectra were obtained on a Hitachi 215 spectrophotometer. ¹H NMR spectra were recorded on JEOL JNM-FX 100 (100 MHz) and JEOL JNM GX-400 (400 MHz), and ¹³C NMR spectra were obtained on the former instrument (25.1 MHz) each in CDCl₃ containing TMS as an internal standard. Mass spectra were operated on a Hitachi M-80 using direct insertion at 70 eV. Wakogel C-200 and LiChroprep Si 60 (Merck) were used for column chromatography. Silica gel 60 HF_{254} (Merck) and precoated silica gel 60 F_{254} were used for preparative and analytical TLC, respectively.

Extraction. Seeds of <u>A.</u> mariesii (900 g), collected in Nagano Prefecture, Japan, were ground and extracted with ether (2 1 x 3). The combined ether extract was concentrated to <u>ca</u> 1 1 and treated successively with aqueous 5% Na_2CO_3 and 5% NaOH (each 500 ml x 3). The ether layer was dried over Na_2SO_4 and evaporated to afford a neutral portion (210 g). Both the aqueous alkaline layers were acidified (pH 3) with 6 N HCl under cooling and extracted with ether to yield a strongly acidic (53 g) and a weakly acidic (9 g) portions. Extraction and fractionation from seeds of <u>A. firma</u> was described in the preceding paper.²

Isolation of mariesiic acids A (5a) and B (6a) from A. mariesii. The strongly acidic portion (53 g) was chromatographed on a silica gel column (36 cm x 7.5 cm i.d.) eluting with hexane-EtOAc (2 : 1, 3 l) and then hexane-EtOAc (1 : 2, 3 l), each 300 ml of the eluant being collected. The fractions 10-20 deposited almost pure crystals (ca 5 g) which were recrystallized from EtOAc to afford 5a.¹⁾ The combined mother liquor of the fractions 10-20 gave an oil (12.5 g), a portion (1.2 g) of which was methylated with an ethereal solution of diazomethane and chromatographed on a 10% AgNO₃-silica gel (130 g) column (ether-CHCl₃, 1 : 2) to give 5b (361 mg) and 6b (182 mg).

<u>5b</u>: a gum; UV: λ_{max} 219 nm (ϵ 16300); IR (CHCl₃): ν_{max} 3610, 3470, 2940, 2880, 1705, 1650, 1440, 1370, 1240, 1130, 1100, 1040, 980, 900, 835 cm⁻¹; H and ¹³C NMR: Tables 1 and 2; MS m/z (rel. int.) 484 (M⁺, 35), 313 (100), 187 (31), 170 (38), 159 (31), 135 (42), 131 (32), 109 (32), 107 (31), 43 (63), 41. (33%).

A solution of <u>6b</u> (103 mg) in 1 M NaOH/MeOH-H₂O (9 : 1) (10 ml) was stirred at room temperature for 2 h. After acidification with 5% HCl (pH 3) the mixture was extracted with ether. The ether extract was washed with brine, dried over Na_2SO_4 and evaporated to afford <u>6a</u> (69 mg) and <u>6b</u> (25 mg) after purification on a silica gel column (hexane-EtOAc, 2 : 3).

On a similar treatment as above, <u>6b</u> gave the triol <u>11</u>, mp 184-188 °C (fine needles from EtOAc); UV: λ_{max} 199 nm (end absorption); IR (KBr): ν_{max} 3275, 2970, 1440, 1375, 1360, 1055, 975, 950, 900, 825 cm⁻¹; ¹H NMR (100 MHz): δ 5.64-5.38 (3H, m, 7-, 12- and 24-H), 4.42 (1H, br t, J=9

Hz, 23-H), 4.00 (2H, s, 25-H), 3.44 (1H, m, $W_{1/2}=6.5$ Hz, 3-H), 1.71 (3H, d, J=1.4 Hz, 26-H), 1.16 (3H, s, CH₃), 0.99-0.94 (CH₃ x 5); HRMS: m/z 456.3617 (M⁺, calcd for C₃₀H₄₈O₃: 456.3605); MS: m/z (rel. int.) 456 (M⁺, 0.4), 438 (20), 423 (24), 405 (15), 313 (41), 295 (71), 173 (49), 145 (48), 141 (100), 123 (97), 121 (87), 107 (48), 105 (50), 55 (94), 43 (64), 41 (51%).

Isolation of methyl firmanoate (1b), methyl isofirmanoate (2b), methyl mariesiate C (7b) and methyl isomariesiate C (8b). The combined fractions 5-9 (4.4 g) was dissolved in ether (88 ml) and methylated with an ethereal solution of diazomethane (\underline{ca} 20 ml) at - 15 °C. The methyl ester was subjected to repeated column chromatography on silica gel using hexane-EtOAc (7 : 1) and benzene-ether (30 : 1) to give $\underline{7b}$ (1.1 g), $\underline{2b}$ (101 mg) and a mixture of $\underline{1b}$ and $\underline{8b}$ (595 mg). The mixture was separated by preparative TLC on 10% AgNO₃-silica gel (ether-CHCl₃, 1 : 40) to afford $\underline{1b}$ (Rf 0.5-0.6, 485 mg) and $\underline{8b}$ (Rf 0.6-0.7, 80 mg). The fractions 5-9 also contained a small amount (10 mg) of $\underline{9b}$ and $\underline{10b}$ in a ratio of \underline{ca} 1 : 2, which was confirmed by comparison of the ¹H NMR spectrum and TLC behavior with those isolated from <u>A. firma</u> described below.

<u>8b</u>: a gum, $[\alpha]_{D} = 154^{\circ}$ (c 3.18, CHCl₃); UV: $\lambda_{max} 235 \text{ nm}$ ($\epsilon 12700$); IR (CHCl₃): $\nu_{max} 2950$, 2860, 1720, 1700, 1695, 1620, 1440, 1380, 1265, 1220, 1115, 885 cm⁻¹; ¹H and ¹³C NMR: Tables 1 and 2; HRMS: m/z 480.3214 (M⁺, calcd for C₃₁H₄₄O₄: 480.3241); MS: m/z (rel. int.) 480 (M⁺, 20), 338 (C₂₄H₃₄O, 100), 311 (C₂₁H₃₁O, 49), 309 (C₂₂H₂₉O, 30), 187 (C₁₄H₁₉, 61), 185 (C₁₄H₁₇, 61), 145 (C₁₁H₁₃, 34), 131 (C₁₀H₁₁, 51), 127 (C₆H₇O₃, 57), 105 (C₈H₉, 34), 91 (29), 69 (36), 55 (34), 41 (35%).

 $\frac{\text{NaBH}_{4}}{\text{mixture was stirred for 30 min.}} \text{ To a solution of } \frac{7\text{b}}{11 \text{ mg}} \text{ in MeOH (20 ml) was added NaBH}_{4} and the mixture was stirred for 30 min. After addition of aqueous NH_{4}Cl, the mixture was extracted with ether. The extract was washed with brine, dried over MgSO_{4} and evaporated. The product was chromatographed on a silica gel column (hexane-ether, 1 : 3) to yield the diols <math>\frac{15}{15}$ (271 mg) and $\frac{14}{14}$ (90 mg).

14: mp 105-106 °C (thin plates from hexane-CHCl₃, monohydrated form); IR (KBr): v_{max} 3530, 3340, 2950, 2860, 1695, 1645, 1440, 1380, 1270, 1065, 880, 840, 755 cm⁻¹; ¹H NMR (100 MHz): 6 6.65 (1H, dq, J=8.8, 1.4 Hz, 24-H), 5.36 (1H, m, 7-H), 4.67, 4.56 (each 1H, narrow m, 30-H), 4.48 (1H, m, 23-H), 3.76 (3H, s, OCH₃), 3.22 (1H, dd, J=7.7, 6.3 Hz, 3-H), 1.89 (3H, d, J=1.4 Hz, 27-H), 1.07-0.81 (CH₃ x 5); ¹³C NMR: 36.0, 27.5, 79.4, 38.5, 46.8, 23.5, 124.3, 144.8, 58.5, 35.3, 24.7, 27.6, 61.5, 162.1, 32.8, 33.7, 49.8, 16.6, 23.5, 33.0, 16.5, 40.2, 67.7, 143.5, 128.5, 168.2, 13.3, 28.9, 16.2, 104.7 (assignments: from C-1 to C-30), 52.1 (OCH₃); HRMS: m/z 484.3552 (M⁺, calcd for C₃₁H₄₈O₄: 484.3554); MS: m/z (rel. int.) 484 (M⁺, 27), 466 (18), 313 (68), 197 (79), 165 (67), 135 (72), 131 (67), 119 (75), 97 (100), 91 (67), 81 (71), 69 (67), 41 (96%).

<u>15</u>: mp 152-153 °C (fine needles from hexane-ether); IR (KBr): v_{max} 3400, 2950, 2860, 1710, 1645, 1440, 1380, 1300, 1250, 1140, 1085, 1035, 1005, 875, 835, 755 cm⁻¹; ¹H NMR (100 MHz): δ 6.58 (1H, dq, J=8.0, 1.4 Hz, 24-H), 5.36 (1H, m, 7-H), 4.68, 4.59 (each 1H, narrow m, 30-H), 4.48 (1H, m, 23-H), 3.72 (OCH₃), 3.21 (1H, dd, J=7.7, 6.3 Hz, 3-H), 1.83 (3H, d, J=1.4 Hz, 27-H), 1.04-0.79 (CH₃ x 5); ¹³C NMR: 36.0, 27.5, 79.4, 38.5, 46.7, 23.6, 123.8, 144.7, 58.1, 35.3, 24.7, 27.8, 61.5, 162.1, 32.9, 33.8, 49.6, 16.6, 23.4, 31.7, 15.4, 38.7, 66.5, 145.0, 126.7, 168.5, 12.7, 28.9, 16.1, 104.4 (assignments: from C-1 to C-30), 52.0 (OCH₃); HRMS: m/z 484.3554 (M⁺, calcd for C₃₁H₄₈O₄: 484.3554); MS: m/z (rel. int.) 484 (M⁺, 26), 466 (33), 313 (72), 135 (83), 131 (73), 119 (81), 97 (100), 91 (71), 81 (75), 69 (71), 41 (98%).

On a similar treatment as above <u>8b</u> gave the diols <u>16</u> and <u>17</u> in a ratio of <u>ca</u> 1 : 3 after chromatography on a LiChroprep RP-8 (Rf 0.43 and 0.37, H_2O -MeOH, 15 : 85).

<u>16</u>: mp 115-116 °C (needles from EtOAc-hexane); IR (KBr): v_{max} 3350, 2965, 2940, 2860, 1715, 1650, 1450, 1375, 1260, 1040, 750 cm⁻¹; ¹H NMR (100 MHz): δ 6.56 (1H, dq, J=7.6, 1.5 Hz, 24-H), 5.21 (1H, narrow m, 15-H), 5.16 (1H, m, 7-H), 4.47 (1H, m, 23-H), 3.75 (OCH₃), 3.19 (1H, dd, J=6.5, 8.0 Hz, 3-H), 1.89 (3H, d, J=1.5 Hz, 27-H), 1.43 (3H, narrow m, 30-H), 0.99 (6H, s), 0.94 (3H, s), 0.87 (3H, s), 0.75 (3H, d, J=6.5 Hz, 21-H); HRMS: m/z 484.3536 (M⁺, calcd for C₃₁H₄₈O₄: 484.3554); MS: m/z (rel. int.) 484 (M⁺, 13), 313 (26), 197 (100), 165 (51), 131 (49), 105 (37), 97 (50), 69 (42), 55 (37), 43 (47), 41 (61%).

 17:
 mp 105-107 °C (needles from ether-hexane); IR (KBr): v_{max} 3400, 2950, 2870, 1715, 1650, 1450, 1375, 1240, 1050, 750 cm⁻¹; ¹H NMR (100 MHz): δ 6.66 (1H, dq, J=8.0, 1.5 Hz, 24-H), 5.22 (1H, narrow m, 15-H), 5.16 (1H, m, 7-H), 4.47 (1H, m, 23-H), 3.73 (OCH₃), 3.20 (1H, dd, J=6.5, 8.0 Hz, 3-H), 1.86 (3H, d, J=1.5 Hz, 27-H), 1.44 (3H, narrow m, 30-H), 0.99 (9H, s), 0.88 (3H, d)

s), 0.76 (3H, d, J=6.9 Hz, 21-H); HRMS: m/z 484.3556 (M^+ , calcd for $C_{31}H_{48}O_4$: 484.3554); MS: m/z (rel. int.) 484 (M^+ , 10), 313 (28), 197 (100), 165 (55), 131 (58), 105 (40), 97 (58), 69 (45), 55 (38), 43 (41), 41 (62%).

<u>Acid catalyzed isomerization of 7b into 8b</u>. To a CDCl_3 solution of <u>7b</u> in a NMR sample tube was added a drop of 10% HCl-MeOH. By monitoring on the ¹H NMR spectrum and TLC, <u>7b</u> was pursued to be isomerized completely into <u>8b</u> at room temperature in 22 h.

Catalytic hydrogenation of 15 and 17. The diol 15 (50 mg) was hydrogenated over PtO, (15 mg) in EtOH (3 ml) at room temperature for 17 h, while the diol 17 for five days. The major products from 15 and 17, showing the same Rf value on TLC, collected by chromatography on 10% AgNO₁-silica gel (EtOAc-hexane, 1 : 2) yielded a mixture of the tetrahydro derivatives <u>18</u> as a solid, which showed the following spectra: ¹H NMR (100 MHz): § 5.30 (m, 7-H), 3.61 (OCH₃), 3.60 (m, 23-H), 3.20 (dd, J=6.5, 8.0 Hz, 3-H), 2.64 (sextet, J=7.1 Hz, 25-H), 1.20-0.69 (methyl groups); ¹³C NMR: § 177.8 and 177.4 (C-26), 145.1 (C-8), 120.4 and 120.1 (C-7), 79.4 (C-3), 68.9 and 68.1 (C-23), 60.2 and 57.4 (C-13), 58.8 and 57.0 (C-9), 51.7 (OCH₃), 50.1 and 49.7 (C-17), 46.1 (C-5), 45.5 (C-14), 42.9, 42.7, 42.5 and 42.4 (C-22 and C-24), 40.3 (t), 38.4 (C-4), 37.3 and 37.0 (C-25), 35.9 and 35.8 (C-1), 35.1 (C-10), 34.6 (t), 33.9 (C-20), 33.2 (t), 32.0 (t), 30.7 (t), 28.8 (C-28), 27.5 (C-2), 27.0 (t), 26.7 (t), 26.3 (t), 23.4 (C-6 and C-19), 20.2 (g), 17.6 (g), 17.5 (g), 17.3 (g), 16.9 (g), 16.0 (C-29), 15.2 and 15.0 (C-21); MS: m/z 488 (M⁺), 456, 261, 260, 243, 222, 215, 135, 133, 131, 121, 119, 105, 95, 69, 55, 43, 41. Considering from the appearance of the carbon signals as double lines, assigned to C-7, C-13, C-17 and carbon atoms in the side chain in the 13 C NMR spectra, the tetrahydro derivatives were deduced to be composed of four diastereoisomers concerning C-14 and C-25.

 $\frac{\text{Isolation of 9b and 10b from A. firma. The chromatographycally most polar fraction (3.9 g) }{\text{of the acidic portion reported}^2} \text{ was methylated with diazomethane at - 15 °C. A portion of the methyl ester (770 mg) was subjected to preparative TLC on 10% <math>\text{AgNO}_3$ -silica gel developing with CHCl_3 -ether (80 : 1) two times to give 9b (Rf 0.37-0.27, 200 mg) and 10b (Rf 0.23-0.10, 453 mg). 9b: a gum, $[\alpha]_D$ + 53.2 °(c 1.28, CHCl_3); UV: λ_{max} 232 nm (ϵ 12800); IR (CCl_4): ν_{max} 3640, 2970, 2880, 1730, 1690, 1620, 1440, 1385, 1365, 1255, 1125, 1100, 1065, 985, 905 cm⁻¹; H and ¹³C NMR: Tables 1 and 2; HRMS: m/z 482.3400 (M⁺, calcd for $C_{31}H_{46}O_4$: 482.3398); MS: m/z (rel. int.) 482 (M⁺, 11), 340 ($C_{24}H_{36}O$, 78), 314 (34), 313 ($C_{22}H_{33}O$, 100), 295 ($C_{22}H_{31}$, 50), 187 ($C_{14}H_{19}$, 32), 169 ($C_{9}H_{13}O_3$, 66), 127 ($C_{6}H_7O_3$, 36%).

 $\frac{10b}{2880}; a gum, [a]_{D} - 157.4 \circ (c 1.98, CHCl_{3}); UV: \lambda_{max} 235 nm (\epsilon 9500); IR (CCl4): v_{max} 3650, 2970, 2880, 1730, 1690, 1620, 1440, 1385, 1370, 1255, 1125, 1060, 985, 905 cm⁻¹; H and ¹³C NMR: Tables 1 and 2; HRMS: m/z 482.3374 (M⁺, calcd for <math>C_{31}H_{46}O_4$: 482.3398); MS: m/z (rel. int.) 482 (M⁺, 28), 467 ($C_{30}H_{43}O_4$, 36), 464 ($C_{31}H_{44}O_3$, 27), 449 ($C_{30}H_{41}O_3$, 41), 353 ($C_{23}H_{29}O_3$, 48), 313 ($C_{22}H_{33}O$, 98), 295 ($C_{22}H_{31}$, 100), 173 ($C_{13}H_{17}$, 62), 127 ($C_{6}H_7O_3$, 86), 123 ($C_{9}H_{15}$, 73), 121 ($C_{9}H_{13}$, 60%).

 $\frac{\text{MnO}_2}{\text{was stirred at room temperature for 30 min.}} \text{ After filtration followed by evaporation, the product was purified on a silica gel column (ether-CHCl₃, 1 : 20) to afford <u>9b</u> (70 mg) identical in all respects (IR, ¹H NMR, MS and TLC) to the natural compound. On a similar treatment as above <u>6b</u> gave <u>10b</u>.$

<u>Jones oxidation of 9b and 10b.</u> To a stirred solution of <u>9b</u> (100 mg) in acetone (10 ml) was added dropwise Jones' reagent under cooling. After usual work-up the product was purified on a silica gel column (ether-CHCl₃, 1 : 50) to yield <u>21</u> (80 mg), a gum; UV: λ_{max} 231.5 nm (c 14700); IR (CCl₄): v_{max} 2970, 1730, 1710, 1700, 1620, 1450, 1435, 1385, 1365, 1255, 1125 cm⁻¹; ¹H NMR (400 MHz): δ 7.08 (1H, g, J=1.5 Hz, 24-H), 5.62 (1H, m, 7-H), 5.26 (1H, dd, J=1.5, 3.3 Hz, 15-H), 2.22 (3H, d, J=1.5 Hz, 27-H), 0.85 (3H, d, J=6.6 Hz, 21-H), 3.81, 1.14, 1.11, 1.10, 0.90, 0.89 (each 3H, s); ¹³C NMR: Table 2; HRMS: m/z 480.3238 (M⁺, calcd for C₃₁H₄₄O₄: 480.3241); MS: m/z (rel. int.) 480 (M⁺, 9), 339 (15), 338 (C₂₄H₃₄O, 58), 323 (C₂₃H₃₁O, 14), 312 (19), 311 (C₂₂H₃₁O, 54), 187 (C₁₄H₁₉, 13), 170 (18), 169 (C₉H₁₃O₃, 100), 127 (C₆H₇O₃, 20%).

On a similar treatment as above <u>10b</u> gave <u>22</u>, a gum; UV: λ_{max} 234.5 nm (ϵ 9900); IR (CCl₄): ^{2980, 1730, 1715, 1700, 1620, 1455, 1440, 1385, 1370, 1260, 1130, 1120, 1070, 990, 905 cm⁻¹; ^{1max} (400 MHz): δ 7.06 (1H, g, J=1.5 Hz, 24-H), 5.68 (1H, m, 7-H), 5.48 (1H, dd, J=8.4, 2.5 Hz, 12-H), 2.20 (3H, d, J=1.5 Hz, 27-H), 0.90 (3H, d, J = 6.6 Hz, 21-H), 3.80, 1.20, 1.11, 1.10, 1.09, 0.95 (each 3H, s); ¹³C NMR: Table 2; HRMS: m/z 480.3227 (M⁺, calcd for C₃₁H₄₄O₄: 480.3241); MS: m/z (rel. int.) 480 (M⁺, 12), 447 (C₃₀H₃₉O₃, 25), 312 (25), 311 (C₂₂H₃₁O, 100), 201 (C₁₄H₁₇O, 21), 169 (23), 127 (C₆H₇O₃, 43), 123 (35), 121 (C₉H₁₃, 32), 107 (C₈H₁₁, 21%).}

<u>Preparation</u> of (2E)-2-methyl-4-oxo-2-pentenoic acid (19). A mixture of acetone (2.5 ml), pyruvic acid (1 ml) and 85% phosphoric acid (2.5 ml) was refluxed for 70 h, as reported⁶⁾. The

reaction mixture was poured into water (15 ml) and extracted with ether (30 ml x 3). The ether layer was extracted with 5% Na_2CO_3 (25 ml x 5) and the aqueous layer was acidified with 6 N HCl. After extraction with ether (50 ml x 3), the product was chromatographed on a silica gel (40 g) column (EtOAc-hexane, 2 : 3) to yield 19 (83 mg), mp 96-98 °C (prisms from CHCl_-hexane); UV: λ_{max} 235 nm (ϵ 12800); IR (KBr): ν_{max} 3170-2800, 1715, 1660, 1610, 1365, 1220, 1135, 1020, 975, $ma_{\Lambda}^{ma_{\Lambda}}$ 800, 740 cm⁻¹; ¹H NMR: 6 7.17 (1H, q, J=1.4 Hz), 2.35 (3H, s), 2.20 (3H, d, J=1.4 Hz), 10.3 (1H, br s, OH); HRMS: m/z 128.0480 (M⁺, calcd for $C_6H_8O_3$: 128.0473); MS: m/z (rel. int.) 128 (M⁺, 9), 113 (13), 110 (40), 85 (15), 82 (16), 43 (100), 41 (15), 39 (18%), together with its <u>z</u> isomer (271 mg), mp 98-100 °C (lit. mp 99-100 °C).

The keto acid <u>19</u> was reduced with NaBH₄ in MeOH to afford $(2\underline{E})$ -4-hydroxy-2-methyl-2pentenoic acid (20), a gum; UV: λ_{max} 210 nm (ϵ 10300); IR (CHCl₃): ν_{max} 3620-2450, 1690, 1650, 1380, 1270, 1160, 1060, 940 cm⁻¹; H NMR: δ 6.80 (1H, dq, J=8.0, 1.5 Hz), 6.10 (2H, br s, OH), 4.68 (1H, dq, J=8.0, 6.0 Hz), 1.88 (3H, d, J=1.5 Hz), 1.33 (3H, d, J=6 Hz); HRMS: m/z 115.0402 $(M^{+}-CH_{3}, calcd for C_{5}H_{7}O_{3}: 115.0395); MS: m/z (rel. int) 115 (M^{+}-CH_{3}, 17), 112 (55), 97 (50), 87$ (76), 69 (82), 45 (43), 43 (74), 41 (100), 39 (54%).

Acknowledgements - We are indebted to Dr. Yasuo Fujimoto, Institute of Physical and Chemical Research, and Mr. Hideaki Kondo, Taisho Pharmaceutical Co., LTD., for the measurement of the ¹H NMR (400 MHz) spectra.

REFERENCES

- S. Hasegawa, T. Miura, Y. Hirose and Y. Iitaka, <u>Chem. Letters</u>, <u>1985</u>, 1589.
 S. Hasegawa, N. Kaneko and Y. Hirose, <u>Phytochemistry</u>, in press.
 <u>MULTAN</u>, P. Main, M. M. Woolfson and G. Germain, <u>Acta Crystallogr</u>., Sect. A, <u>27</u>, 368 (1971).
- 4) The atomic coordinates will be compiled in Cambridge Crystallographic Database and the thermal parameters and Fo, Fc tables may be obtained from one of the authors (Y. I.) on request.
- PLUTO, in Cambridge Crystallographic Database, Cambridge Crystallographic Data 5) Centre, University Chemical Laboratory, Lensfield Road, Cambridge, CB2 1EW.
- 6) R. Scheffold and P. Dubs, <u>Helv. Chem. Acta</u>, <u>50</u>, 798 (1967).

1784