



# Preparation of all stereoisomers of 2-allyl-2-methyl-3-hydroxycyclopentanone by desymmetric processes based on a microbial oxidation and reduction system

Mikio Fujii <sup>a,\*</sup>, Minoru Takeuchi <sup>b</sup>, Hiroyuki Akita <sup>a</sup>, Kaoru Nakamura <sup>b</sup>

<sup>a</sup> Faculty of Pharmaceutical Sciences, Toho University, 2-2-1 Miyama, Funabashi, Chiba 247-8510, Japan

<sup>b</sup> Institute for Chemical Research, Kyoto University, Uji, Kyoto 611-0001, Japan

## ARTICLE INFO

### Article history:

Received 25 May 2009

Revised 13 June 2009

Accepted 16 June 2009

Available online 18 June 2009

## ABSTRACT

All stereoisomers of 2-allyl-3-hydroxy-2-methylcyclopentanones **2–5** were prepared in high conversion and in an optically pure form by microbial reduction and oxidation. The reduction of symmetric diketone **1** by *Geotrichum candidum* NBRC 4597 under anaerobic conditions gave **2** in 83% yield (98% conversion), >99% de, and >99% ee, whereas the reduction of **1** by *G. candidum* NBRC 5767 under aerobic conditions gave **3** in 75% yield (99% conversion), >99% de, and >99% ee. Oxidation of *meso*-diol **6** by *G. candidum* NBRC 5767 under aerobic conditions afforded **4** in 83% yield (99% conversion) and >99% ee, while oxidation of *meso*-diol **7** by *Mucor heimialis* IAM 6095 in the presence of cyclohexanone as a co-oxidant afforded **5** in 68% yield (75% conversion) and >99% ee.

© 2009 Elsevier Ltd. All rights reserved.


## 1. Introduction

Stereoselective and regioselective reduction of 2,2-dialkylcycloalkane-1,3-diones such as 2-allyl-2-methylcyclopentane-1,3-dione **1** is a useful method for the preparation of optically active compounds possessing a chiral quaternary carbon.<sup>1</sup> Desymmetrization of the *meso* compound allows 100% yield of the optically pure compound, while optical resolution gives only 50% yield.<sup>2</sup> Bakers' yeast reduction of 2,2-dialkylcycloalkane-1,3-dione is recognized as a valuable approach for accessing highly optically active 2,2-dialkyl-3-hydroxycycloalkanones through desymmetrization. For example, bakers' yeast reduction of **1** was reported to afford (2*S*,3*S*)-2-allyl-3-hydroxy-2-methylcyclopentanone **2** and (2*R*,3*S*)-2-allyl-3-hydroxy-2-methylcyclopentanone **3** in 75% yield as a diastereomeric mixture with 80% de (**2**:**3** = 9:1).<sup>3</sup> Optically pure **2** is a chiral synthon for corioliol, possessing antibacterial and antitumor activities (Scheme 1).<sup>3b</sup>

We have reported that a single microbe afforded both enantiomers of a secondary alcohol stereoselectively by changing the reaction conditions.<sup>4</sup> Oxygen concentration in the reaction atmosphere, for example, changes the stereochemical course of microbial reduction (Scheme 2). Under anaerobic conditions, reduction of acetophenone by *Geotrichum candidum* NBRC 5767 afforded (*S*)-1-phenylethanol in 98% yield and >99% enantiomeric excess (ee). In contrast, the reduction of ketone by *G. candidum* NBRC 5767 under aerobic conditions afforded (*R*)-1-phenylethanol in 99% yield and >99% ee, this phenomenon is possibly due to deracemization as shown in Scheme 2.<sup>4a</sup> The *S*-enzyme afforded the *S*-alcohol

reversibly but the *R*-enzyme afforded the *R*-alcohol irreversibly.<sup>5</sup> Since oxygen in air enhanced the oxidation of the *S*-alcohol, the *S*-alcohol produced by the reduction was re-oxidized to ketone in aerobic conditions, and finally the *R*-alcohol accumulated through irreversible reduction. In anaerobic conditions, oxidation of the *S*-alcohol was inhibited, and the faster reaction affording the *S*-alcohol was dominant. This procedure was considered useful for providing several enantiomers by only changing the reaction conditions. In this work we attempted to prepare an optically pure chiral synthon **2** of corioliol and its stereoisomers **3–5** using a microbial oxidation and reduction system (Fig. 1).


Compound **1** was reduced under normal conditions by *G. candidum* NBRC 5767 to obtain **2** in 92% conversion with 76% diastereomeric excess (de) (Table 1, entry 1). To control the diastereoselectivity of microbial reduction, a reduction system using XAD-7 under anaerobic conditions was employed (Table 1, entry 2).<sup>6</sup> However, the de of **2** decreased to 8%. Thus reduction under anaerobic conditions gave **2** in 80% conversion, 80% de, and >99% ee together with the minor stereoisomer **3** (>99% ee) (Table 1, entry 3). High diastereoselectivity and excellent enantioselectivity were observed on the



Scheme 1.

\* Corresponding author. Tel./fax: +81 47 472 1825.

E-mail address: [mfujii@phar.toho-u.ac.jp](mailto:mfujii@phar.toho-u.ac.jp) (M. Fujii).



Scheme 2.





Figure 1. Four stereoisomers of 2-allyl-3-hydroxy-2-methylcyclopentanone (2–5).

reduction of **1** by *G. candidum* NBRC 5767. It was found that the obtained **2** was the same major stereoisomer as that of bakers' yeast reduction. On the other hand, the reduction in aerobic conditions gave **3** in 95% conversion with 98% de and 99% ee (Table 1, entry 4). The reduction by *G. candidum* NBRC 5767 has been found to afford two stereoisomers by changing only reaction atmosphere.<sup>4</sup> The time course of the reaction under aerobic conditions is shown in Figure 2. Initially, reduction of **1** by *G. candidum* NBRC 5767 under aerobic conditions afforded **2** dominantly over **3**, the degree of conversion of **2** reached 60% after 6 h, and reduced with increasing reaction time. The degree of conversion of **3** transiently increased and reached 98% after 40 h. In order to improve the diastereoselectivity of the reduction affording **2**, another strain *G. candidum* NBRC 4597 was employed for the reduction of **1** to afford **2** in 98% conversion with 99% de and >99% ee (Table 1, entry 5). Finally, we succeeded in the preparation of **2** and **3** in high yield with satisfactory stereoselectivities. The high stereoselectivity of the reduction system under aerobic conditions was due to the high stereoselectivity of oxidation.<sup>6</sup> Moreover, the

Figure 2. Time course of the reduction of **1** by *G. candidum* 5767 under aerobic conditions.

enzyme catalyzing oxidation of **2** recognized the chirality of the adjacent quaternary carbon at the 2-position of **2**.

Table 1  
Stereochemical control on reduction of **1** by *G. candidum*


Entry	Microbe	Conditions	Conv. (%)	de (%)	ee (%) ( <b>2</b> )	ee (%) ( <b>3</b> )
1	<i>G. candidum</i> NBRC 5767		92	76 ( <b>2</b> )	>99	>99
2	<i>G. candidum</i> NBRC 5767	Ar, XAD-7 <sup>4</sup>	54	8 ( <b>2</b> )	>99	>99
3	<i>G. candidum</i> NBRC 5767	Ar <sup>4</sup>	80	80 ( <b>2</b> )	>99	>99
4	<i>G. candidum</i> NBRC 5767	Aerobic <sup>4</sup>	95	98 ( <b>3</b> )	— <sup>a</sup>	>99
5	<i>G. candidum</i> NBRC 4597	Ar <sup>4</sup>	98	>99 ( <b>2</b> )	>99	— <sup>b</sup>

Conditions: see Ref. 4 (reaction time: 24 h.).

GC analysis: after acetylation of products treated with acetic anhydride and pyridine, GC analysis was performed (Chirasil-DEX CB, 110 °C, 0.25 mm × 25 m, Ar: 2 mL/min). Retention time: **2**-acetate; 23.3 min, **5**-acetate; 24.4 min, **4**-acetate; 22.3 min, **3**-acetate; 19.8 min.

<sup>a</sup> Compound **5** was not observed.

<sup>b</sup> Compound **4** was not observed.



Scheme 3.

Reduction on a preparative scale was performed under both conditions. Reduction of **1** under anaerobic conditions gave **2** in 83% isolated yield and >99% ee (Scheme 3).<sup>7</sup> As with the reduction of **1** by *G. candidum* NBRC 5767 under anaerobic conditions, a prolonged reaction time was employed and **3** was obtained in 75% isolated yield (99% conv.) with >99% de and >99% ee.<sup>8</sup>

Next we attempted to obtain the other stereoisomers of **2**. Stereoselective and regioselective oxidation of *meso*-diols, such as (1*R*,2*S*,3*S*)-2-allyl-2-methylcyclopentane-1,3-diol (**6**) and (1*R*,2*R*,3*S*)-2-allyl-2-methylcyclopentane-1,3-diol (**7**), are also desymmetrization processes capable of producing optically active **4** and **5** in high yield. For the preparation of *meso*-diol substrates, **1** was reduced by excess NaBH<sub>4</sub> to afford **6** (31%), racemic (1*S*,3*S*)-2-allyl-2-methylcyclopentane-1,3-diol (41%), and **7** (15%).<sup>9</sup>


The oxidation of **6** by *G. candidum* NBRC 5767 under aerobic conditions afforded **4** in 99% conversion (83% isolated yield)<sup>10</sup> with >99% ee; further oxidation to diketone **1** was not observed. We succeeded in affording three stereoisomers (**2**, **3**, and **4**) in high yield and high ee using *G. candidum* (Scheme 3). Stereoselective oxidation of **7** by *G. candidum* NBRC 5767 as a pathway to **5** was unsuccessful. In the reduction of **1**, the enzyme catalyzing the oxidation of the alcohol appears to recognize the stereochemistry at the 2-position, leading to the oxidation of **7** to afford **5**. That is,

although alcohols possessing the (2*S*,3*S*) configuration, such as **2** and **6**, were oxidized by *G. candidum* NBRC 5767, alcohols possessing the (2*R*,3*S*) configuration, such as **3** and **7**, could not be oxidized by *G. candidum* NBRC 5767.

The ability of several microbes to afford **5** selectively was tested (Table 2). *Mucor heimalis* IAM 6095 (entry 5) had comparatively higher activity in affording **5** than the other microbes tested. In our experience, oxidation of secondary alcohols by *Mucor* species is not promoted by aerobic conditions. Thus, a biphasic system using cyclohexanone as a co-oxidant in a water–hexane co-solvent system was applied to the oxidation of **7** to enhance oxidation.<sup>11</sup> The addition of 45 mM cyclohexanone to the reaction system increased the conversion rate of **7** into **5** to 45% (entry 8). The addition of 90 mM cyclohexanone further increased the conversion rate up to 75% (entry 9), and optically pure **5** was isolated in 68% yield.<sup>12</sup>

In summary, we demonstrated a microbial oxidation and reduction system affording four different isomers of 2-allyl-3-hydroxy-2-methylcyclopentanones **2–5** in high conversion and in an enantiomerically pure form (Scheme 4). Moreover, by changing the reaction conditions, the reduction of **1** by *G. candidum* NBRC 5767 afforded two stereoisomers (**2** and **3**) and the oxidation of **6** by *G. candidum* NBRC 5767 under aerobic conditions afforded **4** (Scheme 3). In our previous study,<sup>4,5</sup> reversible reductases, S-en-


**Table 2**  
 Screening of microbes to afford **5**



Entry	Microbe	Conditions	Conv. (%)	ee (%)
1	<i>G. candidum</i> NBRC 5767	Aerobic <sup>4,5</sup>	0	–
2	<i>G. candidum</i> NBRC 4597	Aerobic <sup>4,5</sup>	7	38 ( <b>3</b> )
3	<i>G. candidum</i> ATCC 34614	Aerobic <sup>4,5</sup>	3	4 ( <b>3</b> )
4	<i>M. javanicus</i> IAM 6101	Aerobic <sup>4,5</sup>	13	>99 <sup>a</sup> ( <b>5</b> )
5	<i>M. heimalis</i> IAM 6095	Aerobic <sup>4,5</sup>	17	>99 <sup>a</sup> ( <b>5</b> )
6	<i>A. niger</i> IAM 4091	Aerobic <sup>4,5</sup>	0.4	>99 <sup>a</sup> ( <b>5</b> )
7	<i>D. magnusii</i> NBRC 4600	Aerobic <sup>4,5</sup>	3	>99 <sup>a</sup> ( <b>5</b> )
8	<i>M. heimalis</i> IAM 6095	Cyclohexanone: 45 M <sup>11</sup>	45	>99 <sup>a</sup> ( <b>5</b> )
9	<i>M. heimalis</i> IAM 6095	Cyclohexanone: 90 M <sup>11</sup>	75	>99 <sup>a</sup> ( <b>5</b> )

Conditions: see Refs. 4 and 5 for entries 1–7 and Ref. 11 for entries 8 and 9 (reaction time: 3 d).

<sup>a</sup> Compound **3** was not observed.



Scheme 4.

zymes, affording *S*-alcohols, and irreversible reductases, *R*-enzymes, were involved in the reaction to afford both enantiomers independently under different conditions. This work has found that the reversible reductase affording the *S*-alcohol recognizes the (2*S*,3*S*)-configuration of **2** and irreversible *S*-enzymes affording **3** are involved in the reaction. In fact, *S*-enzymes possessing different stereoselectivities on the reduction of ethyl 2-methyl-3-oxobutanoate have been reported.<sup>13</sup> Normally, a single microbe could not afford different isomers in high enantio- and diastereoselectivity. However the unique enantio- selective and diastereoselective oxidase in *G. candidum* was able to afford three isomers in high enantio- and diastereoselectivity. Application of the reversibility of the enzyme is a useful method for the preparation of optically active compounds in high yield.

## References and notes

- (a) Gibian, H.; Kieslich, K.; Koch, H. J.; Kosmol, H.; Rufer, C.; Schroeder, E.; Voessing, R. *Tetrahedron Lett.* **1966**, 21, 2321–2330; (b) Schwarz, S.; Truckenbrodt, G.; Meyer, M.; Zepter, R.; Weber, G.; Carl, C.; Wentzke, M.; Schick, H.; Welzel, H. P. *J. Prakt. Chem.* **1981**, 323, 729–736; (c) Brooks, D. W.; Grothaus, P. G.; Mazdiyasni, H. *J. Am. Chem. Soc.* **1983**, 105, 4472–4473; (d) Kitahara, T.; Miyake, M.; Kido, M.; Mori, K. *Tetrahedron: Asymmetry* **1990**, 1, 775–782; (e) Katoh, T.; Mizumoto, S.; Fudesaka, M.; Nakashima, Y.; Kajimoto, T.; Node, M. *Synlett* **2006**, 2176–2182; (f) Brooks, D. W.; Mazdiyasni, H.; Chakrabarti, S. *Tetrahedron Lett.* **1984**, 25, 1241–1244; (g) Wei, Z.-L.; Li, Z.-Y.; Lin, G.-Q. *Synthesis* **2000**, 1673–1676; (h) Shimoda, K.; Kubota, N.; Hamada, H. *Tetrahedron Lett.* **2006**, 47, 1541–1544.
- (a) Urdiales, E. G.; Alfonso, I.; Gotor, V. *Chem. Rev.* **2005**, 105, 313–354; (b) Schoffers, E.; Golebiowski, A.; Johnson, C. R. *Tetrahedron* **1996**, 52, 3769–3826; (c) Chikamatsu, H.; Taniguchi, M.; Kanemitsu, T. *Bull. Chem. Soc. Jpn.* **1986**, 59, 2663–2665; (d) Iriuchijima, S.; Hasegawa, K.; Tsuchihashi, G. *Agric. Biol. Chem.* **1982**, 46, 1907–1910; (e) Björkling, F.; Boutelje, J.; Gatenbeck, S.; Hult, K.; Norin, T.; Szmulik, P. *Tetrahedron* **1985**, 41, 1347–1352; (f) Schneider, M.; Engel, N.; Boensmann, H. *Angew. Chem., Int. Ed. Engl.* **1984**, 23, 66; (g) Watanabe, H.; Iwamoto, M.; Nakada, M. *J. Org. Chem.* **2005**, 70, 4652–4658; (h) Fujieda, S.; Tomita, M.; Fuhshuku, K.; Ohba, S.; Nishiyama, S.; Sugai, T. *Adv. Synth. Catal.* **2005**, 347, 1099–1109.
- (a) Brooks, D. W.; Mazdiyasni, H.; Grothaus, P. G. *J. Org. Chem.* **1987**, 52, 3223–3232; (b) Brooks, D. W.; Grothaus, P. G.; Irwin, W. L. *J. Org. Chem.* **1982**, 47, 2820–2821; (c) Brooks, D. W.; Grothaus, P. G.; Palmer, J. T. *Tetrahedron Lett.* **1982**, 23, 4187–4190.
- (a) Nakamura, K.; Takenaka, K.; Fujii, M.; Ida, Y. *Tetrahedron Lett.* **2002**, 43, 3629–3631; (b) Nakamura, K.; Fujii, M.; Ida, Y. In *Asymmetric Reduction of Ketones: Synthesis of Both Enantiomers of 1-Phenylethanol by Reduction of Acetophenone with Geotrichum candidum IFO 5767*; Roberts, S. M., Ehrtall, J., Eds.; Catalysts for Fine Chemical Synthesis; John Wiley: New York, 2007; Vol. 5, pp 93–97.
- Nakamura, K.; Fujii, M.; Ida, Y. *J. Chem. Soc., Perkin Trans. 1* **2000**, 3205–3211.
- Nakamura, K.; Fujii, M.; Ida, Y. *Tetrahedron: Asymmetry* **2001**, 12, 3147–3153.
- Compound **2**: From **1** (300 mg, 1.97 mmol) to (2*S*,3*S*)-**2** (255 mg, 1.66 mmol, 83%, >99% ee); [ $\alpha$ ]<sub>D</sub> +106.2 (c 0.43, CHCl<sub>3</sub>), lit.<sup>3a</sup> [ $\alpha$ ]<sub>D</sub> +80.7 (c 0.4, CHCl<sub>3</sub>). IR (neat)  $\nu$ : 3457, 1730, 1640 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz):  $\delta$  0.97 (s, 3H), 1.70–2.56 (m, 7H), 3.98–4.14 (m, 1H), 5.01–5.18 (m, 2H), 5.76–5.95 (m, 1H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz):  $\delta$  19.6, 27.7, 34.0, 35.3, 53.2, 77.3, 118.1, 134.3, 221.0; Anal. Calcd for C<sub>9</sub>H<sub>14</sub>O<sub>2</sub>: C, 70.10; H, 9.15. Found: C, 69.89; H, 9.15.
- Compound **3**: From **1** (100 mg, 0.64 mmol) to (2*R*,3*S*)-**3** (76 mg, 0.49 mmol, 75%, >99% ee); [ $\alpha$ ]<sub>D</sub> –81.0 (c 0.59, CHCl<sub>3</sub>), lit.<sup>3a</sup> [ $\alpha$ ]<sub>D</sub> –86.5 (c 0.26, CHCl<sub>3</sub>); IR (neat)  $\nu$ : 3434, 1732, 1640 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz):  $\delta$  0.98 (s, 3H), 1.78–1.194 (m, 1H) 2.01–2.50 (m, 6H), 4.48 (t, 1H, *J* = 6.4 Hz), 5.01–5.14 (m, 2H), 5.26–5.83 (m, 1H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz):  $\delta$  15.0, 27.5, 34.9, 39.7, 53.0, 75.2, 118.7, 133.5, 220.2; Anal. Calcd for C<sub>9</sub>H<sub>14</sub>O<sub>2</sub>: C, 70.10; H, 9.15. Found: C, 69.94; H, 9.09.
- Gaultieri, F.; Melchiorre, C.; Giannella, M.; Pignini, M. *J. Org. Chem.* **1975**, 40, 2241–2243.
- Compound **4**: From (1*R*,2*R*,3*S*)-**6** (100 mg, 0.64 mmol) to (2*R*,3*R*)-**4** (81 mg, 0.53 mmol, 83%, >99% ee). [ $\alpha$ ]<sub>D</sub> –105.0 (c 0.58, CHCl<sub>3</sub>). Spectral data including <sup>1</sup>H NMR and <sup>13</sup>C NMR were identical to those of **2**.
- Nakamura, K.; Inoue, Y.; Matsuda, T.; Misawa, I. *J. Chem. Soc., Perkin Trans. 1* **1999**, 2397–2402.
- Compound **5**: From (1*R*,2*R*,3*S*)-**7** (120 mg, 0.77 mmol) to (2*R*,3*R*)-**5** (80 mg, 0.52 mmol, 68%, >99% ee). [ $\alpha$ ]<sub>D</sub> +85.4 (c 0.58, CHCl<sub>3</sub>). Spectral data including <sup>1</sup>H NMR and <sup>13</sup>C NMR were identical to those of **3**.
- Kawai, Y.; Kawanobe, K.; Ohno, A. *Bull. Chem. Soc. Jpn.* **1997**, 70, 1683–1686.