

[Chem. Pharm. Bull.]
[34(9)3915—3918(1986)]

Studies on Peptides. CXLII.^{1,2)} Synthesis of Des-1-Ala-des- α -amino-Human Calcitonin Gene-Related Peptide

AKIRA OTAKA,^a NOBUTAKA FUJII,^a SUSUMU FUNAKOSHI,^a ITSUO YAMAMOTO,^b
KANJI TORIZUKA,^b TOSHIHARU NODA,^c KAORU MORITA,^c
and HARUAKI YAJIMA^{*,a}

Faculty of Pharmaceutical Sciences, Kyoto University,^a Department of Nuclear Medicine,
University Hospital, Kyoto University,^b Sakyo-ku, Kyoto 606, Japan and
Research Laboratories, Toyo Jozo Co., Ltd.,^c Mifuku,
Ohito-cho, Shizuoka 410-23, Japan

(Received March 7, 1986)

Subsequent to our synthesis of human calcitonin gene-related peptide (hCGRP) (*J. Chem. Soc., Chem. Commun.*, **1985**, 602), its analog, des-1-Ala-des- α -amino-hCGRP, was synthesized. This peptide, prepared by replacement of the N-terminal Ala-Cys residues with 3-mercaptopropionic acid, followed by air oxidation, was more active than the parent peptide in respect of Ca-lowering and Pi-lowering activities in rat serum.

Keywords—human CGRP; hCGRP analog; *S*-*p*-methoxybenzyl-3-thiopropionic acid; thioanisole-mediated deprotection; serum Ca-lowering activity; serum Pi-lowering activity

Human calcitonin gene related peptide (hCGRP) is a 37-residue peptide amide with one disulfide bridge, characterized by Morris *et al.*³⁾ in 1984. Recently we reported a solution synthesis of hCGRP,⁴⁾ for which seven peptide fragments were selected as building blocks to construct the entire amino acid sequence of this new peptide. Using available fragments [1] to [6] and a newly prepared N-terminal fragment, we have synthesized des-1-Ala-des- α -amino-hCGRP (Fig. 1), which may be resistant to degradation by the action of aminopeptidase. As expected, this analog of hCGRP exhibited much higher activity than the parent molecule in respect of Ca-lowering activity in rat serum.

The N-terminal fragment [7], MBzl-S-CH₂-CH₂-CO-Asp-Thr-Ala-NHNH₂, was syn-




Fig. 1. Synthetic Route to Des-1-Ala-des- α -amino-hCGRP




Fig. 2. Synthetic Scheme for the N-Terminal Fragment [7]




Fig. 3. Biological Activities of the Synthetic hCGRP Analog

a: Effect on Ca in serum.

b: Effect on Pi in serum.

—○—, synthetic hCGRP (80 $\mu\text{g/kg}$ i.v.); —●—, des-Ala-des-amino analog (80 $\mu\text{g/kg}$ i.v.); —×—, vehicle (pH 6.0, 0.38 mmol citric acid buffer containing 0.1% BSA).

thesized according to the scheme shown in Fig. 2. First, a TFA-treated sample of Z(OMe)-Asp(OBzl)-Thr-Ala-OMe⁴⁾ was subjected to hydrogenolysis in order to remove the Bzl ester. Thr, was avoided. The resulting tripeptide ester, H-Asp-Thr-Ala-OMe, was allowed to react with MBzl-S-CH₂-CH₂-COOSu⁶⁾ to give MBzl-S-CH₂-CH₂-CO-Asp-Thr-Ala-OMe, which was converted to [7] by the usual hydrazine treatment. Next, as an amino component, the protected dotriacontapeptide amide, Z(OMe)-(hCGRP 6-37)-NH₂, was prepared by condensations of fragments [1] to [6] as reported previously.⁴⁾ This, after TFA treatment, was condensed with fragment [7] by the azide procedure⁷⁾ and the product was purified by gel-filtration on Sephadex LH-60 using DMF-DMSO (7:3) as an eluant.

The protected peptide amide thus obtained was treated with 1 M TFMSA-thioanisole in TFA⁸⁾ in the presence of *m*-cresol and the desired product was isolated as described in the case of hCGRP synthesis,⁴⁾ i.e., as follows. 1. Reduction with 2-mercaptoethanol. 2. Incubation at pH 8.0 to reverse a possible N→O shift.⁹⁾ 3. Gel-filtration on Sephadex G-25 to remove the reducing reagent. 4. Air-oxidation in a dilute solution at pH 7.5 to establish the intramolecular disulfide bond. 5. Repeated lyophilization to obtain a crude product. 6. Purification by ion-exchange chromatography on CM-Biogel A using ammonium acetate buffer. 7. Purification by high performance liquid chromatography (HPLC) using gradient elution with acetonitrile (25-40%) in 0.1% TFA. The hCGRP analog thus purified exhibited a single spot on thin layer chromatography (TLC) and a single peak in analytical HPLC. Its acid hydrolysate gave the amino acids in ratios predicted by theory.

In respect of the Ca-lowering activity in rat serum, this hCGRP analog (80 $\mu\text{g/kg}$) exhibited much higher activity than the synthetic hCGRP. A similar tendency was also observed in respect of Pi-lowering activity in rat serum (Fig. 3).

Experimental

General experimental procedures described herein are essentially the same as described in the previous synthesis of hCGRP.⁴⁾ R_f values in TLC performed on silica gel (Kieselgel G, Merck) refer to the following solvent systems: R_{f1} CHCl_3 -MeOH- H_2O (8:3:1), R_{f2} n -BuOH-AcOH-pyridine- H_2O (4:1:1:2), R_{f3} n -BuOH-AcOH-AcOEt- H_2O (1:1:1:1). HPLC was conducted with a Waters 204 compact model.

MBzl-S-CH₂-CH₂-CO-Asp-Thr-Ala-OMe—A TFA-treated sample of Z(OMe)-Asp(OBzl)-Thr-Ala-OMe⁴⁾ (1.00 g, 1.74 mmol) in DMF (30 ml) was hydrogenated over a Pd catalyst for 3 h and the catalyst was removed by filtration. Et₃N (0.53 ml, 3.83 mmol) and MBzl-S-CH₂-CH₂-COOSu (0.68 g, 2.09 mmol) were added to the filtrate and the mixture was stirred for 12 h, then concentrated. The residue was dissolved in AcOEt. The organic phase was washed with 5% citric acid and H_2O -NaCl, dried over Na₂SO₄ and concentrated. Trituration of the residue with ether afforded a powder which was recrystallized from AcOEt and ether; yield 0.56 g (61%), mp 140–142 °C, $[\alpha]_D^{18}$ -10.2° (c =0.6, DMF), R_{f1} 0.47. Anal. Calcd for C₂₃H₃₃N₃O₉S: C, 52.36; H, 6.30; N, 7.97. Found: C, 52.09; H, 6.44; N, 7.70.

MBzl-S-CH₂-CH₂-CO-Asp-Thr-Ala-NHNH₂—The above ester (0.55 g, 0.95 mmol) in DMF-MeOH (1:1, 50 ml) was treated with 80% hydrazine hydrate (0.57 ml, 10 eq) at room temperature for 24 h. The solvent was removed by evaporation and the residue was treated with EtOH to afford a powder, which was recrystallized from DMF and EtOH; yield 0.44 g (87%), mp 178–180 °C, $[\alpha]_D^{18}$ -16.0° (c =1.0, DMF), R_{f1} 0.21. Amino acid ratios in a 6 N HCl hydrolysate: Asp 1.06, Thr 0.97, Ala 1.00 (recovery of Ala 81%). Anal. Calcd for C₂₂H₃₃N₅O₈S·H₂O: C, 48.43; H, 6.47; N, 12.84. Found: C, 48.69; H, 6.46; N, 13.17.

Protected Des-1-Ala-des- α -amino-hCGRP, MBzl-S-CH₂-CH₂-CO-(hCGRP 3–37)-NH₂—Z(OMe)-(hCGRP 6–37)-NH₂ (270 mg, 60 μ mol) was treated with TFA (3.0 ml) in the presence of anisole (0.3 ml) in an ice-bath for 2 h, then dry ether was added. The resulting powder was collected by filtration, dried over KOH pellets *in vacuo* for 3 h and dissolved in DMF-DMSO (2:1, 5 ml) containing Et₃N (17 μ l, 120 μ mol). The azide [prepared from 157 mg (298 μ mol) of MBzl-S-CH₂-CH₂-CO-Asp-Thr-Ala-NHNH₂] in DMF (5 ml) and Et₃N (124 μ l, 894 μ mol) were added to the above ice-chilled solution and the mixture, after being stirred at 4 °C for 48 h, was poured into H₂O (200 ml). The resulting powder was washed with MeOH and purified by gel-filtration on Sephadex LH-60 (3.2 $\times$ 136 cm) using DMF-DMSO (7:3) as an eluant. The desired fractions corresponding to the front peak (tube Nos. 50–76, 10 ml each, determined by ultraviolet (UV) absorption measurement at 275 nm) were combined and the solvent was removed by evaporation. The residue was treated with AcOEt to form a powder; yield 182 mg (62%), mp 271–273 °C, $[\alpha]_D^{18}$ -19.7° (c =0.3, DMSO), R_{f1} 0.42. Amino acid ratios in a 6 N HCl hydrolysate: Asp 4.37, Thr 3.78, Ser 3.10, Pro 0.98, Gly 4.33, Ala 3.67, Val 3.46, Leu 3.36, Phe 2.00, Lys 2.01, His 1.00, Arg 2.11 (recovery of Phe 81%). Anal. Calcd for C₂₃₃H₃₃₅N₄₉O₅₇S₄·7H₂O: C, 56.09; H, 7.05; N, 13.76. Found: C, 56.18; H, 6.97; N, 13.96.

Des-1-Ala-des- α -amino-hCGRP—The above protected peptide (80 mg, 16.5 μ mol) was treated with 1 M TFMSA-thioanisole in TFA (6.6 ml) in the presence of *m*-cresol (0.34 ml, 200 eq) in an ice-bath for 2 h, then dry ether was added. The resulting powder was collected by centrifugation, dried over KOH pellets *in vacuo* for 1 h and dissolved in 6 M guanidine-HCl in 0.1 M Tris-HCl buffer (pH 8.0, 4.0 ml) containing 2-mercaptoethanol (0.2 ml). The pH of the solution was adjusted to 8.0 with MeNH₂ and the solution was incubated under an argon atmosphere at room temperature for 24 h. This solution was applied to a column of Sephadex G-25 (2.8 $\times$ 138 cm), which was eluted with 1 N AcOH. The fractions corresponding to the main peak (tube Nos. 47–70, 9 ml each, determined by UV absorption measurement at 206 nm) were combined and the entire solution was diluted with ice-chilled H₂O to 1000 ml. The pH of the solution was adjusted to 7.5 with 5% NH₄OH and the solution was kept standing at 23 °C for 7 d, during which time the Ellman test values (412 nm)¹⁰⁾ dropped from 0.091 to 0.019. The pH of the solution was adjusted to 5.0 with AcOH and then the solvent and the salt were removed by repeated lyophilization to give a powder; yield 40.2 mg (66%).

The crude air-oxidized product thus obtained was dissolved in 0.02 M AcONH₄ buffer (pH 5.8, 3 ml) and the solution was applied to a column of CM-Biogel A (2 $\times$ 7 cm), which was eluted first with the same buffer (192 ml) and then with a linear gradient formed from 0.2 M AcONH₄ buffer (pH 6.8, 250 ml) through a mixing flask containing the starting buffer (250 ml). The fractions corresponding to the main peak (tube Nos. 69–84, 4 ml each, determined by measuring UV absorption at 206 nm and by the Folin-Lowry test¹¹⁾) were combined and the solvent and the ammonium salt were removed by repeated lyophilization to give a fluffy powder; yield 16.1 mg (40%). Subsequent purification was performed by reversed-phase HPLC on a Cosmosil 5C₁₈ column (10 $\times$ 250 mm). A part of the above CM-purified sample (6.0 mg) was dissolved in 0.1% TFA (0.6 ml) and the solution was applied to the column, which was eluted with a gradient of acetonitrile (25 to 40% in 1 h) in 0.1% TFA at a flow rate of 1.8 ml per min (Fig. 4-a). The eluate corresponding to the main peak (retention time 44 min) was collected and the solvent was removed by lyophilization to give a fluffy white powder; yield 1.78 mg. The rest of the sample was similarly purified; total yield 4.8 mg. The over all yield from the protected peptide was 7.8%. $[\alpha]_D^{20}$ -101.5° (c =0.1, 1 N AcOH), R_{f2} 0.32, R_{f3} 0.59. The retention time was 20 min in HPLC on an analytical Nucleosil 5C₁₈ column (4 $\times$ 150 mm) on gradient elution with acetonitrile (25 to 40%) in 0.1% TFA at a flow rate of 1 ml per min (Fig. 4-b). Amino acid ratios in a 6 N HCl hydrolysate (numbers in parentheses are theoretical): Asp 3.71 (4), Thr 2.88 (4), Ser 2.79 (3), Pro 1.03 (1), Gly 4.35 (4),




Fig. 4. HPLC of the Synthetic hCGRP Analog

a: CM-purified sample.
b: purified sample.

Cys N. D., Ala 3.13 (3), Val 3.97 (5), Leu 3.13 (3), Phe 2.00 (2), Lys 1.98 (2), His 0.89 (1), Arg 2.15 (2) (recovery of Phe 90%).

References and Notes

- 1) Part CXLI: M. Sakurai, K. Akaji, N. Fujii, M. Moriga, M. Aono, K. Mizuta, H. Adachi, M. Noguchi, K. Inoue, R. Hosotani, T. Tobe, and H. Yajima, *Chem. Pharm. Bull.*, **34**, 3447 (1986).
- 2) Amino acids used in this investigation are of the L-configuration. The following abbreviations are used: Ad = 1-adamantyl, Z(OMe) = *p*-methoxybenzyloxycarbonyl, Z = benzyloxycarbonyl, Mts = mesitylenesulfonyl, Bzl = benzyl, MBzl = *p*-methoxybenzyl, TFMSA = trifluoromethanesulfonic acid, TFA = trifluoroacetic acid, Su = *N*-hydroxysuccinimidyl, DMSO = dimethylsulfoxide, DMF = dimethylformamide, BSA = bovine serum albumin.
- 3) H. R. Morris, M. Panico, T. Etienne, J. Tippins, S. I. Girgis, and I. MacIntyre, *Nature* (London), **308**, 746 (1984).
- 4) N. Fujii, H. Yajima, A. Otaka, S. Funakoshi, M. Nomizu, K. Akaji, I. Yamamoto, K. Torizuka, K. Kitagawa, T. Akita, K. Ando, T. Kawamoto, Y. Shimonishi, and T. Takao, *J. Chem. Soc., Chem. Commun.*, **1985**, 602.
- 5) J. Martinez and M. Bodanszky, *Int. J. Peptide Protein Res.*, **12**, 277 (1978) and references cited therein.
- 6) N. Fujii, K. Akaji, Y. Hayashi, and H. Yajima, *Chem. Pharm. Bull.*, **33**, 362 (1985).
- 7) J. Honzl and J. Rudinger, *Collect. Czech. Chem. Commun.*, **26**, 2333 (1961).
- 8) H. Yajima, N. Fujii, H. Ogawa, and H. Kawatani, *J. Chem. Soc., Chem. Commun.*, **1974**, 107; H. Yajima and N. Fujii, *J. Am. Chem. Soc.*, **103**, 5867 (1981).
- 9) S. Sakakibara, in "Chemistry and Biochemistry of Amino Acids, Peptides and Proteins," Vol. 1, ed. B. Weinstein and M. Dekker, New York, 1971, p. 51.
- 10) G. L. Ellman, *Arch. Biochem. Biophys.*, **82**, 70 (1959).
- 11) O. H. Lowry, N. J. Rosebrough, A. L. Farr, and R. J. Randall, *J. Biol. Chem.*, **193**, 215 (1951).