

A Stable Enediol Sulphite: an Example of Steric Protection of Unstable Compounds



By A. DE GROOT, J. A. BOERMA, and HANS WYNBERG*

(Department of Organic Chemistry of the University, Bloemsingel 10, Groningen, The Netherlands)

DURING our studies on the 3,3,6,6-tetramethylthiepan system^{1,2} we attempted to prepare 5-chloro-3,3,6,6-tetramethyl-4-oxothiepan, by refluxing the hydroxy-ketone (I)¹ for 2 hr. with thionyl chloride and dimethylaniline, in chloroform as solvent. Instead of the chloro-ketone, the enediol sulphite (II) was isolated as a stable solid (m.p. 86–88°, 84% yield) which was characterized as 4,5,7,8-tetrahydro-4,4,8,8-tetramethylthiepano[4,5,-d]-1,3,2-dioxathiole 2-oxide by elemental analysis (C₁₀H₁₆O₃S₂) and by spectra. The i.r. spectrum showed an absorption at 1670 cm.⁻¹, indicating a double bond. The n.m.r. spectrum (10%, CCl₄) showed a singlet at τ 7.36 (4 methylene protons) and two singlets at τ 8.73 and 8.68 (12 methyl protons). The u.v. spectrum showed no absorption above 220 m μ . The mass spectrum showed a

parent peak at 248 and the fragmentation pattern is in accord with the assigned structure. Pyrolysis of the enediol sulphite at 200–225° gave the diketone (III)¹ in 75% yield.

Enediol sulphites were postulated as intermediates in the reaction of benzoin with thionyl chloride by Fieser and Okumura.³ The latter⁴ prepared the dimesityl compound (IV) which showed an absorption maximum at 260 m μ (ϵ 10650). This, however, is attributable to the *cis*-hexamethylstilbene chromophore (*cf.* refs. 5 and 6), rather than to the enediol sulphite group, as was suggested by Okumura.⁴



The remarkable stability of compound (II), the first known example of a stable alicyclic enediol sulphite, is an example of stabilisation by steric protection such as the reported protection by *t*-butyl groups or α -gem-dimethyl groups of several reactive compounds.⁷

(Received, February 5th, 1968; Com. 143.)

¹ A. de Groot and H. Wynberg, *J. Org. Chem.*, 1966, **31**, 3954.

² A. de Groot, B. Evenhuis, and H. Wynberg, *J. Org. Chem.*, 1968, **33**, in the press.

³ L. F. Fieser and Y. Okumura, *J. Org. Chem.*, 1962, **27**, 2247.

⁴ Y. Okumura, *J. Org. Chem.*, 1963, **28**, 1075.

⁵ M. T. O'Shaughnessy and W. H. Rodebush, *J. Amer. Chem. Soc.*, 1940, **62**, 2906.

⁶ K. A. Muszkat, D. Gégion, and E. Fischer, *Chem. Comm.*, 1965, 447.

⁷ A. K. Hoffmann, *J. Amer. Chem. Soc.*, 1961, **83**, 4671; M. S. Newman, A. Arkell, and T. Fukunaya, *ibid.*, 1960, **82**, 2498; E. E. van Tamelen and S. P. Pappas, *ibid.*, 1963, **85**, 3297; C. Hoogzand and W. Hübel, *Tetrahedron Letters*, 1961, 637; J. B. Pazos and F. D. Greene, *J. Amer. Chem. Soc.*, 1967, **89**, 1030; J. C. Sheehan and J. H. Beesan, *ibid.*, p. 362; J. E. Bennett, H. Sieper, and P. Tavs, *Tetrahedron*, 1967, **23**, 1697; U. E. Wiersum and H. Wynberg, *Tetrahedron Letters*, 1967, 2951.