



Synthesis and Biological Evaluation of the Four Racemic Stereoisomers of the Structure Proposed for Sorgolactone, the Germination Stimulant from *Sorghum bicolor*

Kenji Mori^{a*}, Junichi Matsui^{a†}, Masahiko Bando^b, Masaru Kido^b and Yasutomo Takeuchi^c

^aDepartment of Chemistry, Faculty of Science, Science University of Tokyo, Kagurazaka 1-3, Shinjuku-ku, Tokyo 162, Japan

^bSecond Tokushima Institute of New Drug Research, Otsuka Pharmaceutical Co., Ltd., Kawauchi, Tokushima 771-01, Japan

^cWeed Science Center, Utsunomiya University, Mine-machi, Utsunomiya, Tochigi 321, Japan

Abstract: The synthesis of the racemates of the structure **1** proposed for sorgolactone and its three stereoisomers was achieved by confirming the stereostructures of the intermediate ($\pm$)-**6** and the final product ($\pm$)-**1** by X-ray analyses. Biological evaluation of the products employing clover broomrape (*Orobancha minor*) seeds revealed that the order of activity as a germination stimulant was ($\pm$)-strigol = ($\pm$)-**9** $\geq$ ($\pm$)-**12** > ($\pm$)-**1** > ($\pm$)-**11**. © 1997 Elsevier Science Ltd.

Parasitic weeds of the genera *Orobancha* and *Striga* are known to cause severe yield losses in grains and legumes in Africa, Asia and the U. S. A.^{1,2} The seeds of such weeds remain dormant in soil until exudates from their host plant induce germination. (+)-Strigol was first isolated from cotton root exudates and shown to be a strong stimulant for the germination of such seeds³. It was later isolated from the host plants such as maize, proso millet and sorghum⁴. Synthesis of (+)-strigol by Sih⁵ and that of ($\pm$)-strigol by Raphael⁶ evoked much interest in the chemistry of germination stimulants, which culminated in the determination of the absolute configuration of the naturally occurring (+)-strigol by X-ray analysis⁷. The synthetic endeavor in this area is still remarkable^{8,9}.

In 1992 Hauck et al. isolated sorgolactone from *Sorghum bicolor*, a genuine host plant, as the germination stimulant for parasitic weeds¹⁰. They proposed **1** as the structure of sorgolactone based on its ¹H NMR analysis and CD comparison with strigol¹⁰. We herein report the first synthesis of ($\pm$)-**1**. Two attempted syntheses of **1** have been reported recently^{11,12}.

Our synthesis of ($\pm$)-**1** is summarized in Figure 1. The sodium salt of the known diketo acid **2** (a racemic and diastereomeric mixture)¹³ was reduced with sodium borohydride in the presence of cerium(III) chloride¹⁴ to give, after acidification, a racemic and diastereomeric mixture of the hydroxy lactone **3**¹⁵. Treatment of **3** with carbon tetrabromide and triphenylphosphine yielded a racemic and diastereomeric mixture of the unstable bromolactone **4**, which was reduced with zinc-copper couple (Zn:Cu = 91:5, Kanto Chemical Co.)¹⁶ and acetic acid in THF to furnish the crude reduction product. This was dissolved in chloroform and stirred for 60 hr at room temperature to cause the double bond migration to the more stable tetrasubstituted position¹⁶. The product was further purified by MPLC [Lobar LiChroprep[®] Si 60 (40-63 μ m)] to give an oily fast moving




Figure 1. Synthesis of (±)-1

Reagents: (a) NaOH, NaBH₄, CeCl₃·7H₂O, CH₂Cl₂/H₂O (67%).—(b) CBr₄, Ph₃P, CH₂Cl₂ (87%).—(c) 1) Zn—Cu, AcOH, THF; 2) CHCl₃, stirring (40% as a mixture of 5 and 6); 3) MPLC separation.—(d) NaH, HCO₂Et, Et₂O (quant.).—(e) K₂CO₃, *N*-methylpyrrolidone; SiO₂ chromatog. separation [42% for (±)-1, 41% for (±)-9, 39% for (±)-11 and 45% for (±)-12].




Figure 2. Perspective views of ($\pm$)-6 and ($\pm$)-1


diastereomer and a crystalline slowly moving one. The latter (m.p. 43–46 °C) was submitted to the X-ray analysis¹⁷ to reveal its structure as (±)-**6**. Its computer-generated stereoview is shown in Figure 2. The oily lactone must therefore be (±)-**5**, which leads to (±)-**1**. These two stereoisomeric lactones were separately processed to give (±)-**1** and its three stereoisomers.

Formylation of ($\pm$)-**5** with sodium hydride and ethyl formate gave ($\pm$)-**7**, m.p. 114–116 °C. Treatment of ($\pm$)-**7** with ($\pm$)-4-bromo-2-methyl-2-buten-4-olide [($\pm$)-**8**] in the presence of potassium carbonate gave a

mixture of ($\pm$)-1 and ($\pm$)-9, which could be separated by silica gel chromatography to give two crystalline products. The structure of one of the products with m.p. 127–129 °C was solved by X-ray analysis¹⁸, and its stereoview is shown in Figure 2. This product was thus ($\pm$)-1¹⁹, and the other with m.p. 117–119 °C was ($\pm$)-9. Similarly, the stereoisomeric lactone ($\pm$)-6 was converted to ($\pm$)-11, m.p. 131–133 °C, and ($\pm$)-12, m.p. 116–118 °C. These structures were tentatively assigned on the basis of their *R_f* values.

The ¹H NMR spectra of ($\pm$)-1, ($\pm$)-9, ($\pm$)-11 and ($\pm$)-12 in CDCl₃ or C₆D₆ were measured and compared with the copies (sent by Dr. C. Hauck) of the spectra of the natural sorgolactone. As shown in Table 1, although they were all very similar to each other, the spectrum of ($\pm$)-1 was almost identical with that of the natural product. Unfortunately, Dr Hauck's spectra of sorgolactone were of poor quality due to the scarcity of the sample and the presence of impurities. Because the natural product itself is no more available, we conclude that the structure of sorgolactone is most probably 1 and its reisolation must be attempted to completely settle the problem by direct TLC comparison of ($\pm$)-1 with the reisolated sample.

Bioactivity of strigol, its stereoisomers or structural analogues as a germination stimulant is known to be strongly dependent on its structure including relative and absolute stereochemistry^{20–22}. We therefore carried out the bioassay of four final products by using seeds of clover broomrape (*Orobancha minor*) as the test parasitic weed²³. All of four products stimulated the germination of the *Orobancha* seeds, and the order of the stimulant activity was ($\pm$)-strigol $\approx$ ($\pm$)-9 $\geq$ ($\pm$)-12 > ($\pm$)-1 > ($\pm$)-11 as shown in Table 1. The details of this work will appear in *Liebigs Ann.*



m.p.	127–129 °C	117–119 °C	131–133 °C	116–118 °C
TLC(hexane-EtOAc 1:1)	<i>R_f</i> 0.42	<i>R_f</i> 0.34	<i>R_f</i> 0.42	<i>R_f</i> 0.33
1H-NMR Data (Our data were measured at 500 MHz as CDCl ₃ solutions.)				
	<i>lit.</i> ¹⁰			
8-Me	1.1 (3H)	1.06 (3H, d, <i>J</i> =7.0 Hz)	1.04 (3H, d, <i>J</i> =7.0 Hz)	1.13 (3H, d, <i>J</i> =7.0 Hz)
7-H		1.24 (1H, m)	1.23 (1H, m)	1.35 (1H, m)
7-H'		1.56 (1H, m)	1.55 (1H, m)	1.55 (1H, m)
6-H		1.70, 1.78 (2H, m)	1.68, 1.77 (2H, m)	1.72 (2H, m)
5-H		1.94 (2H, m)	1.93 (2H, m)	1.96 (2H, m)
4'-Me	2.0 (3H)	2.03 (3H, t, <i>J</i> =1.5 Hz)	2.01 (3H, t, <i>J</i> =1.5 Hz)	2.03 (3H, t, <i>J</i> =1.5 Hz)
4-H	2.4 (1H)	2.34 (1H, d, <i>J</i> =16.5 Hz)	2.33 (1H, d, <i>J</i> =16.5 Hz)	2.32 (1H, m)
8-H		2.38 (1H, m)	2.36 (1H, m)	2.37 (1H, d, <i>J</i> =16.5 Hz)
4-H'	2.7 (1H)	2.75 (1H, dd, <i>J</i> = 15.0, 9.0 Hz)	2.72 (1H, dd, <i>J</i> = 15.5, 9.0 Hz)	2.70 (1H, ddd, <i>J</i> = 16.5, 9.5, 3.4 Hz)
3a-H		3.63 (1H, m)	3.60 (1H, m)	3.62 (1H, m)
8b-H	5.5 (1H)	5.49 (1H, d, <i>J</i> =8.0 Hz)	5.48 (1H, d, <i>J</i> =7.5 Hz)	5.35 (1H, d, <i>J</i> =7.5 Hz)
3'-H	6.1 (1H)	6.15 (1H, t, <i>J</i> =1.5 Hz)	6.14 (1H, s)	6.14 (1H, t, <i>J</i> =1.5 Hz)
2'-H	6.9 (1H)	6.92 (1H, t, <i>J</i> =1.5 Hz)	6.93 (1H, t, <i>J</i> =1.5 Hz)	6.92 (1H, t, <i>J</i> =1.5 Hz)
9-H	7.4 (1H)	7.41 (1H, d, <i>J</i> =3.0 Hz)	7.42 (1H, d, <i>J</i> =3.0 Hz)	7.41 (1H, d, <i>J</i> =2.5 Hz)
Concentration (M)	Relative germination of <i>Orobancha minor</i> (%)			
10 ⁻⁵	82, 90	95, 95	70, 57	92, 95 (cf.)
10 ⁻⁶	77, 65	93, 93	12, 18	92, 95 (±)-strigol
10 ⁻⁷	40, 51	90, 90	7, 3	87, 78 94, 94
10 ⁻⁸	21, 23	84, 87	3, 0	80, 70 86, 85

Table 1. Physical and biological data of ($\pm$)-1 and its stereoisomers

Acknowledgment: We thank Dr. C. Hauck (Novartis, Basel) for kindly sending to us the files of ^1H NMR spectrum of the natural **1** in C_6D_6 and a copy of his doctoral dissertation in which the ^1H NMR spectra of **1** in CDCl_3 and C_6D_6 are recorded. This work was financially supported by Kanebo Co., Ltd..

References and Notes

[†]Research fellow on leave from Kanebo Co., Ltd. (1996-1998).

- Parker, C.; Riches, C. R. *Parasitic Weeds of the World, Biology and Control*, CAB International, Wallingford, Oxon., U. K., 1993.
- Sand, P. F.; Eplee, R. E.; Westbrook, R. G. *Witchweed Research and Control in the United States*, Weed Science Society of America, 1990.
- Cook, C. E.; Whitchard, L. P.; Wall, M. E.; Egley, G. H.; Coggon, P.; Luhan, P. A.; McPhail, A. T. *J. Am. Chem. Soc.* **1972**, *94*, 6198-6199.
- Siame, B. A.; Weerasuriya, Y.; Wood, K.; Ejeta, G.; Butler, L. G. *J. Agric. Food Chem.* **1993**, *41*, 1486-1491.
- Heather, J. B.; Mittal, R. S. D.; Sih, C. J. *J. Am. Chem. Soc.* **1976**, *98*, 3661-3669.
- MacAlpine, G. A.; Raphael, R. A.; Shaw, A.; Taylor, A. W.; Wild, H.-J. *J. Chem. Soc., Perkin Trans. 1* **1976**, 410-416.
- Brooks, D. W.; Bevinakatti, H. S.; Powell, D. R. *J. Org. Chem.* **1985**, *50*, 3779-3781.
- Thuring, J. W. J. F.; Neffens, G. H. L.; Wegman, M. A.; Klunder, A. J. H.; Zwanenburg, B. *J. Org. Chem.* **1996**, *61*, 6931-6935.
- Kranz, A.; Samson-Schulz, E.; Henning, L.; Welzel, P.; Müller, D.; Mayer-Figge, H.; Sheldrick, W. S. *Tetrahedron* **1996**, *52*, 14827-14840.
- Hauck, C.; Müller, S.; Schildeknecht, H. *J. Plant Physiol.* **1992**, *139*, 474-478.
- Schröer, J.; Welzel, P. *Tetrahedron* **1994**, *50*, 6839-6858.
- Mikló, K.; Jaszberenyi, J. Cs.; Kádas, I.; Árvai, G.; Töke, L. *Tetrahedron Lett.* **1996**, *37*, 3491-3494.
- Kádas, I.; Árvai, G.; Töke, L. Tóth, G.; Szöllösy, Á.; Bihari, M. *Tetrahedron* **1994**, *50*, 2895-2906.
- Luche, J.-L. *J. Am. Chem. Soc.* **1978**, *100*, 2226-2227.
- All the new compounds were characterized by spectroscopic (IR and NMR) and elemental (combustion or HRMS) analyses.
- Frischmuth, K.; Samson, E.; Kranz, A.; Welzel, P.; Meuer, H.; Sheldrick, W. S. *Tetrahedron* **1991**, *47*, 9793-9806.
- X-ray analysis of ($\pm$)-**6**: Crystal size, 0.2 x 0.5 x 0.8 mm. All the data were obtained on Rigaku AFC-5S automated four circle diffractometer with graphite-monochromated Mo K α radiation. Crystal data: $\text{C}_{12}\text{H}_{16}\text{O}_2$, $M_r = 192.26$, monoclinic, space group $P2_1/a$, $a = 9.020(6)$ Å, $b = 7.494(6)$ Å, $c = 15.673(3)$ Å, $\beta = 96.23(3)^\circ$, $V = 1053.1(8)$ Å³, $Z = 4$, $D_x = 1.213$ g/cm³, $F(000) = 416$ and $\mu(\text{Mo K}\alpha) = 0.756$ cm⁻¹. Of the 1521 independent reflections collected, 840 reflections with $I > 3.0\sigma(I)$ were used for the structure determination. The final refinement converged with $R = 0.058$ and $R_w = 0.072$ for 127 parameters. Atomic coordinates have been deposited at the Cambridge Crystallographic Data Center.
- X-ray analysis of ($\pm$)-**1**: Crystal size, 0.3 x 0.4 x 0.5 mm. All the data were obtained on Rigaku AFC-5S automated four circle diffractometer with graphite-monochromated Mo K α radiation. Crystal data: $\text{C}_{18}\text{H}_{20}\text{O}_5$, $M_r = 316.35$, monoclinic, space group $P2_1/n$, $a = 6.957(3)$ Å, $b = 17.541(3)$ Å, $c = 13.777(4)$ Å, $\beta = 97.78(3)^\circ$, $V = 1665.8(9)$ Å³, $Z = 4$, $D_x = 1.261$ g/cm³, $F(000) = 672$ and $\mu(\text{Mo K}\alpha) = 0.86$ cm⁻¹. Of the 2285 independent reflections collected, 1248 reflections with $I > 3.0\sigma(I)$ were used for the structure determination. The final refinement converged with $R = 0.056$ and $R_w = 0.063$ for 208 parameters. Atomic coordinates have been deposited at the Cambridge Crystallographic Data Center.
- Properties of ($\pm$)-**1**: m.p. 127-129 °C; IR (KBr) 2928, 2860, 1790, 1746, 1728, 1678, 1182, 1096, 1021, 957 cm⁻¹; ^1H NMR (500 MHz, C_6D_6) δ 0.94 (d, $J = 7.0$ Hz, 3H), 1.02 (m, 1H), 1.31 (m, 1H), 1.35 (t, $J = 1.5$ Hz, 3H), 1.43 (m, 1H), 1.53 (m, 1H), 1.60 (m, 1H), 1.70 (m, 1H), 2.27 (d, $J = 16.5$ Hz, 1H), 2.33 (br s, 1H), 2.41 (dd, $J = 16.5$ and 8.5 Hz, 1H), 3.19 (m, 1H), 5.09 (d, $J = 8.0$ Hz, 1H), 5.31 (t, $J = 1.5$ Hz, 1H), 5.81 (t, $J = 1.5$ Hz, 1H), 7.48 (d, $J = 2.5$ Hz, 1H); ^{13}C NMR (75.0 MHz, CDCl_3) δ 10.7, 18.5, 20.6, 26.0, 27.8, 31.1, 36.4, 41.3, 88.0, 100.4, 114.3, 135.8, 137.2, 141.0, 141.3, 150.1, 170.2, 171.3; MS (EI) m/z 316(M⁺), 219, 201, 173, 145, 131, 105, 97, 91.
- Hauck, C.; Schildeknecht, H. *J. Plant Physiol.* **1990**, *136*, 126-128.
- Mangnus, E. M.; Zwanenburg, B. *J. Agric. Food Chem.* **1992**, *40*, 697-700.
- Bergmann, C.; Wegmann, K.; Frischmuth, K.; Samson, E.; Kranz, A.; Weigelt, D.; Koll, P.; Welzel, P. *J. Plant Physiol.* **1993**, *142*, 338-342.
- Takeuchi, Y.; Omigawa, Y.; Ogasawara, M.; Yoneyama, K.; Konnai, M.; Worsham, A. D. *Plant Growth Regul.* **1995**, *16*, 153-160.

(Received in Japan 30 January 1997; revised 19 February 1997; accepted 21 February 1997)