

Toxic 12,13-Epoxytrichothec-9-enes from *Fusarium* sp.


By JOHN FREDERICK GROVE

(University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW)

Summary The structures and absolute configurations of a number of compounds obtained from a group of closely-related *Fusarium* sp. have been determined by correlation with 4 β ,15-diacetoxy-12,13-epoxy-trichothec-9-en-3 α -ol (diacetoxyscirpenol).

THE naturally-occurring esters of the group of sesquiterpene alcohols containing the tetracyclic 12,13-epoxytrichothec-9-ene nucleus¹ constitute an important class of mycotoxins.² The structure and absolute configuration of diacetoxyscirpenol (I; R¹ = Ac, R² = OH) have been established^{3,4} by relating it to verrucarol (I; R¹ = R² = H), whose absolute configuration is known.⁵ A number of these toxic compounds have now been related to diacetoxyscirpenol and their structures and absolute configurations thereby determined.

Structure (II; R¹ = OH, R² = R³ = Ac, R⁴ = H) was assigned,⁶ mainly on spectroscopic evidence, to the C₁₈H₂₄O₉



diacetate from *F. scirpi*;⁷ that this also represents the absolute configuration follows⁸ from the direct conversion

of the enone (II; $R^1 = H$, $R^2 = R^3 = R^4 = Ac$) into the diacetyl-derivative (II; $R^1 = OAc$, $R^2 = R^3 = R^4 = Ac$) of (II; $R^1 = OH$, $R^2 = R^3 = Ac$, $R^4 = H$) by acetoxylation with lead tetra-acetate. The enone (II; $R^1 = H$, $R^2 = R^3 = R^4 = Ac$) has been obtained from diacetoxyscirpenol by acetylation and oxidation with t-butyl chromate⁴ and from the acetylated 8α -(3-methylbutyryloxy)-derivative (III; $R^1 = Me_2CH\cdot CH_2\cdot CO$, $R^2 = R^5 = H$, $R^3 = R^4 = Ac$)^{9,10} by oxidation with selenium dioxide.⁹

Partial hydrolysis of the diacetate (II; $R^1 = OH$, $R^2 = R^3 = Ac$, $R^4 = H$) gave the 4β -monoacetate (II; $R^1 = OH$, $R^2 = R^4 = H$, $R^3 = Ac$), identical with fusarenone.¹¹ Hydrolysis of both ester groups furnished the parent tetraol (II; $R^1 = OH$, $R^2 = R^3 = R^4 = H$), identical with nivalenol.¹² The exceptional lability of the ester groups of (II; $R^1 = OH$, $R^2 = R^3 = Ac$, $R^4 = OH$), attributed to participation by the neighbouring 7α - and 3α -hydroxy-substituents, suggests that both fusarenone and nivalenol may be artifacts derived by hydrolysis of the diacetate during the lengthy isolation procedures.^{11,12}

Reduction of the 8-oxo-group of the diacetate (II; $R^1 = OH$, $R^2 = R^3 = Ac$, $R^4 = H$) with sodium borohydride took place from the less crowded β -face to give the $3\alpha,7\alpha,8\alpha$ -triol (III; $R^1 = R^5 = H$, $R^2 = OH$, $R^3 = R^4 = Ac$). The diacetyl derivative (III; $R^1 = R^3 = R^4 = R^5 = Ac$, $R^2 = OH$) was identical with a monoacetyl-derivative of the $C_{21}H_{28}O_{10}$ triacetate from *F. scirpi*⁷ which (n.m.r.) has structure (III; $R^1 = R^3 = R^4 = Ac$, $R^2 = OH$, $R^5 = H$).⁸

All the naturally-occurring biologically-active trichothecanes contain an ethylenic double bond at C-9 and a 12,13-epoxy-group. The elucidation of the structures of the toxins based on (II; $R^1 = OH$) and (III; $R^2 = OH$) illustrates the remarkable capacity of some *Fusaria* for oxidation of the trichothecane nucleus, which is mevalonate-derived.

This work was supported by a grant from the Agricultural Research Council during the tenure of a Comyns Berkeley Bye-Fellowship from Gonville and Caius College.

(Received, August 5th, 1969; Com. 1199.)

¹ W. O. Godtfredsen, J. F. Grove, and Ch. Tamm, *Helv. Chim. Acta*, 1967, **50**, 1666.

² J. F. Grove and P. H. Mortimer, *Biochem. Pharmacol.*, 1969, **18**, 1473.

³ A. W. Dawkins, J. F. Grove, and B. K. Tidd, *Chem. Comm.*, 1965, 27; A. W. Dawkins, *J. Chem. Soc. (C)*, 1966, 116.

⁴ E. Flury, R. Mauli, and H. P. Sigg, *Chem. Comm.*, 1965, 26; H. P. Sigg, R. Mauli, E. Flury, and D. Hausen, *Helv. Chim. Acta*, 1965, **48**, 962.

⁵ A. T. McPhail and G. A. Sim, *J. Chem. Soc. (C)*, 1966, 1394.

⁶ B. K. Tidd, *J. Chem. Soc. (C)*, 1967, 218.

⁷ P. W. Brian, A. W. Dawkins, J. F. Grove, H. G. Hemming, D. Lowe, and G. L. F. Norris, *J. Exp. Bot.*, 1961, **12**, 1.

⁸ J. F. Grove, *J. Chem. Soc. (C)*, in the press.

⁹ J. R. Bamburg, N. V. Riggs, and F. M. Strong, *Tetrahedron*, 1968, **24**, 3329.

¹⁰ S. G. Yates, H. L. Tookey, J. J. Ellis and H. J. Burkhardt, *Phytochem.*, 1968, **7**, 139.

¹¹ H. Tsunoda, N. Toyazaki, N. Marooka, N. Nakano, H. Yoshiyama, K. Okubo, and M. Isoda, *Rep. Fd. Res. Inst. (Min. Agr. Forest., Japan)*, 1968, **23**, 89.

¹² T. Tatsuno, M. Saito, M. Enomoto, and H. Tsunoda, *Chem. Pharm. Bull.*, 1968, **16**, 2519.