

and **4** extruded CF_2 ; at almost equal rates ($k_2/k_3 = 1.05$).

Our observed activation energy for the geometrical isomerization of **2** (49.7 kcal/mol) can be compared with that for the analogous nonfluorinated molecule (59.1 kcal/mol) which was determined by Flowers and Frey in 1960.¹⁵ It can thus be seen that the presence of the CF_2 group significantly facilitates the isomerization process. However it must be emphasized that precise interpretation of these results at this time is premature. There is, as of yet, no experimental evidence as to the mechanism of the cis-trans interconversion, i.e., whether cleavage of bond $\text{C}_1\text{--C}_2$ or of bond $\text{C}_2\text{--C}_3$ is responsible for the observed isomerization. The earlier mentioned exo-endo epimerizations^{10,11} do, of course, provide strong credibility for $\text{C}_2\text{--C}_3$ cleavage, but work is currently in progress in our laboratories to fully elucidate the mechanism of the isomerization process.

Acknowledgment. Grateful acknowledgement is given to the National Science Foundation for partial support of this research and to Mr. David Muthard for assistance in the preparation of the cyclopropanes.

References and Notes

- (1) (a) R. Hoffmann and W. P. Stohrer, *J. Am. Chem. Soc.*, **93**, 6941 (1971); (b) R. Hoffmann, *Tetrahedron Lett.*, 2907 (1970).
- (2) H. Günter, *Tetrahedron Lett.*, 5173 (1970).
- (3) J. M. Simpson and H. G. Richey, Jr., *Tetrahedron Lett.*, 2545 (1973).
- (4) (a) W. Kirmse and H. R. Murawski, *J. Chem. Soc., Chem. Commun.*, 122 (1977); (b) W. Kirmse and M. Zeppenfeld, *ibid.*, 124 (1977).
- (5) R. J. Pearson, Jr., A. Choplin, and V. W. Laurie, *J. Chem. Phys.*, **62**, 4859 (1975).
- (6) A. R. Mochel, C. O. Britt, and J. E. Boggs, *J. Chem. Phys.*, **58**, 3221 (1973).
- (7) A. T. Perretta and V. W. Laurie, *J. Chem. Phys.*, **62**, 2469 (1975).
- (8) A. Skancke, E. Flood, and J. E. Boggs, unpublished results.
- (9) E. Vogel, unpublished results, as reported in ref 2.
- (10) P. B. Sargeant, *J. Am. Chem. Soc.*, **91**, 3061 (1969).
- (11) C. W. Jefford, J. Mareda, J. C. E. Gehret, T. Karbengele, W. D. Graham, and U. Burger, *J. Am. Chem. Soc.*, **98**, 2585 (1976).
- (12) Very little kinetic data is presently available for perfluoro or partially fluorinated cyclopropane thermal processes: (a) F. Cassas, J. A. Kerr, and A. F. Trotman-Dickenson, *J. Chem. Soc.*, 3655 (1964); (b) F. F. Herbert, J. A. Kerr, and A. F. Trotman-Dickenson, *ibid.*, 5710 (1965); (c) R. A. Mitsch and F. W. Neuvar, *J. Phys. Chem.*, **70**, 546 (1966).
- (13) (a) D. J. Burton and D. G. Naee, *J. Am. Chem. Soc.*, **95**, 8467 (1973); (b) D. J. Burton and D. G. Naee, *Syn. Commun.*, **3**, 197 (1973).
- (14) Surface catalyzed decomposition of the reaction mixture was difficult to eradicate. Conditioning of the Pyrex vessel with starting materials at temperatures $>300^\circ\text{C}$ for a period of 3–4 weeks created a vessel which allowed good first-order isomerization rates to be measured.
- (15) M. C. Flowers and H. M. Frey, *Proc. R. Soc. London, Ser. A*, **257**, 122 (1960); M. C. Flowers and H. M. Frey, *ibid.*, **260**, 424 (1961).
- (16) J. M. Birchall, R. N. Haszeldine, and D. W. Roberts, *J. Chem. Soc., Chem. Commun.*, 287 (1967).
- (17) B. Atkinson and D. McKeagan, *J. Chem. Soc., Chem. Commun.*, 189 (1966).
- (18) P. B. Sargeant and C. G. Krespan, *J. Am. Chem. Soc.*, **91**, 415 (1969).

William R. Dolbier, Jr.,* Harold O. Enoch

Department of Chemistry, University of Florida
Gainesville, Florida 32611


Received April 8, 1977

New Synthetic Strategy for the Preparation of Linear Phenolic Natural Products


Sir:

We have developed a conceptually simple synthetic strategy which significantly facilitates the preparation of linear polynuclear phenolic systems, a structural feature present in many antibiotics and other natural products. This strategy, shown in Scheme I, involves repetitive execution¹ of a single regioselective reaction sequence which adds a new phenolic ring to the existing aromatic system in each cycle. To illustrate the versatility of this synthetic strategy, 2-methyl-6-methoxybenzoic acid (**1**) was carried through two cycles yielding the

Scheme I



Scheme II



polyfunctionalized anthracenecarboxylic ester **8** (Scheme II).

The aglycone portion of numerous naturally occurring compounds consists of linear tri- and tetracyclic systems with two or more aromatic rings.² Many of the parent compounds, such as the anthracyclines adriamycin,³ daunorubicin,⁴ and carminomycin,⁵ olivomycins A, B, C, and D,⁶ chromomycins A 1–4,^{6,7} and aureolic acid,⁸ have, in addition to their antibiotic activity, significant anticancer properties. Of these, only daunorubicin^{9,10} has been synthesized.

Generally, only three approaches, the use of the Friedel-Crafts acylation,^{9,11} variant forms of the Diels-Alder reaction,^{12–14} or cyclization of polyketides,^{15,16} have been employed to construct linear polynuclear systems. Alternate synthetic methodology is much needed to permit the regioselective preparation of linear phenolic systems not achievable by these other approaches. The described repetitive synthetic strategy and the derived reaction sequence which accomplishes it should be adaptable to a broader range of these systems.

The starting material chosen for the synthesis, 2-methyl-6-methoxybenzoic acid¹⁶ (**1**), was carboxylated on the methyl group by quenching the orange dilithium anion, formed by treatment of **1** with lithium diisopropylamide at -78°C with dimethyl carbonate. Aqueous workup gave directly homophthalic acid **2**, mp $165\text{--}166^\circ\text{C}$ (lit.¹⁷ mp $165\text{--}166^\circ\text{C}$), in 90% yield.^{18,19} A three-step sequence involving acylation of **2** with acetic anhydride and pyridine, followed by basic hydrolysis with concomitant decarboxylation, and, finally, ring closure of the intermediate using acetic anhydride and perchloric acid yielded isocoumarin (**3**), mp 109°C (lit.¹⁷ mp $109.5\text{--}110.5^\circ\text{C}$), in 68% overall yield.^{20,21} Slow addition of a dilute benzene solution of ethyl bromoacetate to a refluxing benzene solution of **3** and zinc gave naphthoate **4**, mp 59°C ,

in 73% yield.²² Methylation of **4** was accomplished using dimethyl sulfate and potassium carbonate; subsequent hydrolysis of the ethyl ester with potassium hydroxide in dimethyl sulfide²³ gave naphthoic acid (**5**), mp 149–50 °C.

The ortho disposed carboxyl and methyl functionalities used in the conversion of benzoate system **2** to naphthoate system **5** are retained and permit the annelation sequence to be repeated. Naphthoic acid (**5**), upon treatment with lithium diisopropylamide, was converted to a deep purple dilithium anion which was carboxylated and hydrolyzed to give naphthalene-acetic acid (**6**).²⁴ The reaction sequence described earlier for conversion of homophthalic acid (**2**) to isocoumarin (**3**) was employed again for the preparation of naphtho[2,3-*c*]pyran (**7**), mp 168–70 °C, in 56% overall yield from naphthoic acid (**5**). Reformatsky reaction on naphtho[2,3-*c*]pyran (**7**), gave 9,10-dimethoxy-1-hydroxy-3-methylanthracenecarboxylic acid (**8**), mp 159–161 °C, which bears a hydroxylation pattern found in naturally occurring linear polynuclear aromatic systems.

Studies to broaden the scope of this approach for synthesis of hydroxylated polynuclear aromatic compounds and to synthesize selected natural products are in progress.

Acknowledgment. The authors wish to thank the National Cancer Institute of DHEW, Grant No. CA 18141, for support of this work.

References and Notes

- (1) The use of repetitive reaction sequences as a synthetic strategy to construct complex molecules is not totally new, as it has been employed extensively, even automated, in polypeptide synthesis (G. R. Marshall and R. B. Merrifield, "Biochemical Aspects of Reactions on Solid Supports", G. Stark, Ed., Academic Press, New York, N.Y., 1971, Chapter 3). Another application of this strategy is the synthesis of naturally occurring 1,5,10,*n*-polyene containing systems by repetitive Claisen–Cope rearrangement (S. J. Rhoads and N. R. Raulins, *Org. React.*, **22**, 1 (1975)). Recently an elegant synthesis of Semicorin E was reported which employed repetitive addition of a nitrile oxide to an acetylene (R. V. Stevens and E. B. Reid, *Tetrahedron Lett.*, 4193 (1975)).
- (2) For an extensive list of these compounds see "Naturally Occurring Quinones", R. H. Thompson, Academic Press, New York, N.Y., 1971, and "Handbook of Naturally Occurring Compounds", Vol. I, T. K. Devon and A. I. Scott, Ed., Academic Press, New York, N.Y., Chapters 25–27.
- (3) F. Arcamone, G. Franceschi, and S. Penco, *Tetrahedron Lett.*, 1007 (1969).
- (4) F. Arcamone, G. Franceschi, P. Orezzi, G. Cassinelli, W. Barbiere, and R. Mondelli, *J. Am. Chem. Soc.*, **86**, 5334 (1964).
- (5) M. G. Brazhnikova, V. B. Zbarsky, V. I. Ponomareko, and N. P. Potapova, *J. Antibiot.*, **27**, 254 (1974).
- (6) Yu. A. Berlin, S. E. Esipor, M. N. Kolosov, and M. M. Shemyakin, *Tetrahedron Lett.*, 1643–1647 (1966).
- (7) Yu. A. Berlin, O. A. Kiseleva, M. N. Kolosov, M. M. Shemyakin, V. B. Soifer, I. V. Vasina, I. V. Yartseva, and V. D. Kuznetov, *Nature*, **218**, 193 (1968).
- (8) Yu. A. Berlin, M. N. Kolosov, and L. A. Piotrovich, *Tetrahedron Lett.*, 1329 (1970).
- (9) C. M. Wong, R. Schwenk, D. Popien, and T.-L. Ho, *Can. J. Chem.*, **51**, 466 (1973).
- (10) A. S. Kende, Y.-g. Tsay, and J. E. Mills, *J. Am. Chem. Soc.*, **98**, 1967 (1976).
- (11) For a use of this reaction in an approach to the synthesis of adriamycin, see D. G. Miller, S. Trenbeath, and C. J. Sih, *Tetrahedron Lett.*, 1637 (1976), and R. D. Gleim, S. Trenbeath, R. S. D. Mittal, and C. J. Sih, *ibid.*, 3385 (1976).
- (12) L. W. Butz and A. W. Rytina, *Org. React.*, **5**, 136 (1949).
- (13) P. G. Sammes and T. W. Wallace, *J. Chem. Soc., Chem. Commun.*, 524 (1973).
- (14) P. G. Sammes, *Tetrahedron*, **32**, 405 (1976).
- (15) For reviews, see (a) T. Money, *Chem. Rev.*, **70**, 553 (1970), and T. M. Harris, C. M. Harris, and K. B. Hindley, *Fortschr. Chem. Org. Naturst.*, **31**, 219 (1974).
- (16) T. M. Harris and P. J. Wittek, *J. Am. Chem. Soc.*, **97**, 3270 (1975).
- (17) M. Matsui, K. Mori, and S. Arasaki, *Agr. Biol. Chem.*, **28**, 896 (1964).
- (18) F. Hauser and R. Rhee, *Synthesis*, 245 (1977). Coaddition of **1** and dimethylcarbonate to lithium diisopropylamide at –78 °C gives an equally good yield.
- (19) Satisfactory elemental analyses were obtained for previously unreported compounds. Mass and ¹H NMR spectra were, in each instance, in accord with the assigned structures.
- (20) H. L. Slaters, S. Weber, and N. L. Wendler, *Chimia*, **21**, 468 (1967).
- (21) Attempts to abbreviate the sequence by direct acylation of the dilithium anion with acetic anhydride, ethyl acetate, acetyl chloride, or phenyl acetate failed.
- (22) This is a modification of a reaction previously reported by M. Pailer and O. Vostrowsky, *Monatsh. Chem.*, **102**, 951 (1971).

- (23) The ortho-disubstituted ester was virtually resistant to hydrolysis in ethanolic sodium or potassium hydroxide.
- (24) Compound **6** could not be purified; however, comparison of the relative intensity of the ¹H NMR for the aromatic methyl group for the starting material (δ 2.42 ppm) with that for the methylene group of the phenyl acetic acid product (singlet at δ 3.81 ppm) indicated that the conversion of **5** to **6** had occurred in 85–90% yield.

Frank M. Hauser,* Richard Rhee

Department of Chemistry, Oregon Graduate Center
Beaverton, Oregon 97005

Received March 21, 1977

A Secondary Isotope Effect in the Cysteine-Promoted Dehalogenation of 5-Bromo-2'-deoxyuridine. Evidence for Transient 5,6-Dihydropyrimidine Intermediates

Sir:

There has been long-standing interest in 5-halogenated pyrimidines because of their use as biochemical tools and chemotherapeutic agents.^{1,2} The metabolism of 5-bromo- and 5-iodouracils involves their dehalogenation, and numerous model systems have been investigated in attempts to understand the mechanism of these reactions.^{3–12} Most studies have centered about the bisulfite mediated halide release and its replacement by proton; the same reaction is prompted by thiols, which is more relevant to enzymic conversions, but less well understood. The proposed mechanism for the thiol mediated dehalogenation of 1-substituted 5-bromouracils is depicted in Scheme I; a similar pathway is believed to exist for dehalogenation of corresponding 5-iodouracils.¹⁰ The initial, and perhaps rate-determining,^{9,10} step is believed to involve attack of thiolate at the 6 position of the heterocycle **1** and protonation of C-5 to produce the 5-bromo-6-thiol-5,6-dihydrouracil **2**. Two general pathways have been proposed^{9,10,12} to account for subsequent steps leading to the dehalogenated product. The first, E2 Hal, involves abstraction of bromonium (Br⁺) ion from **2** to provide intermediate **3** and a sulfonyl halide. The latter would react with a thiol to provide the halide ion and




Figure 1. Secondary isotope effect on the cysteine-promoted dehalogenation of BrdUrd. Experimental points refer to the ³H/¹⁴C ratio of BrdUrd (●), dUrd (○), and (△) 5-CysdUrd. The lines are theoretical¹⁵ for a secondary isotope effect of $k_T/k_H = 1.18$.