

Communications to the Editor

[Chem. Pharm. Bull.]
[24(2) 360-362 (1976)]

UDC 547.913.4.02 : 581.192

New Eremophilane-Type Lactones from *Ligularia Fauriei* (Fr.) Koidz.

The structures of three new sesquiterpenes isolated from *Ligularia Fauriei* (Fr.) Koidz. were shown to be 1, 2, and 3, respectively.


In connection with structural studies on constituents of plants of the genus *Ligularia* (Compositae) and related plants,¹⁾ three new sesquiterpenes have been isolated from the benzene extract of *Ligularia Fauriei* (Fr.) Koidz. We wish to describe the structure determination leading to 1, 2, and 3 for these substances.

The molecular formula of $C_{15}H_{22}O_4$ ^{2,3)} was given for a compound (1), mp 217–218° (dec.), $[\alpha]_D +82^\circ$ ($c=0.98$, EtOH). The presence of an α,β -unsaturated γ -lactone moiety^{4,5)} and hydroxyl group(s) was suggested from the infrared (IR) ($\nu_{\max}^{\text{Nujol}}$ 3480, 3230, and 1721 cm^{-1}) and the ultraviolet (UV) ($\lambda_{\max}^{\text{EtOH}}$ 219 nm, ϵ 11800) spectra. The proton magnetic resonance (PMR) spectrum (acetone- d_6) showed the presence of a secondary methyl (δ 0.80, d -like; $C_{(4)}\text{-CH}_3$), a tertiary methyl (δ 1.12, s, $C_{(5)}\text{-CH}_3$), an olefinic methyl (δ 1.83, s, $C_{(11)}\text{-CH}_3$), and a proton (δ 4.64, br. s, $C_{(6a)}\text{-H}$) attached to a carbon bearing an oxygen atom. Acetylation of 1 with acetic anhydride in pyridine yielded a diacetate (4), mp 120.5–121°, $C_{19}H_{26}O_6$ ²⁾ which gave no hydroxyl absorption in its IR spectrum. In the PMR spectrum (CDCl_3) of 4 two methyl signals newly appeared (δ 1.98, s, and δ 2.02, s, each $\text{CH}_3\text{CO-O-}$). Therefore two hydroxyl groups must be present in 1.

Reduction of 1 with sodium borohydride in methanol gave two products (5 and 6). The more polar product (5) [mp 207–207.5°, $[\alpha]_D +213^\circ$ ($c=1.19$, CHCl_3), $C_{15}H_{22}O_3$ (M^+ at m/e 250); $\nu_{\max}^{\text{Nujol}}$ 3440, 1739, 1709(sh), and 1690(sh) cm^{-1} ; $\lambda_{\max}^{\text{EtOH}}$ 218 nm, ϵ 14400; PMR (CDCl_3): δ 0.80 (d -like; $C_{(4)}\text{-CH}_3$), δ 1.13 (s, $C_{(5)}\text{-CH}_3$), δ 1.85 (d, $J_{8\beta,13}=2$ Hz; $C_{(11)}\text{-CH}_3$), δ 4.80 (s, $C_{(6a)}\text{-H}$), δ ca. 5.1 (m, $C_{(8\beta)}\text{-H}$), and δ 2.23 (s, OH, disappeared on addition of D_2O); (acetone- d_6): δ 0.78 (d -like; $C_{(4)}\text{-CH}_3$), δ 1.10 (s, $C_{(5)}\text{-CH}_3$), δ 1.77 (d, $J_{8\beta,13}=2$ Hz; $C_{(11)}\text{-CH}_3$), δ 4.70 (s, $C_{(6a)}\text{-H}$), and δ ca. 5.1 (m, $C_{(8\beta)}\text{-H}$)] proved to be identical (mp, mixed mp, $[\alpha]_D$, IR, UV, PMR, and mass spectra) with 6 β -hydroxyeremophil-7(11)-en-12,8 α -olide (5).⁵⁻⁷⁾ The less polar product (6) [mp 107–111°, $C_{15}H_{22}O_2$ (M^+ at m/e 234); $\nu_{\max}^{\text{Nujol}}$ 1734, 1670, and 1025 cm^{-1} ; PMR (CDCl_3): δ 0.80 (d -like; $C_{(4)}\text{-CH}_3$), δ 1.05 (s, $C_{(5)}\text{-CH}_3$), δ 1.80 (d, $J_{8\beta,13}=2$ Hz; $C_{(11)}\text{-CH}_3$), δ ca. 2.9 (d, $J=15$ Hz; an A-part of an AB-type quartet due to the allylic methylene protons at $C_{(6)}$),⁸⁾ and δ 4.61 (m, $C_{(8\beta)}\text{-H}$)] was found to be identical with eremophil-7(11)-en-12,8 α -olide (6).^{7,9)} Similar reduction of 5 with sodium borohydride gave 6. The structure of 1 must contain one hydroxyl group more than that of 5.

In the PMR spectrum of 5 a multiplet due to a proton at $C_{(8\beta)}$ appeared at δ ca. 5.1, while the corresponding signals were absent in that of 1. The presence in 5 of long-range spin-

- 1) T. Sato, Y. Moriyama, H. Nagano, Y. Tanahashi, and T. Takahashi, *Bull. Chem. Soc. Japan*, **48**, 112 (1975). And the references cited therein.
- 2) Elemental analysis of the compound gave a satisfactory result.
- 3) The mass spectrum of 1 showed the $M-18$ ion peak at m/e 248, while no molecular ion peak was observed.
- 4) K. Takeda, H. Minato, M. Ishikawa, and M. Miyawaki, *Tetrahedron*, **20**, 2655 (1964).
- 5) H. Ishii, T. Tozyo, and H. Minato, *J. Chem. Soc. (C)*, **1966**, 1545.
- 6) L. Novotný, V. Herout, and F. Šorm, *Coll. Czech. Chem. Comm.*, **29**, 2189 (1964).
- 7) The $C_{(8\beta)}\text{-H}$ configuration of 5 was determined by X-ray study: C. Kabuto, N. Takada, S. Maeda, and Y. Kitahara, *Chem. Lett.*, **1973**, 371.
- 8) The B-part signals were overlapped with other signals.
- 9) L. Novotný, J. Jizba, V. Herout, and F. Šorm, *Coll. Czech. Chem. Comm.*, **27**, 1393 (1962); M. Horák, O. Motl, J. Pliva, and F. Šorm, "Die Terpene," Teil II, S I 28, Akademie-Verlag, Berlin, 1963.



coupling (2 Hz) of the $C_{(8\beta)}$ -H to the olefinic $C_{(11)}$ -CH₃ protons (δ 1.85) was shown by proton magnetic double resonance (PMDR) experiments. The other spectral data of **1** and **5** were almost identical. These observations led to the location of an extra hydroxyl group on $C_{(8\beta)}$ for **1**.¹⁰⁾ The structure of **1** should be represented by 6 β , 8 β -dihydroxyeremophil-7(11)-en-12,8 α -olide (**1**).

The compound (**2**), mp 253.5–254°, $[\alpha]_D +94^\circ$ ($c=0.88$, CHCl₃), C₁₅H₁₈O₅²⁾ (M^+ at m/e 278), showed the IR ($\nu_{\max}^{\text{Nujol}}$ 3260, 1787, and 1710 cm⁻¹) and the UV ($\lambda_{\max}^{\text{EtOH}}$ 214 nm, ϵ 12400) spectra suggesting the presence of an α,β -unsaturated γ -lactone and a γ -lactone moiety along with a hydroxyl group. The PMR spectrum (acetone-*d*₆) indicated the presence of a tertiary methyl (δ 1.28, s, $C_{(5)}$ -CH₃), an olefinic methyl (δ 1.86, d, $J_{6\beta,13}=2$ Hz; $C_{(11)}$ -CH₃), a proton (δ 5.18, q, $J_{6\beta,13}=2$ Hz; $C_{(6\beta)}$ -H), and a hydroxyl group (δ 6.51, s, disappeared on addition of D₂O); signals due to a secondary methyl were absent. On acetylation with acetic anhydride in pyridine **2** gave a monoacetate (**7**), mp 183–184°, C₁₇H₂₀O₆²⁾ (M^+ at m/e 320), which showed no hydroxyl absorption in its IR spectrum.

These spectral data could be best interpreted on the basis of the structure (**2**), providing that **2** belongs to a sesquiterpene of eremophilane-type. This received support from the following evidences. (i) Treatment of furanoeremophilan-14 β ,6 α -olide (**8**)¹¹⁾ in dioxane with dicyanodichlorobenzoquinone (DDQ)¹²⁾ at 80° for 1 hr yielded an unstable unsaturated enol lactone (**9**) [an amorphous compound, C₁₅H₁₆O₄ (M^+ at m/e 260); $\nu_{\max}^{\text{neat}}$ 1785 and 1665 cm⁻¹, $\lambda_{\max}^{\text{EtOH}}$ 280 nm;¹³⁾ PMR (CDCl₃): δ 1.38 (s, $C_{(5)}$ -CH₃), δ 2.12 (br. s, $C_{(11)}$ -CH₃), δ 2.65 (dd, $J=2$ and 12 Hz; $C_{(4\alpha)}$ -H), δ 2.96 (m, $C_{(10\beta)}$ -H), δ 5.28 (q, $J_{6\beta,13}=1.5$ Hz; $C_{(6\beta)}$ -H), and δ 5.65 (d, $J_{9,10\beta}=3$ Hz; $C_{(9)}$ -H)]. The same compound (**9**) was obtained by dehydration of **2** with phosphorus oxychloride in pyridine (identification: IR, PMR, and mass spectra). (ii) The lactone (**8**) in methanol was irradiated with a high pressure mercury lamp under an oxygen atmosphere for 1 hr in the presence of Rose Bengal to afford a compound identical (mp, mixed mp, IR and PMR spectra) with **2**. These observations led to the structure of 8 β -hydroxyeremophil-7(11)-ene-12, 8 α ; 14 β , 6 α -diolide (**2**)^{14,15)} for **2**.

The presence of an α,β -unsaturated γ -lactone and a γ -lactone grouping was suggested for the compound (**3**) [mp 186–186.5° (decomp.), $[\alpha]_D +93^\circ$ ($c=0.91$, CHCl₃), C₁₅H₁₈O₄²⁾ (M^+ at m/e 262); $\nu_{\max}^{\text{Nujol}}$ 1786, 1739, and 1684 cm⁻¹; $\lambda_{\max}^{\text{EtOH}}$ 217 nm (ϵ 19300); PMR (CDCl₃): δ 1.29 (s, $C_{(5)}$ -CH₃), δ 2.00 (t, $J_{6\beta,13}=2$ Hz; $J_{8\beta,13}=2$ Hz; $C_{(11)}$ -CH₃), δ 4.75 (m, $C_{(8\beta)}$ -H), δ 5.05 (br. s,

10) An alternative structure with 8 α -OH configuration would show a different PMR spectrum from that of **5** (with 8 β -H configuration).

11) Y. Ishizaki, Y. Tanahashi, T. Takahashi, and K. Tori, *Chem. Comm.*, **1969**, 551.

12) Oxidation of atractylon and linderene with DDQ has been described: K. Takeda, M. Ikuta, M. Miyawaki, and K. Tori, *Tetrahedron*, **22**, 1159 (1966).

13) The ϵ value could not be determined precisely due to fragility of the compound.

14) Closely related PMR spectral data for **1**, **2**, **3**, **5**, and **6**, suggested that the configurations at $C_{(8)}$ of these compounds are the same ($C_{(8\beta)}$ -H or $C_{(8\beta)}$ -OH). PMDR experiments showed the presence of long-range spin-couplings of the olefinic $C_{(11)}$ -CH₃ protons to the $C_{(6\beta)}$ -H (at δ 5.18 for **2** and at δ 5.05 for **3**) and to the $C_{(8\beta)}$ -H (at δ 4.75 for **3**).

15) The lactones (**2** and **1**) might be considered as artifacts produced, during isolation, from **8** and 6 β -hydroxy-furanoeremophilane, respectively. An examination on this point is under way.

$C_{(6\beta)}-H]$. Reduction of **2** with sodium borohydride in methanol yielded **3**. The structure of **3** must be shown as eremophil-7(11)-ene-12,8 α ; 14 β ,6 α -diolide (**3**),¹⁴ since **1** was reduced with sodium borohydride to form **5** with a loss of the hydroxyl group at $C_{(8\beta)}$. The spectral data described above are compatible with the structure (**3**).

Finally, 6 β -hydroxyeremophil-7(11)-en-12,8 α -olide (**5**)⁵⁻⁷ was also isolated from the same plant.

Compounds **2** and **3** constitute the first examples of eremophilane-type sesquiterpenes having two lactone rings in their molecules.

Acknowledgement We wish to thank Dr. H. Ishii, Shionogi Research Laboratory, Osaka, for a generous gift of an authentic sample of 6 β -hydroxyeremophil-7(11)-en-12,8 α -olide (**5**).

Department of Chemistry,
Faculty of Science,
The University of Tokyo,
Bunkyo-ku, Tokyo

YOSHIHIKO MORIYAMA
TAKEYOSHI TAKAHASHI

Received September 13, 1975

[Chem. Pharm. Bull.]
[24(2) 362-365 (1976)]

UDC 547.869.1.04 : 547.789.1.04

Opening of the Cephalosporin Dihydrothiazine Ring

2-Ethoxy-3-cephem-1 β -oxide (**3**) was found to be thermally unstable and easily converted into isothiazolones (**4**, **5**, **6**, **9**, **10**), and the β -lactam derivative (**8**) under varying reaction conditions. Furthermore, 2-ethoxy-3-cephem (**2**) was treated with *tert*-butyl hypochlorite, giving the azetidinone-oxazoline acetate (**14**).

In a previous paper,¹⁾ we described a new rearrangement reaction of 2-methylthio- or 2-methoxycephalosporins into azlactone derivatives similar to the penicillin-penicillenate rearrangement reaction.²⁾ This reaction suggested that introduction of a heteroatom substituent at C(2) of cephem molecules would facilitate ring opening of the dihydrothiazine moiety. Further, we wish to add herein other transformation reactions of 2-alkoxycephems involving S(1)-C(2) bond fission.

Treatment of methyl 7 β -benzamido-3-methyl-3-cephem-4-carboxylate³⁾ (**1**) with 1.2 equivalents of *tert*-butyl hypochlorite in ethanol-containing methylene chloride (0°, 1 day, 45% yield) gave a 2 α -ethoxy-3-cephem (**2**, mp 178—179.5°⁴⁾).⁵⁾ Successive oxidation of **2** with 1 equivalent of *meta*-chloroperbenzoic acid in chloroform (0°, 1 hr, 73% yield) afforded a 2 α -ethoxy-1 β -oxide [**3**, mp 137—138°; IR $\nu_{\max}^{\text{Nujol}}$ cm⁻¹: 3320, 1785, 1736, 1660, 1543; NMR (CDCl₃) δ ppm: 1.20 (3H, t, $J=7$, -OCH₂CH₃), 2.12 (3H, s, 3-CH₃), 3.80 (3H, s, -COOCH₃), 4.60 (1H, d, $J=5.5$, H-6), 4.63 (1H, s, H-2), 6.25 (1H, dd, $J=5.5$ and 10, H-7)]. The 2 α -ethoxy-1-oxide **3** thereby obtained was found to be thermally unstable in protic solvents; and **3** was easily converted into an isothiazolone diethylacetal [**4**, mp 142—143°; IR $\nu_{\max}^{\text{CHCl}_3}$ cm⁻¹: 3430, 1734, 1660, 1524;

1) A. Yoshida, S. Oida, and E. Ohki, *Chem. Pharm. Bull.* (Tokyo), **23**, 2518 (1975).

2) H. Bundgaard, *J. Pharm. Sci.*, **60**, 1273 (1971) and the related references cited therein.

3) A. Yoshida, S. Oida, and E. Ohki, *Chem. Pharm. Bull.* (Tokyo), **23**, 2507 (1975).

4) All compounds were characterized by infrared (IR), nuclear magnetic resonance (NMR) and mass spectrometry and also by elementary analysis.

5) cf. D.O. Spry, *Tetrahedron Letters*, **1972**, 3717.