View Article Online View Journal

# ChemComm

# Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: K. Sato, S. Tanaka, K. Yamamoto, Y. Tashiro, T. Narumi and N. Mase, *Chem. Commun.*, 2018, DOI: 10.1039/C8CC03591A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemcomm

# **Journal Name**



# COMMUNICATION

# Direct synthesis of N-terminal thiazolidine-containing peptide thioesters from peptide hydrazides<sup>†</sup>

Received 00th January 20xx, Accepted 00th January 20xx Kohei Sato, \*a Shoko Tanaka, a Kazuki Yamamoto, Yosuke Tashiro, ab Tetsuo Narumi abc and Nobuyuki Mase abc

DOI: 10.1039/x0xx00000x

www.rsc.org/

Published on 05 June 2018. Downloaded by Hacettepe Universitesi on 05/06/2018 06:32:14.

We report a simple and promising synthetic method to oxidize peptide hydrazides containing N-terminal thiazolidine as a protected cysteine. This yields the corresponding thioester *via* a peptide azide without decomposition of the thiazolidine ring. The newly developed protocol was validated by the synthesis of the bioactive peptide LacZ $\alpha$ .

Chemical synthesis of proteins has attracted considerable attention as an important tool in the field of structural biology,<sup>1</sup> drug discovery<sup>2</sup> and chemical biology.<sup>3</sup> Although the total- or semi-synthesis of a protein remains challenging, recent advances, including chemoselective ligations and use of related protecting groups are bringing the chemical synthesis of proteins into the mainstream in synthetic chemistry. In particular, native chemical ligation (NCL), developed by Kent et al.<sup>4,5</sup> is a powerful technique for the chemoselective ligation between a peptide thioester and an N-terminal cysteinyl peptide, yielding the ligated product with high efficiency. A key to the success of the chemical synthesis of a large protein is how more than two peptide segments can be assembled in a sequential and convergent manner.<sup>6</sup> In sequential NCL strategy, the N-terminal Cys residue of the middle segment must be protected in order to prevent cyclization caused by intramolecular reactions with the thioester moiety (Figure 1). Among the protecting groups that have been developed,<sup>7</sup> the 1,3-thiazolidine-4-carboxo (Thz) group is a reliable choice because of its facile convertibility to Cys and its commercial availability, and it has been applied extensively to the synthesis of various proteins.6,7c





Fig. 1 (A) Conversion of peptide hydrazide bearing protected cysteine to the corresponding thioester. (B) Direct conversion of Thz-peptide hydrazide to the corresponding thioester using TFA as a solvent and subsequent sequential NCLs. PG: protecting groups; Tbeoc: 2-(*tert*-butyldisulfanyl)ethyloxycarbonyl; Dobz: *p*-dihydroxyborylbenzyloxycarbonyl; Tfacm: trifluoroacetamidomethyl; Tfa: trifluoroacetyl.

<sup>&</sup>lt;sup>a.</sup> Department of Engineering, Graduate School of Integrated Science and Technology, Shizuoka University, 3-5-1 Johoku, Hamamatsu, Shizuoka 432-8561, Japan. Email: sato.kohei@shizuoka.ac.jp

<sup>&</sup>lt;sup>b.</sup> Graduate School of Science and Technology, Shizuoka University, 3-5-1 Johoku, Hamamatsu, Shizuoka 432-8561, Japan

<sup>&</sup>lt;sup>c</sup> Research Institute of Green Science and Technology, Shizuoka University, 3-5-1 Johoku, Hamamatsu, Shizuoka 432-8561, Japan

<sup>&</sup>lt;sup>+</sup> Electronic Supplementary Information (ESI) available: Experimental details of syntheses including HPLC charts. See DOI: 10.1039/x0xx00000x

#### COMMUNICATION

prepared by conventional 9-fluorenylmethyloxycarbonyl (Fmoc) solid-phase peptide synthesis (SPPS) on hydrazineincorporated resin and can be converted into a thioester by treatment with NaNO<sub>2</sub> under acid conditions, followed by thiolysis of the resulting peptidyl azide (Figure 1A). This technique has seen widespread use and the number of ligations carried out using hydrazides as an acyl donor represents 30% of the total ligations reported in 2016.9 This method however has not been used widely in the synthesis of N-terminal Thz-peptide thioesters due to the decomposition of the Thz moiety in the hydrazide in the acidic activation step.<sup>10a</sup> To circumvent this problem, alternative protections for Cys have been examined. The functionalized Cys derivatives that are called for by these methods must be prepared (Figure 1A).<sup>10</sup> Thus, there is an increasing demand for the development of practical methods for the direct synthesis of Thz-containing peptide thioesters from peptide hydrazides (Figure 1B).

Peptides with a Thz group are stable during cleavage reactions mediated by HF or TFA, and the moiety can be converted to the corresponding Cys residue under acidic aqueous conditions with MeONH<sub>2</sub>.<sup>7c</sup> Based on these facts, we speculated that Thz-containing peptide hydrazides would be compatible with the NaNO<sub>2</sub>-mediated activation step using a TFA-based cocktail. Since TFA is usually used as a cleavage reagent in Fmoc SPPS, we also envisioned that the cleavage and conversion of the hydrazide could be achieved in a one-pot reaction.

In our initial attempts, the peptide hydrazide H-LYRAG-NHNH<sub>2</sub> (1a) lacking a Thz group at N-terminus was used as a substrate (Table 1). The peptide hydrazide was elongated on 2-

Table 1 Conversion of peptide hydrazide to thioester during global deprotection <sup>a</sup>								
H-LYRAG-N-N-N Cl								
addition of NaNO2 aq (5 eq)				3% MESNa in				
20 min, -10 °C								
then cold Et <sub>2</sub> O precipitation 2a pH 7.3, 30 min, rt H-LYRAG-S(CH <sub>2</sub> ) <sub>2</sub> SO <sub>3</sub> H								
<b>3a</b> (X = H); <b>4a</b> (X = NO <sub>2</sub> )								
entry	global deprotection $(v/v)$					HPLC area ratio [%]		
	TFA	TIS	$H_2O$	<i>m</i> -cresol	thioanisole	1	3	4
1	95	2.5	2.5	0	0	58	20	17
2 <sup>b</sup>	95	2.5	2.5	0	0	61	16	12
3	90	2.5	2.5	5	0	54	39	0
4	90	2.5	2.5	0	5	0	77	13
5 <sup>c</sup>	80	2.5	2.5	5	10	0	87	0
6 <sup>de</sup>	80	2.5	2.5	5	10	0	93	0

<sup>*a*</sup> All reactions were performed as follows: the peptidyl resin was treated under global deprotection conditions at rt for 2 h. The reaction mixture was then incubated at -10 °C for 20 min after addition of 10% aqueous NaNO<sub>2</sub>. The peptide azide precipitated by the addition of cold Et<sub>2</sub>O was subjected to thiolysis with MESNa (3% *w/w*) in 6 M Gn·HCl, 0.2 M sodium phosphate buffer (pH 7.3) at rt for 0.5 h. The product was analysed by HPLC. <sup>*b*</sup> The activating step with NaNO<sub>2</sub> was performed at 0 °C. <sup>*c*</sup> 69% isolated yield. <sup>*d*</sup> N-Terminal Thz-containing peptide (H-**Thz-LYRAG**-NHNH<sub>2</sub>) was used as a substrate. <sup>*e*</sup> 68% isolated yield. HPLC charts: see ESI.<sup>+</sup>

# Journal Name

chlorotrityl chloride resin using a previously AFREQUEED protocol.<sup>11</sup> The peptidyl resin was Derested 9/With 0.37PA triisopropylsilane (TIS)-H<sub>2</sub>O (95:2.5:2.5) to give the peptide hydrazide **1a**. Aqueous NaNO<sub>2</sub> solution (approximately 5 equivalents relative to the hydrazide) was added to the resulting mixture, affording the corresponding azide intermediate **2a**. After a 20 min reaction at -10 °C followed by trituration with Et<sub>2</sub>O, the formed precipitate was dissolved into a buffer containing sodium 2-mercaptoethanesulfonate (MESNa).

HPLC analysis indicated that the desired thioester 3a had been formed, while the hydrazide 1a and the nitrated thioester 4a were also observed (Table 1, entry 1). Higher temperatures during the azidation were not effective (entry 2). The nitration occurs in the phenol ring of Tyr residue, but was not observed during the reaction of peptide hydrazides with NaNO2 in aqueous media.<sup>12</sup> This side reaction was suppressed by the addition of *m*-cresol which is commonly used as an additive in global deprotection of peptides (entry 3). Surprisingly, the addition of thioanisole, which is also used as an additive for peptide deprotection, accelerated the conversion of the hydrazide (entry 4).<sup>13</sup> Use of the both additives resulted in complete conversion and no nitrated product was formed (entry 5). This condition was applied to the conversion of the peptide hydrazide with a Thz group at the N-terminus, successfully producing the N-terminal Thz-containing peptide thioester in 68% isolated yield. Side reactions, including opening of the thiazolidine and subsequent intramolecular or intermolecular ligations were not observed (entry 6). To the best of our knowledge, this is the first example of the direct synthesis of an N-Thz peptide thioester from the peptide hydrazide without any additional protecting groups.

With the optimal conditions in hand, we evaluated the effects of the C-terminal amino acids with the exception of Asp, Asn and Gln, of which thioesters are applicable to NCL reactions<sup>17b</sup> but the corresponding hydrazides cannot be prepared due to intramolecular cyclization between the hydrazide and side-chain during SPPS.<sup>8a</sup> As summarized in Table 2, in all cases, a peptide hydrazides was successfully converted to the corresponding thioester with a modest to high purity (63-95%). Sterically hindered amino acids (Val and Ile) afforded relatively low 70% and 63% yields, respectively, as a result of Curtius rearrangement of the azide to an isocyanate, affording thiocarbamate derivatives (entries 3 and 4).<sup>14</sup> This side reaction also occurred at a higher level than 5% with Glu, Arg and Trp (entries 11, 18 and 19). This probability of formation of a Curtius rearrangement product is consistent with a previous report.<sup>14</sup> The epimerization levels of C-terminal amino acids (Ala, Phe, Ser, Cys, Tyr, and His) were assessed. Peptides containing Ser and His were more sensitive to epimerization whereas the epimerization was suppressed under more acidic thiolysis conditions (entries 8 vs 9 and 15 vs 16).

The synthetic applicability of our protocol was confirmed through the synthesis of the LacZ alpha subunit (LacZ $\alpha$ ) as shown in Figure 2.<sup>15</sup> Since LacZ $\alpha$  has no Cys residue, use of an NCL-desulfurization strategy, formally enabling the assembly of peptides at Xaa-Ala junctions, was envisioned.<sup>16</sup> Peptide Published on 05 June 2018. Downloaded by Hacettepe Universitesi on 05/06/2018 06:32:14.

#### Journal Name

#### globa deprotection thiolysis -Xaa-NHNH-LYRA -LYRA-Xaa-SR then azidation 3a-a Curtius rearrangemen thiolysis HPLC area ratio [%] epimerization entry Хаа product 3 5<sup>b</sup> (L:D) [%] 87 1 Gly 3a n.d. 90 >99:1 2 Ala 3b <1 3 Val 70 8 3c 4 lle 3d 63 13 5 Leu 3e 89 3 6 Phe 3f 90 2 >99:1 7 Pro 3g 89 n.d. 8 Ser 3h 67 n.d. 88:12 9<sup>d</sup> Ser 3h 90 n.d. >99:1 10 72 Thr 3i n.d. 11 Glu 3i 76 7 \_ 12 Cys 3k 79 n.d. >99:1 95 13 Met 31 <1 14 Tyr 3m 82 2 >99:1 15 His 3n 72 n.d. 84:16 16 His 3n 82 n.d. >99:1 17 30 93 n.d. Lys \_ 18 Arg 3p 78 5 19 81 10 Trp 3q

Table 2 Scope of the conversion of peptide hydrazides to peptide thioesters

<sup>*a*</sup> All reactions were performed under similar conditions to that of Table 1 except for global deprotection conditions (TFA-TIS-H<sub>2</sub>O-*m*-cresol-thioanisole = 80:2.5:2.5:5:10). <sup>*b*</sup> n.d. = not detected. <sup>*c*</sup> Standard material as an epimerized peptide was separately synthesized using the corresponding D-isomer, and epimerization ratio was determined by HPLC analysis. <sup>*d*</sup> Buffer solution containing 10% MESNa (pH 4) was used. <sup>*e*</sup> TFA-TIS-H<sub>2</sub>O-*m*-cresol-thioanisole = 75:2.5:2.5:5:15 was used as a global deprotection cocktail.

segments **6–9** covering the entire sequence of LacZ $\alpha$  were prepared as peptide hydrazide for segments **6–8** or peptide acid for **9** by the conventional Fmoc-SPPS (Figure 2B). After construction of the peptide segments, the peptide hydrazides were converted to the corresponding thioesters by the established protocol. This procedure afforded the desired peptide thioesters in 31% isolated yield for **6**, 34% for **7** or 64% for **8**. Notably, no decomposition of N-terminal Thz group in segments **7** and **8** during the activating step was observed.

The first NCL between **9** bearing a Cys residue at the Nterminus and **8** with a Glu-thioester at the C-terminus in 6 M guanidine hydrochloride (Gn·HCl), 0.2 M sodium phosphate buffer (pH 7.3) in the presence of 4-mercaptophenylacetic acid (MPAA) (100 mM) and tris(2-carboxyethyl)phosphine hydrochloride (TCEP·HCl) (50 mM) at 37 °C was complete within 7 h but HPLC analysis of the products of the ligation followed by conversion of Thz to Cys by the addition of MeONH<sub>2</sub>·HCl showed two peaks with the expected molecular weight of the product **10** (Figure 3A, denoted by **10**'). The formation of  $\gamma$ -linked side products had been observed previously in an NCL at the Glu-Cys



COMMUNICATION



Fig. 2 (A) Primary sequence of synthesized LacZ $\alpha$ . (B) Synthetic scheme for LacZ $\alpha$  by sequential NCLs-desulfurization.

site.<sup>17</sup> Kent et al. demonstrated that the use of an increased concentration of MPAA and/or high pH can prevent such by-product formation,<sup>17b</sup> but although we tried to optimize ligation conditions, pH or concentration of peptides and MPAA, suppression of the side reaction could not be achieved.

After being purified by HPLC, the ligated peptide 10 was subjected to the next NCLs. We initially envisioned a one-pot reaction consisting of sequential ligations between peptides 7, 10 and 6 followed by desulfurization, yielding the desired product 12. Peptides 7 and 10 in 6 M Gn·HCl, 0.2 M sodium phosphate, 40 mM methyl thioglycolate as a non-aryl thiol additive,<sup>18</sup> 20 mM TCEP·HCl (pH 7.2) were coupled smoothly. The coupled material was converted to the corresponding Cys form with the aid of MeONH<sub>2</sub>·HCl, and the peptide thioester  $\mathbf{6}$ was added to the resulting mixture to yield the full-length polypeptide. Howev er, the thioester 6 reacted preferentially with the residual MeONH<sub>2</sub> and no ligated product was obtained. We examined the use of [Pd(allyl)Cl]<sub>2</sub> as a reagent to convert Thz to a free Cys,<sup>19</sup> but this did not give the desired product. These results led us revise our synthetic design, which is based on an NCL-deprotection between 7 and 10 followed by removal of MeONH<sub>2</sub> and subsequent one-pot NCL-desulfurization between peptide 6 and 11 using imidazole-aided NCL conditions (Figure 2B).<sup>20</sup>

The reaction sequence for the synthesis of LacZ $\alpha$  is shown in Figure 3A–C. Ligation between the Thz-containing thioester peptide **7** and Cys-peptide **10** followed by deprotection of the Thz ring was conducted under the optimum conditions, affording the three-component ligated product **11** (Figure 3B). After purification by HPLC, the ligated peptide was subjected to the next NCL reaction with thioester **6** under imidazole-aided NCL conditions, allowing us to conduct subsequent free-radicalbased desulfurization of three Cys residues in a one-pot reaction which proceeded efficiently to afford LacZ $\alpha$  (**12**) in 85% isolated yield (Figures 3C and D).

Finally,  $\beta$ -galactosidase activity of the complex between the synthesized LacZ $\alpha$  and expressed LacZ omega subunit (LacZ $\omega$ ) in the conversion of *o*-nitrophenyl- $\beta$ -D-galactopyranoside (ONPG) was evaluated (Figure 4A). Incubation of ONPG with the

ccepted Manu

#### **Journal Name**



Fig. 3 HPLC analyses of reactions for **12**: (A) ligation of **8** and **9** followed by deprotection of Thz (t = 7 h and 11 h); (B) ligation of **7** and **10** followed by deprotection of Thz (t = 5 h and 3 h); (C) ligation of **6** and **11** followed by desulfurization (t = 11 h and 16 h). (D) HPLC analysis and mass spectrum of purified **12**. The calculated mass is 6990.83 Da. Deconvolution of the mass spectrum yielded an observed mass of 6991.98 Da as a proton adduct. Reaction and HPLC conditions: see ESI.† \* MPAA.

synthetic material and bacterial homogenate of *Escherichia coli* DH5 $\alpha$  as a source of LacZ $\omega$  resulted in a colour change resulting from the formation of *o*-nitrophenol through the enzymatic reaction, whereas absence of LacZ $\alpha$  or LacZ $\omega$  did not affect the colour of the reaction solution (Figure 4B). This result clearly indicated that the synthesized LacZ $\alpha$  can exhibit  $\alpha$ -complementation.<sup>15</sup>

In summary, we have developed the simple and practical protocol for converting N-Thz-peptide hydrazides into the corresponding thioesters. A key to our developed protocol is the use of TFA as a reaction medium for  $NaNO_2$ -mediated activation in the presence of thioanisole and *m*-cresol, which can accelerate the desired reaction and suppress the side reaction, respectively. Application of the developed protocol to the synthesis of more-complex proteins will be presented in due course.

This research was supported in part by a Grant-in-Aid for Young Scientists (B) (No. 16K21077) from the Japan Society for the Promotion of Science (JSPS) and research grants from the Futaba Electronics Memorial Foundation.

## **Conflicts of interest**

There are no conflicts to declare.

## Notes and references

- (a) G. Debelouchina and T. W. Muir, *Q. Rev. Biophys.*, 2017, 50, e7; (b) V. Muralidharan and T. W. Muir, *Nat. Methods*, 2006, *3*, 429.
- 2 (a) J. M. Chalker, *Chem. Biol. Drug Des.*, **2013**, *81*, 122; (b) F. Liu and J. P. Mayer, *Top. Curr. Chem.*, **2015**, *362*, 183.
- 3 P. Siman and A. Brik, Org. Biomol. Chem., 2012, 13, 5684.
- 4 P. E. Dawson, T. W. Muir, I. Clark-Lewis and S. B. H. Kent, *Science*, **1994**, *266*, 776.
- 5 a) S. S. Kulkarni, J. Sayers, B. Premdjee and R. J. Payne, *Nat. Rev. Chem.* **2018**, *2*, 122; (b) S. B. H. Kent, *Bioorg. Med.*



Fig. 4 (A) Enzymatic hydrolysis of ONPG by  $\beta$ -D-galactosidase. (B)  $\alpha$ -Complementation assay using ONPG with synthesized LacZ $\alpha$  and crude proteins of *E. coli* DH5 $\alpha$  expressing LacZ $\omega$ . Full length of LacZ expressed by *E. coli* W3110 was used as a positive control.

*Chem.*, **2017**, *25*, 4926; (c) S. B. H. Kent, *Chem. Soc. Rev.*, **2009**, *38*, 338.

- 6 L. Raibaut, N. Olivier and O. Melnyk, Chem. Soc. Rev., 2012, 41, 7001.
- 7 (a) J. A. Camarero, G. J. Cotton, A. Adeva and T. W. Muir, J. Pept. Res., 1998, 51, 303; (b) D. Bang, N. Chopra and S. B. H. Kent, J. Am. Chem. Soc., 2004, 126, 1377; (c) D. Bang and S. B. H. Kent, Angew. Chem. Int. Ed., 2004, 43, 2534; (d) S. Ueda, M. Fujita, H. Tamamura, N. Fujii and A. Otaka, ChemBioChem, 2005, 6, 1983.
- 8 (a) G.-M. Fang, Y.-M. Li, F. Shen, Y.-C. Huang, J.-B. Li, Y. Lin, H.-K. Cui and L. Liu, *Angew. Chem. Int. Ed.* **2011**, *50*, 7645; (b) J.-S. Zheng, S. Tang, Y.-K. Qi, Z.-P. Wang and L. Liu, *Nat. Protoc.*, **2013**, *8*, 2483; (c) J.-S. Zheng, S. Tang, Y.-C. Huang and L. Liu, *Acc. Chem. Res.*, **2013**, *46*, 2475.
- 9 V. Agouridas, O. E. Mahdi, M. Cargoët and O. Melnyk, *Bioorg. Med. Chem.*, 2017, 25, 4938.
- 10 (a) G.-M. Fang, J.-X. Wang and L. Liu, *Angew. Chem. Int. Ed.*,
  2012, *51*, 10347; (b) J. Li, Y. Li, Q. He, Y. Li, H. Li and L. Liu, *Org. Biomol. Chem.*, 2014, *12*, 5435; (c) M. Pan, Y. He, M.
  Wen, F. Wu, D. Sun, S. Li, L. Zhang, Y. Li and C. Tian, *Chem. Commun.*, 2014, *50*, 5837; (d) S. Tang, Y.-Y. Si, Z.-P. Wang, K.-R. Mei, X. Chen, J.-Y. Cheng, J.-S. Zheng and L. Liu, *Angew. Chem. Int. Ed.*, 2015, *54*, 5713; (e) Y.-C. Huang, C.-C. Chen, S.
  Gao, Y.-H. Wang, H. Xiao, F. Wang, C.-L. Tian and Y.-M. Li, *Chem. Eur. J.*, 2016, *22*, 7623.
- 11 G. Stavropoulos, D. Gatos, V. Magafa and K. Barlos, *Lett. Pept. Sci.*, **1995**, *2*, 315.
- 12 The nitration is likely to occur through the coupling of the phenoxyl radical with nitrogen dioxide derived from oxidation of nitric oxide. These species could be generated by homolysis of aromatic *O*-nitroso derivatives. See a reference: D. Koley, O. C. Colón and S. N. Savinov, *Org. Lett.*, 2009, 11, 4172.
- 13 Although we have yet to elucidate the role of thioanisole in the conversion, one potential explanation for the effect of thianisole is that thianisole reacts with nitrosonium cation to afford sulfonium that might work as a more effective oxidant than nitrosonium itself. See a reference: G. I. Borodkin, V. A. Podryvanov, M. M. Shakirov and V. G. Shubin, *J. Chem. Soc., Perkin Trans.* 2, **1995**, 1029.
- 14 A. A. Vinogradov, M. D. Simon and B. L. Pentelute, *Org. Lett.* **2016**, *18*, 1222.
- 15 D. H. Juers, B. W. Matthews and R. E. Huber, *Protein Sci.*, **2012**, *21*, 1792.
- 16 Q. Wan and S. J. Danishefsky, Angew. Chem. Int. Ed., 2007, 46, 9248.
- 17 (a) M. Villain, H. Gaertner and P. Botti, *Eur. J. Org. Chem.*,
   2003, 3267; (b) B. Dang, T. Kubota, K. Mandal, F. Bezanilla and S. B. H. Kent, *J. Am. Chem. Soc.*, 2013, 135, 11911.
- 18 Aryl thiol hampers radical-mediated desulfurization. See reference 10e.
- 19 M. Jbara, S. K. Maity, M. Seenaiah and A. Brik, *J. Am. Chem. Soc.*, **2016**, *138*, 5069.
- 20 K. Sakamoto, S. Tsuda, M. Mochizuki, Y. Nohara, H. Nishio and T. Yoshiya, *Chem. Eur. J.*, **2016**, *22*, 17940.

14

Published on 05 June 2018. Downloaded by Hacettepe Universitesi on 05/06/2018 06:32:

This journal is © The Royal Society of Chemistry 20xx



37x17mm (300 x 300 DPI)