```
A NOVEL PHOTOREARRANGEMENT OF 2,5-DIARYL-1,4-DITHIIN-1,1-DIOXIDE
ACCOMPANIED BY EXTRUSION OF SULFUR DIOXIDE
```

Keiji KOBAYASHI* and Kiyoshi MUTAI Department of Chemistry, College of Arts and Sciences, The University of Tokyo, Komaba, Meguro-ku, Tokyo 153

Photolysis of the title compound afforded 2,5-diarylthiophene. The mechanism involving the valence isomerization to thioketone was supported by the photolysis in n-butylamine, which gave pyrrole derivatives.

Photochemical ring transformation of five membered heterocyclic systems has been the subject of extensive studies.¹⁾ In particular, the fascinating photorearrangement of arylthiophenes has attracted considerable interest since the work of Wynberg and co-workers.²⁾ We report herein an unusual formation of rearranged thiophene in photochemical ring contraction of a six membered heterocyclic compound, 2,5-diaryl-1,4-dithiin-1,1-dioxide (la and lb).

A solution of la in methanol (ca. 1.5×10^{-3} M) was irradiated with a high pressure mercury lamp through a Pyrex filter under nitrogen for 30 min. After removal of the solvent the photolysates were chromatographed on alumina to give 2,5-diphenylthiophene (2a) (4%) and 2,4-diphenylthiophene (3a) (1%). Although intractable mixture other than unreacted la (25%) was also eluted on chromatographic separation, the thiophene products were easily collected by monitoring their strong fluorescence. The isomeric thiophenes were identified by comparison with authentic samples.²⁾ Irradiation of p-chlorophenyl derivative <u>lb</u> similarly gave rearranged thiophene 2b (3%) as well as thiophene 3b (1%) solely due to the loss of sulfur dioxide. The structure of 2b, mp 161-162 °C, NMR (CDCl₂): § 7.20 (2H, s), 7.42(8H, broad q, J=8 Hz), was deduced from the spectral properties.

Although photochemical elimination of sulfur dioxide is frequently reported in the literature,³⁾ the extrusion concomitant with the rearrangement seems pretty rare. Based on the photorearrangement of arylthiophene explored by Wynberg²) one might suspect that 2a is a secondary photoproduct derived from initially formed 3a. However, this possibility is ruled out by the finding that no detectable amount of 2a was obtained when 3a was irradiated. Thus 2a should be formed directly from la via a photochemical process.

Among a number of possible mechanisms envisaged the most plausible one seems to involve the valence isomerization to thicketone $\frac{4}{2}$ bearing a thiete dioxide moiety. Subsequent intramolecular cyclization of 4 followed by the extrusion of sulfur dioxide would give rise to 2a. Alternatively, 4 might eliminate sulfur dioxide prior to the cyclization, leading to cyclopropenyl thioketone 5 as an



intermediate. This rationale by analogy of the van Tamelen-Whitesides mechanism⁴) of thiophene photorearrangement was supported by the photolysis carried out in n-butylamine as solvent. Thus, when <u>la</u> was irradiated in n-butylamine under nitrogen at 0 °C for 8 h, l-butyl-2,5-diphenylpyrrole (<u>6</u>) (2%), l-butyl-2,4-diphenylpyrrole (<u>7</u>) (5%), and 2,4-diphenylthiophene (<u>3</u>) (1%) were obtained by chromatography. The structure of the isomeric pyrroles was identified by direct comparison with authentic samples.^{5,6}) The formation of the pyrroles is most reasonably interpreted in terms of the conversion of the thioketone intermediate into <u>8</u> or <u>9</u> by a ground state aminolysis.⁷)

It is interesting to note that the photochemical behaviour of <u>la</u> is remarkably different from that of the corresponding sulfoxide.⁸⁾ The scope of the photochemistry in thiophenes appears now to extend to involve the thiophene equivalent of six membered heterocyclic systems.

References

- 1) A. Padwa, "Rearrangements in Ground and Excited States," ed by P. de Mayo, Academic Press, New York (1980), Vol. 3, p. 501.
- 2) H. Wynberg, H. van Driel, R. M. Kellogg, and J. Buter, J. Am. Chem. Soc., <u>89</u>, 3487 (1967).
- 3) R. S. Givens, "Organic Photochemistry," ed by P. Padwa, Marcel Dekker, New York (1981), p. 317.
- 4) E. E. van Tamelen and T. H. Whitesides, J. Am. Chem. Soc., <u>93</u>, 6129 (1971).
- 5) K. E. Schlte, J. Reisch, and H. Walker, Chem. Ber., 98, 98 (1965).
- 6) R. M. Rodebaugh and N. H. Cromwell, Tetrahedron Lett., 1967, 2895.
- 7) A. Couture and A. L. Combier, Tetrahedron, 27, 1095 (1971).
- 8) K. Kobayashi and K. Mutai, Tetrahedron Lett., 22, 5201 (1981).

(Received July 21, 1983)