

Tetrahedron Letters 39 (1998) 6899--6900

Highly Diastereoselective Samarium-Reformatsky Reaction of Chiral α-Bromoacetyl-2-oxazolidinones with Aldehydes

Shin-ichi Fukuzawa*, Masahiro Tatsuzawa, and Kohki Hirano

Department of Applied Chemistry, Chuo University, Bunkyo-ku, Tokyo 112-8551, Japan Received 9 April 1998; revised 3 July 1998; accepted 10 July 1998

Abstract : Samarium(11) iodide mediated highly diastereoselective Reformatsky-type reaction of chiral α -bromoacetyl-2-oxazolidinones with aldehydes. © 1998 Elsevier Science Ltd. All rights reserved.

Samarium(II) iodide often promises highly stereoselective reactions as the result of chelation control of the samarium atom with the oxygen or nitrogen moiety in organic molecules.¹ Highly stereoselective intra- and intermolecular 1,2- and 1,3-asymmetric inductions are typical examples of this chelation control.² We and other research groups have recently reported a diastereoselective ketyl-alkene coupling reaction using chiral esters, acetals, and aldehydes as chiral auxiliaries.³ We now succeeded in the highly diastereoselective intermolecular samarium-Reformatsky reaction using a chiral α -bromoacetyl-2-oxazolidinone as the chiral auxiliary.⁴⁻⁶ We now present the preliminary results of this reaction.

Scheme 1



To a THF (20 ml) solution of Sml₂ (0.1 M, 2.0 mmol) was added a mixture of α -bromoacetyl-2oxazolidinone (2) (1.0 mmol) and aldehyde (1.0 mmol) at -78 °C and the resulting mixture was stirred at the same temperature for 0.5 h during which time the color of the solution turned from deep green to yellow brown. After the usual acidic workup, the β -hydroxy carboximide (3) was isolated by column chromatography (SiO₂, hexane/ethyl acetate = 3 : 1). The diastereomeric excess (de) of **3** was determined by GC and/or HPLC. We examined the stereoselectivity of the reaction of isobutyraldehyde with three chiral oxazolidinone auxiliaries. X = Pr¹ (78 % de), X = Ph (78 % de), and X = PhCH₂ (78 % de). Table 1 summarizes the results of the samarium-Reformatsky reaction of chiral α -bromoacetyl-2-oxazolidinones (2; X = Pr¹) with various aldehydes (1). The yields of the products were good to excellent and the de values were high for the straight chain, branched aliphatic aldehydes and aromatic aldehydes (up to 99% de). Absolute configuration of some of the products were determined by transforming into the corresponding ethyl esters. ^{7,8}

The origin of the high stereoselectivity has not been clarified but the reaction should proceeded *via* the samarium enolate. Further studies are now in progress.

Acknowledgment: This work was partially supported by Sumitomo Foundation Ltd.

Entry	1	%, yield of 3 ^b	%, de ^c	$[\alpha]_D^{25}$ of 4d Config.
1	≻сно	92	78	+14.42 (c= 1.09, <i>R</i> MeOH)
2	сно	95	76	
3	<->Сно	78	84	
4) сно	87	>99	+43.48 (c≍1.48, <i>ND</i> CHCl ₃)
5	C ₇ H ₁₅ CHO	81	82	+10.43 (c=1.24, <i>S</i> CHCl ₃)
6	СНС) 86	86	
7	Пр-сно	67	64	+37.50 (c= 0.30, <i>R</i> CHCl ₃)

Table 1. Reaction of α -Bromoacetyl-2-oxazolidinones with Aldehydes^a

^a **1** (1.0 mmol), **2** (1.0 mmol), Sml₂ (2.0 mmol); THF (20 ml), -78 °C, 0.5 h. ^b Isolated yield. ^c Determined by GC and/or HPLC.

References

- (1) For a review, Molander, G. A.; Harris, C. H. Chem. Rev. 1996, 96, 307.
- (2) (a) Kito, M.; Sakai, T.; Yamada, K.: Matsuda, F. Shirahama, H. *Synlett*, **1993**, 158. (b) Kawatsura, M.; Matsuda, F.; Shirahama, H. *J. Org. Chem.* **1994**, *59*, 6900. (c) Kawatsura, M.; Hosaka, K.; Matsuda, F.; Shirahama, H. *Synlett*, **1995**, 729. (d) Kawatsura, M.; Dekura, F.; Shirahama, H.; Matsuda, F. *Synlett*, **1996**, 373. (e) Evans, D. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1990**, *112*, 6447.
- (3) (a) Fukuzawa, S.; Seki K.; Tatsuzawa, M.; Mutoh, K. J. Am. Chem. Soc. 1997,119, 1482. (b) Molander G.
 A.; McWilliams, J. C.; Noll, B. C. J. Am. Chem. Soc. 1997,119, 1265. (c)Taniguchi, N.; Uemura, M. Tetrahedron Lett, 1997, 38, 7199.
- (4) For a review on chiral oxazolidinones in asymmetric synthesis. Ager, D. J.; Prakash, I.; Schaad, D. R. *Chem. Rev.* **1996**, *96*, 835-875.
- (5) Zinc promoted Reformasky reaction with α-bromoacetyl-2-oxazolidinone has been studied, but only one experimental result are reported. Ito, Y.; Terashima, S. *Tetrahedron*, **1991**, *47*, 2821.
- (6) Diastereoselective intramolecular samarium-Reformatsky reaction has been reported. Molander G. A.; Etter, J. B.; Harring, L. S.; Thorel, P.-J. J. Am. Chem. Soc. **1992**,114, 1656.
- (7) Evans, D.A.; Bartroli, J.; Shih, T.L. J. Am. Chem. Soc. 1981, 114, 2127.
- (8) (a) Sugiura, T.; Yoshikawa, M.; Yoneda, T.; Tai, A. Bull. Chem. Soc. Jpn. 1990, 63, 1080. (b) Basile, T.; Tagliavini, E.; Trombini, C.; Umani-Ronchi, A. Synthesis, 1990, 305. (c)Soai, K.; Yamanoi, T.; Hikima, H.; Oyamada, H. Chem. Commun. 1985, 138.