

A Preparation of β -Oxoester Enolate Equivalents from SmI_2 and α -Bromoalkanoates


Kiitiro Utimoto,* Toshiki Matsui, Tsutomu Takai, and Seiji Matsubara*

Department of Material Chemistry, Faculty of Engineering, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-01

(Received November 18, 1994)

A treatment of ethyl bromoacetate with two molar equivalent of samarium diiodide at -50°C for 15 min in THF produces a β -oxoester enolate equivalent; the reagent reacts with ketones or aldehydes to give δ -hydroxy- β -oxoesters in excellent yields.


Samarium diiodide (SmI_2) has been recognized to be an effective electron donating reagent in both Reformatsky type reactions¹ and Barbier type reactions.² Those have been performed by the treatment of a mixture of carbonyl compounds and organic halides with SmI_2 to produce the corresponding alcohols. Attempts to prepare organosamarium(III) species from SmI_2 and organic halides before an addition of carbonyl compounds, often afforded unsatisfactory results.³ One of the reasons might come from the instability of samarium(III) species. We assumed that Reformatsky type reagent might be obtained from α -bromoester and SmI_2 . In contrast to the assumption, a treatment of cyclohexanone **3a** with a mixture of ethyl α -bromoacetate **1a** and two molar equivalent of SmI_2 gave δ -hydroxy- β -oxoester **4a** in good yield (eq 1). The result suggested the formation of β -oxoester enolate **2a**.⁴ This paper describes a novel route to δ -hydroxy- β -oxoester.^{5,6}



A solution of ethyl bromoacetate (**1a**, 0.17 g, 1.0 mmol) in THF (3.0 ml) was added to a THF solution of SmI_2 (0.1 M, 20 ml, 2.0 mmol) at -60°C . The mixture was stirred for 15 min at -50°C . The color of the reaction mixture turned from dark blue to light yellow during the stirring. A THF solution of cyclohexanone (**3a**, 0.5 g, 0.5 mmol) was added to the reaction mixture. The resulting mixture was stirred for 15 min at the same temperature and for another 15 min at 0°C . The reaction mixture was worked up with aqueous condition and the crude product was purified by silica gel column chromatography; ethyl 5-(1-hydroxycyclohexyl)-3-oxobutanoate **4a** was obtained in 98% yield (112 mg). Other carbonyl compounds afforded corresponding δ -hydroxy- β -oxoesters in good yields; the results are summarized in Table 1. Benzaldehyde, that is easily reduced with SmI_2 , also gave the corresponding product in high yield (entry 9). The use of α,β -unsaturated aldehyde gave the 1,2-adduct exclusively (entry 10). The results showed a complete consumption of SmI_2 before an addition of aldehyde under the reaction conditions. The intermediate **2a** also added to β -tetralone to give the corresponding product in good yield (entry 5).⁷

The intermediate **2** was effective to give diketo ester **6** by


Table 1. Preparation of δ -Hydroxy- β -ketoesters by Self-condensed Reformatsky Type Reagent ^a



Entry	R	Carbonyl compound	Time / h	Yield / % ^b
1	H	Cyclohexanone	0.5	98
2	H	Cyclopentanone	1.0	>98
3	H	Propanal	1.0	87
4	H	4-Heptanone	1.0	>98
5	H	β -Tetralone	0.5	82
6	H	2-Methylcyclohexanone	1.0	95 (10/90) ^c
7	H	4- <i>t</i> -Butylcyclohexanone	1.0	>98 (12/88) ^c
8	Me	Cyclohexanone	0.2	95 ^d
9	H	Benzaldehyde	0.5	>98
10	H	(<i>E</i>)-2-Hexenal	0.5	>98

a) β -Hydroxyester was not detected by nmr in all cases. b) Isolated yield. c) A ratio of diastereomers. d) Isolated as a diastereomeric mixture.

Table 2. Preparation of *t*-Butyl 3,5-Diketoalkanoate^a




Entry	R	Yield / % ^b	Product
1	Et-	26 ^c	Et-CO-CH2-CO-CO2t-Bu
2	EtOCH2-	65	EtOCH2-CO-CH2-CO-CO2t-Bu
3		75	

a) $\text{BrCH}_2\text{CO}_2t\text{-Bu}$ (1.0 mmol), SmI_2 (2.0 mmol), and anhydride (**5**, 0.48 mmol) were used. b) Isolated Yield. c) *t*-Butyl 4-ethoxycarbonyl-3-oxobutanoate was also isolated in 23% yield.


the reaction with the mixed anhydrides **5**⁸ as shown in Table 2.⁹ In these cases, the intermediate was prepared by a condensation of the *t*-butyl ester that are often tolerate to nucleophilic attack.

In these reactions, alkoxy group on the substrate (ethoxy and tetrahydrofuranyl in entries 2 and 3) promoted the reaction.¹⁰

The intermediate **2** was stable under $-50\text{ }^{\circ}\text{C}$, but the thermal isomerization was observed when the temperature of reaction mixture was raised to $0\text{ }^{\circ}\text{C}$. Quenching the reagent at $-50\text{ }^{\circ}\text{C}$ with DCl in D_2O afforded 4-deuterio compound **8** quantitatively (eq 2). In contrast, after the reagent was warmed up to $0\text{ }^{\circ}\text{C}$, quenching it with DCl in D_2O afforded 2-deuterio compound **9** quantitatively. The intermediate **2a** isomerized into the stable enolate **7** that did not give any adduct with ketone or aldehyde.¹¹



An attempt of cross condensation was also realized by a proper combination of α -haloesters. An equimolar mixture of methyl 2-chloropropionate and t-butyl bromoacetate was treated with SmI_2 and followed by an addition of benzaldehyde; t-butyl 5-hydroxy-5-phenyl-3-ketopentanoate was isolated in 84% yield as a diastereomeric mixture (eq 3).¹²



References and Notes

- P. Girard, J. L. Namy, and H. B. Kagan, *J. Am. Chem. Soc.*, **102**, 2693 (1980); J. Inanaga, Y. Yokoyama, Y. Handa, and M. Yamaguchi, *Tetrahedron Lett.*, **32**, 6371 (1991); K. Otsubo, K. Kawamura, J. Inanaga, and M. Yamaguchi, *Chem. Lett.*, **1987**, 1487; G. A. Molander and J. B. Etter, *J. Am. Chem. Soc.*, **109**, 6556 (1987).
- H. B. Kagan, J. Collin, J. L. Namy, C. Bied, F. Dallemer, and A. Lebrun, *J. Alloys Comp.*, **192**, 191 (1993); J. J. Inanaga, O. Ujikawa, Y. Handa, K. Otsubo, and M. Yamaguchi, *J. Alloys Comp.*, **192**, 197 (1993); G. A. Molander, *Chem. Rev.*, **92**, 29 (1992); J. A. Soderquist, *Aldrichmica Acta*, **24**, 5 (1991).
- J. L. Namy, J. Collin, C. Bied, and H. B. Kagan, *Synlett*, **1992**, 733; D. P. Curran, T. L. Fevig, C. P. Jasperse, M. J. Tottleben, *Synlett*, **1992**, 943; J. Inanaga, M. Ishikawa, and M. Yamaguchi, *Chem. Lett.*, **1987**, 1485.
- Self condensation of lithium enolate to produce β -oxoester: D. F. Sullivan, R. P. Woodbury, and M. W. Rathke, *J. Org. Chem.*, **42**, 2038 (1977); M. W. Rathke and D. F. Sullivan, *J. Am. Chem. Soc.*, **95**, 3050 (1973).
- Dianion of β -oxoesters: S. N. Huckin and L. Weiler, *Can. J. Chem.*, **52**, 2157 (1974).
- A part of the results described here has been reported; the 40th Symposium on Organometallic Chemistry, Japan, Sapporo, 1993, Abstr., PB109; the 67th Annual Meeting of the Chemical Society of Japan, Tokyo, 1994, Abstr., 4J228.
- The dianion, obtained from ethyl acetoacetate and lithium diisopropylamide,⁵ acted as a strong base to β -tetralone and resulted in an enolization.
- G. Friour, A. Alexakis, G. Cahiez, and J. Normant, *Tetrahedron*, **40**, 683 (1984).
- δ,β -Diketo esters: M. Yamaguchi, K. Shibato, and I. Hirao, *Chem. Lett.*, **1985**, 1145; S. N. Huckin and L. Weiler, *Can. J. Chem.*, **52**, 1343 (1973); T. M. Harris and C. M. Harris, *Tetrahedron*, **33**, 2159 (1977).
- Coordination of alkoxy group to samarium reagent may accelerate the nucleophilic addition; see, X. Chen, E. R. Hortelaro, E. L. Eliel, and S. V. Frye, *J. Am. Chem. Soc.*, **114**, 1778 (1992).
- I. Shibata, M. Nishio, A. Baba, and H. Matsuda, *Chem. Lett.*, **1993**, 1219.
- Financial supports by the Ministry of Education, Science and Culture (Grant-in-Aid #05235106, #06403025, #06241241) and the Ono Pharmaceutical Company are deeply acknowledged.