This article was downloaded by: [State University NY Binghamton] On: 29 September 2014, At: 11:12 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lcar20

Synthesis of Some Novel Glucosyl Triazoles from 2,3,4,6-Tetra-O-pivaloyl-D-glucopyranosyl Azide

Chao Shen ^{a b} , Hui Zheng ^b , Pengfei Zhang ^b & Xinzhi Chen ^b ^a Department of Chemical and Biological Engineering , Zhejiang University , Hangzhou, 310027, China

^b College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou, 310036, China Published online: 26 Aug 2010.

To cite this article: Chao Shen , Hui Zheng , Pengfei Zhang & Xinzhi Chen (2010) Synthesis of Some Novel Glucosyl Triazoles from 2,3,4,6-Tetra-O-pivaloyl-D-glucopyranosyl Azide, Journal of Carbohydrate Chemistry, 29:4, 155-163, DOI: <u>10.1080/07328303.2010.487231</u>

To link to this article: <u>http://dx.doi.org/10.1080/07328303.2010.487231</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Synthesis of Some Novel Glucosyl Triazoles from 2,3,4,6-Tetra-O-pivaloyl-Dglucopyranosyl Azide

Chao Shen,^{1,2} Hui Zheng,² Pengfei Zhang,² and Xinzhi Chen²

¹Department of Chemical and Biological Engineering, Zhejiang University Hangzhou 310027, China ²College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310036, China

In the present study, we have synthesized a series of novel glucosyl triazoles for the first time. The glucosyl triazoles **4a–e** were synthesized by reaction of some azidoglycosides with various terminal alkynes via a copper-catalyzed [3+2] cycloaddition ("click chemistry") and were deprotected to afford the corresponding glucosyl triazoles **5a–e** in good yields. The structures of the new compounds were determined by IR, NMR spectroscopy, and mass spectrometry. The antitumor (human cervical cancer cell) activity was evaluated for the target compounds.

Keywords Click chemistry; Glucosyl triazoles; Azidoglycosides

INTRODUCTION

Organic compounds bearing triazoles of different pharmacodynamic nuclei were found to possess various biological activity. Glucosyl triazoles and their derivatives also have received considerable attention because of their bioactivity and many applications in organic and medicinal chemistry.^[1] For example, Poulsen's group has reported that a series of glucosyl triazole benzene sulfonamide derivatives are a novel class of carbonic anhydrase inhibitors.^[2] The classical method for the preparation of 1,2,3-triazoles is the Huisgen reaction.^[3] This powerful and reliable Cu-catalyzed 1,3-dipolar cycloaddition has

Received January 7, 2010; accepted April 17, 2010.

Address correspondence to Pengfei Zhang, College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310036, China. E-mail: zpf100@163.com

found widespread applications in combinatorial chemistry for drug discovery,^[4] material science,^[5] and bioconjugation.^[6] Since that time triazole-linked glycoconjugates have become increasingly useful and important in glycobiology.^[7] Glycoconjugates exert important effects on many complex biological processes,^[8] including the cellular recognition in the processes of immune response,^[9] inflammation,^[10] tumor metastasis,^[11] and viral infections. In addition, glycosylation of proteins and lipids are key factors in modulating their structures and functions. Therefore, the preparation of these compounds has become increasingly important. Herein, we report a convenient method for the synthesis of novel glucosyl triazole derivatives from D-glucose by means of a three-step sequence involving stereospecific displacement with azide anion, copper-catalyzed [3+2] cycloaddition, and deprotection of the pivaloyl group.

RESULTS AND DISCUSSION

During our previous research,^[12] we developed a novel route for the synthesis of thiazol-2(3H)-imine-linked glycoconjugates, which were screened for their antitumor activities against Hela, HCT-8, and Bel-7402. Some of the compounds showed moderate cytotoxicity against HCT-8. That study forms part of our research program to evolve new methodologies for the construction of novel glycoconjugates and to discover new lead molecules with antitumor activities. Further synthesis and biological investigation of new glucosyl triazoles containing different active sugar moieties for antitumor activity are reported here.

The glucosyl triazole **4a** was used as a model to study the rate of triazole formation.^[13] Using a procedure adapted from the literature,^[14,15] the 2,3,4,6-tetra-O-pivalovl-D-glucopyranosyl azide **2** is available from compound 1 by means of stereospecific displacement with azide anion. Then reaction of phenylacetylene with the azide 2 in the presence of a CuSO₄/ascorbate mixture affords exclusively the glucosyl triazole 4a with complete retention of anomeric stereochemistry and in high yield. The effects of various solvents were studied for the preparation of **4a** and comparative results are presented in Table 1. As a result, the reaction of azide **2** proceeded smoothly in *tert*-butanol within a short period at room temperature to afford the desired products in very high yield (Table 1, entry 8). In other cases, the yields varied 75% to 91%. These results encouraged further studies with various other alkynes such as 2-ethynylpyridine, 4-bromophenylacetylene, 4-fluorophenyla-cetylene, and 4methylphenylacetylene. These alkynes readily reacted with glucosyl azides under identical conditions to produce triazole-linked glycosides in high yields (Table 2, entries 2–4). The structure of all new compounds was confirmed by ¹H NMR, ¹³C NMR, IR, and elemental analysis. The O-pivaloyl group of glucosyl triazoles **4a-e** was subsequently removed to obtain the fully deprotected sugar analog **5a-e** in quantitative or near-quantitative yields (Scheme 1).

Pivo Pivo 2	V O O O O O O O O	───► Piv F	OPiv OPivO OPiv OPiv OPiv Aa	N N
Entry	Solvent	Temp (°C)	Time (h)	Yield (%) ^b
1 2 3 4 5 6 7 8	CH ₃ CN CH ₂ Cl ₂ Toluene THF Methanol Isopropanol <i>tert</i> -Butanol <i>tert</i> -Butanol	40 40 40 40 40 40 40 25	2.0 2.0 2.0 2.0 2.0 2.0 2.0 2.0 2.5	83 80 75 78 88 91 95 95

Table 1: Study of formation of glucosyl triazole 4a^a

^aAll reactions were carried out using **2** (0.5 M), phenylacetylene (0.5 M), 20 mol% CuSO₄, 40 mol% sodium ascorbate (relative to substrate). ^bIsolated yield after chromatography.

Next, some of the newly synthesized glucosyl triazoles **5a–e** were screened for their antitumor activity against Hela (cervical carcinoma), HCT-8 (colon carcinoma), and Bel-7402 (liver carcinoma). The cytotoxicity was evaluated by MTT assay.^[16] However, all the compounds showed very weak cytotoxicity against Bel-7402, Hela, and HCT-8.

In conclusion, we have synthesized a series of triazole derivatives bearing sugar moieties, providing a convenient method for constructing such glucosyl triazole derivatives under very mild conditions. The preliminary biological

Entry	Ar	Product	Time (h)	Yield (%) ^b	Mp (°C)
1	Ph	4a	2.5	92	230–231
2	p-Br-C ₆ H ₄	4b	2.5	88	218–220
3	p-F-C ₆ H ₄	4c	2.5	92	208–210
3	p-CH ₃ -C ₆ H ₄	4d	2.5	95	215–217
4	C ₅ H ₄ N	4e	2.5	96	214–216

Table 2: Synthesis of different D-glucose-derived 1,2,3-triazoles^a

^aAll reactions were carried out using **2** (0.5 M), alkynes (0.5 M), 20 mol% CuSO₄, 40 mol% sodium ascorbate (relative to substrate).

^blsolated yield after chromatography.



Scheme 1: A general glucosyl triazoles synthesis.

evaluation of compounds **5a–e** showed no antitumor activities. Further syntheses and biological studies of new triazoles containing different active sugar moieties for biological activity are continued by our group.

EXPERIMENTAL

General Methods

1,2,3,4,6-Penta-*O*-pivaloyl-D-glucopyranose (1) was prepared according to our previous reports.^[12] Other chemicals and solvents were either purchased or purified by standard techniques. Analytical TLC was performed on a Merck precoated TLC (silica gel 60 F254) plate. Melting points were recorded on an X₄-Data microscopic melting point apparatus and are uncorrected. IR spectra were recorded on a Nicolet 380 FT-IR spectrophotometer using KBr discs. Optical rotation values were measured on a Perkin Elmer P241 polarimeter. ESI-MS were acquired on a Bruker Esquire 3000 plus spectrometer. ¹H and ¹³C NMR spectra of samples in CDCl₃ or CD₃OD were recorded on a Bruker Avance 400 spectrometer using TMS as internal standard. Elemental analyses were performed on Carlo-Erba 1106.

General procedure for the synthesis of glucosyl triazoles via click chemistry

To a mixture of the corresponding 2,3,4,6-tetra-O-pivaloyl-D-gluco pyranosyl azide **2** (541 mg, 1.0 mmol) and alkyne (1.0 mmol) in *tert*-butyl alcohol (1 mL), 0.5 mL of an aqueous solution of CuSO₄ (50 mg, 0.2 mmol) and 0.5 mL of an aqueous solution of sodium ascorbate (79.2 mg, 0.4 mmol) were added. The reaction mixture was stirred vigorously for 24 h. The suspended product was filtered and washed with water. The crude product was purified by column chromatography on silica gel, eluting with hexane/ethyl acetate mixtures of increasing polarity to afford the triazole products.

4-Phenyl-1-(2,3,4,6-tetra-O-pivaloyl-1-D-glucopyranosyl)-1,2,3triazole (4a)

Yellow solid; yield: 92%; mp 216–220°C; $[\alpha]_D^{20} = -18.0$ (*c* 0.1, CH₃OH); ¹H NMR (400 MHz, DCCl₃): δ 8.05 (s, 1H), 7.85 (d, J = 8.6 Hz, 2H), 7.46 (t, 2H), 7.38 (t,1H), 5.87 (d, J = 8.0 Hz, 1H), 5.65 (t, J = 9.6 Hz, 2H), 5.60 (d, J = 3.8 Hz, 1H), 5.36 (d, J = 3.3 Hz, 1H), 4.74 (m, 1H), 4.31 (m, 1H), 4.04 (m, 1H), 0.92–1.25 (m, 36H, Piv-CH₃); ¹³C NMR (100 MHz, DCCl₃): δ 177.78, 176.52, 147.43, 144.60, 129.00, 127.95, 127.58, 125.01, 85.12, 74.64, 71.14, 69.15, 66.18, 60.42, 37.76, 25.79–26.12; IR (KBr, cm⁻¹) υ : 2970, 2077, 1744, 1637, 1481, 1400, 1368, 1280, 1140, 1037.

4-Pyridyl-1-(2,3,4,6-tetra-O-pivaloyl-1-D-glucopyranosyl)-1,2,3triazole (4b)

Yellow solid; yield: 95%; mp 179–181°C; $[\alpha]_D^{20} = -43.2$ (*c* 0.1, CH₃OH); ¹H NMR (400 MHz, DCCl₃): δ 8.60 (s, 1H), 8.37 (d, J = 3.6 Hz, 1H), 8.11 (t, 1H), 7.78 (t, 1H), 7.76 (d, J = 12.2 Hz, 1H), 5.97 (d, J = 9.2 Hz, 1H), 5.55 (m, 2H), 5.32 (m, 1H), 4.21 (m, 2H), 4.05 (d, J = 2.4 Hz, 1H), 1.89 (m, 1H), 0.94–1.21 (m, 36H, Piv-CH₃); ¹³C NMR (100 MHz, DCCl₃): δ 177.71, 177.12, 177.10, 155.3, 149.20, 137.21, 131.41, 129.52, 123.64, 97.62, 78.21, 73.69, 69.15, 69.82, 62.42, 36.76, 25.79–26.12; IR (KBr, cm⁻¹) υ : 3431, 2974, 1742, 1601, 1479, 1400, 1281, 1141, 1034. Anal. Calcd for C₃₃H₄₈N₄O₉: C, 61.55; H, 7.82; N, 8.78. Found: C, 61.58; H, 7.80; N, 8.75.

4-(4-Bromophenyl)-1-(2,3,4,6-tetra-O-pivaloyl-1-Dglucopyranosyl)-1,2,3-triazole (4c)

Yellow solid; yield: 88.1%; mp 220–222°C; $[\alpha]_D^{20} = -38.2$ (*c* 0.1, CH₃OH); ¹H NMR (400 MHz, DCCl₃): δ 7.97 (s, 1H), 7.70 (d, J = 5.6 Hz, 2H), 7.56 (t, 2H), 5.99 (d, J = 9.6 Hz, 1H), 5.55 (m, 2H), 5.36 (d, J = 4.2 Hz, 1H), 4.21 (m, 2H), 4.06 (m, 1H), 0.92–1.25 (m, 36H, Piv-CH₃); ¹³C NMR (100 MHz, DCCl₃): δ

160 C. Shen et al.

177.71, 176.74,131.9, 129.72, 127.24, 122.36, 85.94, 77.18, 71.84, 69.96, 66.94, 61.14, 38.76, 25.53–26.92; IR (KBr, cm⁻¹) υ : 3422, 2974, 1742, 1637, 1481, 1403, 1368, 1140, 1040. Anal. Calcd for $C_{34}H_{48}BrN_3O_9$: C, 56.62; H, 6.73; N, 5.93. Found: C, 56.65; H, 6.70; N, 5.91.

4-(4-Fluorophenyl)-1-(2,3,4,6-tetra-O-pivaloyl-1-Dglucopyranosyl)-1,2,3-triazole (4d)

White solid; yield: 89.1%; mp 190–192°C; $[\alpha]_D^{20} = -21.6 (c \ 0.1, CH_3OH)$; ¹H NMR (400 MHz, DCCl₃): δ 8.02 (s, 1H), 7.82 (d, 2H), 7.82 (t, 2H), 5.96 (d, J = 9.6 Hz, 1H), 5.58 (m, 2H), 5.38 (d, J = 2.4 Hz, 1H), 4.20 (m, 2H), 4.11 (d, J = 12.4 Hz, 1H), 0.95–1.26 (m, 36H, Piv-CH₃); ¹³C NMR (100 MHz, DCCl₃): δ 178.71, 176.99, 132.1, 128.72, 127.24, 122.36, 85.94, 77.20, 71.86, 69.98, 66.90, 61.20, 38.73, 25.53–26.92; IR (KBr, cm⁻¹) υ : 3436, 2976, 1740, 1641, 1485, 1406, 1360, 1141, 1039. Anal. Calcd for C₃₄H₄₈FN₃O₉: C, 61.76; H, 7.22; N, 6.43. Found: C, 61.73; H, 7.20; N, 6.45.

4-(4-Methylphenyl)-1-(2,3,4,6-tetra-O-pivaloyl-1-Dglucopyranosyl)-1,2,3-triazole (4e)

White solid; yield: 92.6%; mp 208–210°C; $[\alpha]_D^{20} = +3.2$ (*c* 0.1, CH₃OH); ¹H NMR (400 MHz, DCCl₃): δ 8.00 (s, 1H), 7.74 (d, J = 5.6 Hz, 2H), 7.25 (t, 2H), 5.86 (d, J = 11.6 Hz, 1H), 5.55 (d, J = 4.2 Hz, 1H), 5.34 (m, 1H), 4.71 (m, 2H), 4.45 (d, J = 2.2 Hz, 1H), 4.30 (d, J = 6.2 Hz, 1H), 2.39 (s, 1H), 0.92–1.25 (m, 36H, Piv-CH₃); ¹³C NMR (100 MHz, DCCl₃): δ 177.46, 8176.18, 176.18, 144.29, 137.99, 129.11, 126.46, 126.24, 115.86, 86.16, 76.82, 76.19, 66.02, 65.94, 60.03, 38.38, 29.19–26.47; IR (KBr, cm⁻¹) υ : 3443, 2352, 2118, 1746, 1649, 1540, 1398, 1280, 1146. Anal. Calcd for C₃₅H₅₁N₃O₉: C, 63.86; H, 7.74; N, 6.50; Found: C, 63.86; H, 7.74; N, 6.50.

Representative procedure for the depivaloylation of glucosyl triazoles

The corresponding glucosyl triazole (1 mmol) was dissolved in 10 mL of methanol. Freshly prepared sodium methoxide was then added portion-wise and the reaction was monitored by TLC. Upon completion, Amberlite resin (H⁺ form) was added until the solution was at neutral pH, and the solution was filtered and concentrated in vacuo. The crude product was taken up in hot water and treated with activated charcoal. The product was recrystallized from water to give a white solid. 4-Phenyl-1-D-glucopyranosyl-1,2,3-triazole (**5a**) was reported by Basu et al.^[16]

4-Pyridyl-1-D-glucopyranosyl-1,2,3-triazole (5b)

Yellow solid; yield: 97%; mp 198–200°C; $[\alpha]_D^{20} = -11.2$ (*c* 0.055, CH₃OH); ¹H NMR (400 MHz, CD₃OD): δ 8.53 (s, 1H), 8.37 (d, J = 10.2 Hz, 1H), 8.13 (t, 1H), 7.81 (t, 1H), 7.73 (d, J = 6.4 Hz, 1H), 5.72 (d, J = 9.6 Hz, 1H), 4.26 (t, 1H), 4.03 (d, J = 8.2 Hz, 1H), 3.78–3.71 (m, 4H); ¹³C NMR (100 MHz, CD₃OD): δ 177.69, 177.10, 177.08, 155.30, 149.20, 137.21, 131.43, 129.49, 123.64, 95.62, 76.09, 71.26, 69.15, 62.42; Anal. Calcd for C₁₃H₁₆N₄O₅: C, 50.68; H, 5.96; N, 18.02. Found: C, 50.70; H, 5.93; N, 18.16.

4-(4-Bromophenyl)-1-D-glucopyranosyl-1,2,3-triazole (5c)

Yellow solid; yield: 96%; mp 215–217°C; $[\alpha]_D^{20} = -7.8$ (c 0.062, CHCl₃); ¹H NMR (400 MHz, CD₃OD): δ 8.01 (s, 1H), 7.81 (d, J = 11.2 Hz, 2H), 7.72 (d, J = 4.6 Hz, 2H), 5.81 (d, J = 9.2 Hz, 1H), 4.41 (t, 1H), 4.21 (d, J = 8.2 Hz, 1H), 3.65–3.70 (m, 4H); ¹³C NMR (100 MHz, CD₃OD): δ 177.66, 176.45, 131.9, 129.72, 127.24, 122.36, 85.94, 77.08, 71.76, 69.77, 66.80, 61.19; Anal. Calcd for C₁₄H₁₆BrN₃O₅: C, 43.89; H, 4.02; N, 10.56. Found: C, 43.86; H, 4.10; N, 10.52.

4-(4-Fluorophenyl) -1-D-glucopyranosyl -1,2,3-triazole (5d)

White solid; yield: 95%; mp 182–185°C; $[\alpha]_D^{20} = -15.8 (c \ 0.068, CHCl_3)$; ¹H NMR (400 MHz, CD₃OD): δ 8.12 (s, 1H), 7.86 (d, J = 6.2 Hz, 2H), 7.73 (d, J = 3.2 Hz, 2H), 5.79 (d, J = 9.8 Hz, 1H), 4.66 (t, 1H), 4.21 (d, J = 3.6 Hz, 1H), 3.69–3.72 (m, 4H); ¹³ C NMR (100 MHz, CD₃OD): δ 177.53, 176.31, 131.09, 129.14, 127.12, 122.28, 85.44, 77.11, 72.03, 69.66, 66.52, 61.28; Anal. Calcd for C₁₄H₁₆FN₃O₅: C, 51.56; H, 4.81; N, 12.23. Found: C, 51.52; H, 4.78; N, 12.19.

4-(4-Methylphenyl)-1-D-glucopyranosyl -1,2,3-triazole (5e)

White solid; yield: 95%; mp 216–219°C; $[\alpha]_D^{20} = -22.1 (c \ 0.075, \text{CHCl}_3)$; ¹H NMR (400 MHz, CD₃OD): δ 8.03 (s, 1H), 7.56 (d, J = 5.6 Hz, 2H), 7.31 (d, J = 5.6 Hz, 2H), 5.66 (d, J = 9.4Hz, 1H), 4.70 (t, 1H), 4.18 (d, J = 2.6 Hz, 1H), 3.68–3.72 (m, 4H), 2.35 (s, 3H); ¹³C NMR (100 MHz, CD₃OD): δ 177.21, 176.22, 175.58, 144.28, 138.20, 128.02, 126.24, 126.11, 115.75, 86.25, 76.74, 76.09, 66.12, 65.34, 60.23, 38.14. Anal. Calcd for C₁₅H₁₉N₃O₅: C, 56.26; H, 5.82; N, 13.11. Found: C, 56.30; H, 5.81; N, 13.14.

Biological activity assays

Cell culture

Three different human carcinoma cell lines were used for cytotoxicity determination: Hela (cervical carcinoma), HCT-8 (colon carcinoma), and Bel-7402

162 C. Shen et al.

(liver carcinoma). They were obtained from the American Type Culture Collection (ATCC) and cultured in RPMI-1640 medium supplemented with 10% FBS, 100 units/mL of penicillin, and 100 μ g/mL of streptomycin. Cells were maintained at 37°C in a humidified atmosphere of 5% CO₂ in air.

Cytotoxicity Analysis

Briefly, cells were plated in 96-well culture plates (10^4 cells per well) and incubated overnight at 37° C in a 5% CO₂ incubator. Then, compounds were added to the wells to achieve final concentrations ranging from 10^{-7} to 10^{-4} M. Control wells were prepared by addition of culture medium. Wells containing culture medium without cells were used as blanks. The plates were incubated at 37° C in a 5% CO₂ incubator for 44 h. Upon completion of the incubation, MTT dye solution ($20 \ \mu$ L, 5 mg/mL) was added to each well. After 4 h of incubation, 2-propanol ($100 \ \mu$ L) was added to solubilize the MTT formazan. The OD was measured at a wavelength of 570 nm by a microplate spectrophotometer. The IC₅₀ value was determined from plots of% viability against the dose of complexes added.

ACKNOWLEDGEMENT

This work was supported by the National Science and Technology Ministry of China (No. 2007BAI34B05) and Zhejiang Provincial Natural Science Foundation of China (No. Y4090052).

REFERENCES

1. (a) Brian, H.M.; Kuijpers, G.S.; Soede, A.C.; Laverman, P.; Boerman, O.C.; Delft, F.L.; Rutjes, F.P. Preparation and evaluation of glycosylated arginine-glycine-aspartate (RGD) derivatives for integrin targeting. *Bioconjugate Chem.* **2007**, *18*, 1847–1854; (b) Wilkinson, B.L.; Long, H.; Sim, E.; Fairbanks, A.J. Synthesis of Arabino glucosyl triazoles as potential inhibitors of mycobacterial cell wall biosynthesis. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 6265–6267.

2. Wilkinson, B.A.; Innocenti, A.; Vullo, D.; Supuran, C.T.; Poulsen, S.A. Inhibition of carbonic anhydrases with glycosyltriazole benzene sulfonamides. *J. Med. Chem.* **2008**, *51*, 1945–1953.

3. (a) Huisgen, R. 1,3-Dipolar cycloadditions. Angew. Chem., Int. Ed. **1963**, 2, 565–598; (b) Huisgen, R. In 1,3-Dipolar Cycloaddition Chemistry; Padwa, A., Ed.; Wiley: New York, **1984**, 1–176.

4. (a) Moorhouse, A.D.; Santos, A.M.; Gunaratnam, M.; Moore, M.; Neidle, S.; Moses, J.E. Stabilization of G-quadruplex DNA by highly selective ligands via click chemistry. *J. Am. Chem. Soc.* **2006**, *128*, 15972–15973.

5. Rozkiewicz, D.I.; Janczewski, D.; Verboom, W.; Ravoo, B.J.; Reinhoudt, D.N. "Click" chemistry by microcontact printing. *Angew. Chem.*, *Int. Ed.* **2006**, *45*, 5292–5296.

6. Burley, G.A.; Gierlich, J.; Mofid, M.R.; Nir, H.; Tal, S.; Eichen, Y.; Carell, T. Directed DNA metallization. J. Am. Chem. Soc. **2006**, 128, 1398–1399.

7. Maschauer, S.; Einsiedel, J.; Haubner, R.; Hocke, C.; Ocker, M.; Hubner, H.; Kuwert, T.; Gmeiner, P.; Prante, O. Labeling and glycosylation of peptides using click chemistry: a general approach to ¹⁸F-glycopeptides as effective imaging probes for positron emission tomography. *Angew. Chem. Int. Ed.* **2009**, *48*, 1–5.

8. Bertozzi, C.R.; Kiessling, L.L. Chemical glycobiology. *Science* **2001**, *291*, 2357–2363.

9. Rudd, P.M.; Elliott, T.; Cresswell, P.; Wilson, I.A.; Dwek, R.A. Glycosylation and the immune system. *Science* **2001**, *291*, 2370–2375.

10. Giannis, A. The sialyl Lewis X group and its analogues as ligands for selectins: chemoenzymatic syntheses and biological functions. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 178–180.

11. Feizi, T. Oligosaccharides that mediate mammalian cell-cell adhesion. *Curr. Opin.* Struct. Biol. **1993**, *3*, 701–710.

12. Zhao, Q.; Shen, C.; Zheng, H.; Zhang, J.C.; Zhang, P.F. Synthesis, characterization, and cytotoxicity of some novel glycosyl thiazol-2-imines as antitumoral agents. *Carbohydr. Res.* **2010**, *345*, 437–441.

13. Zhou, G.B.; Guan, Y.Q.; Shen, C.; Wang, Q.; Liu, X.M.; Zhang, P.F.; Li, L.L. A novel and convenient synthesis of thiazol-2(3*H*)-imine-linked glycoconjugates. *Synthesis* **2008**, *13*, 1994–1996.

14. Kolb, H.C.; Finn, M.G.; Sharpless, K.B. Click chemistry: diverse chemical function from a few good reactions. *Angew. Chem. Int. Ed.* **2001**, *40*, 2004–2021.

15. Wilkinson, B.L.; Bornaghi, L.F.; Poulsen, S.A.; Houston, T.A. Synthetic utility of glycosyl triazoles in carbohydrate chemistry. *Tetrahedron* **2006**, *62*, 8115–8125.

16. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods.* **1983**, *65*, 55–63.