

Identification of Cyclic Depsipeptides and Their Dedicated Synthetase from *Hapsidospora irregularis*

Shuwei Zhang,[†] Yixing Qiu,^{†,‡} Thomas B. Kakule,[§] Zhenyu Lu,[⊥] Fuchao Xu,[†] John G. Lamb,[⊥] Christopher A. Reilly,[⊥] Yong Zheng,^{||} Shing Wo Simon Sham,[#] Wei Wang,^{*,†,‡,||} Lijiang Xuan,^{||} Eric W. Schmidt,[§] and Jixun Zhan^{*,†,‡,||}

[†]Department of Biological Engineering, Utah State University, 4105 Old Main Hill, Logan, Utah 84322, United States


[‡]TCM and Ethnomedicine Innovation & Development Laboratory, School of Pharmacy, Hunan University of Chinese Medicine, Changsha, Hunan 410208, People's Republic of China

[§]Department of Medicinal Chemistry, L. S. Skaggs Pharmacy Institute, and [⊥]Department of Pharmacology and Toxicology, University of Utah, Salt Lake City, Utah 84112, United States

^{||}State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 501 Haik Road, Shanghai 202203, People's Republic of China

[#]Department of Chemistry and Biochemistry, Utah State University, 0300 Old Main Hill, Logan, Utah 84322, United States

Supporting Information




ABSTRACT: Seven cyclic depsipeptides were isolated from *Hapsidospora irregularis* and structurally characterized as the calcium channel blocker leucalacin and six new analogues based on the NMR and HRESIMS data. These new compounds were named leucalacins B–G. The absolute configurations of the amino acids and 2-hydroxyisocaproic acids were determined by recording the optical rotation values. Biological studies showed that calcium influx elicited by leucalacin F in primary human lobar bronchial epithelial cells involves the TRPA1 channel. Through genome sequencing and targeted gene disruption, a noniterative nonribosomal peptide synthetase was found to be involved in the biosynthesis of leucalacin. A comparison of the structures of leucalacin and its analogues indicated that the A₂ and A₄ domains of the leucalacin synthetase are substrate specific, while A₁, A₃, and A₅ can accept alternative precursors to yield new molecules.

Cyclic depsipeptides are a group of natural products that have shown a wide range of biological activities and show immunosuppressant,^{1,2} antibiotic,^{3,4} anti-inflammatory,^{5,6} and antitumor^{7–9} properties. They are assembled from proteinogenic/nonproteinogenic amino acids and hydroxy acids through amide and ester linkages. Fungi are an important source of biologically active molecules. Leucalacin (1) was previously isolated from the fungus *Hapsidospora irregularis* FERM BP-2511 as a new cyclic depsipeptide and potent calcium channel blocker, which inhibits the binding of *H*-nitrendipine to porcine heart membranes *in vitro* and lowers the blood pressure of spontaneous hypertensive rats.^{10,11} Calcium channel blockers block the entry of calcium into cells. These agents are used to treat hypertension, angina pectoris, myocardial infarction, and abnormal heart rhythms.^{12,13}

Current clinic calcium channel blockers are mainly divided into three groups: phenylalkylamines (e.g., verapamil), benzothiazepines (e.g., diltiazem), and dihydropyridines (e.g., cilnidipine). Compound 1 represents a new type of calcium channel blocker. It is of significant interest to find more analogues of 1 for bioactivity screening. To this end, we reinvestigated the chemical constituents of *H. irregularis*. Herein we report the isolation and structure elucidation of six new depsipeptides from this fungus. The absolute configurations of amino and α -hydroxy acids were determined by hydrolysis of the parent depsipeptides and measurement of the optical rotation values of the isolated moieties. The leucalacin synthetase gene was discovered by genome sequencing. Its

Received: September 4, 2016



involvement in leualacin biosynthesis was confirmed by double crossover gene disruption through *Agrobacterium*-mediated transformation. Production of the six leualacin analogues revealed that this giant fungal noniterative peptide synthetase has relaxed substrate specificity, which presents great potential for precursor-directed biosynthesis of new cyclic depsipeptides.

RESULTS AND DISCUSSION

H. irregularis FERM BP-2511 was grown in K-2 broth at 30 °C for 8 days, and HPLC analysis revealed the production of 1–7, in addition to the previously characterized cephalosporin P1, isocephalosporin P1, Sch210971, and Sch210972 (Figure S1).^{14,15} The strain was cultured in K2 broth with and without 0.5% tyrosine (3 L each), from which 1 and six new cyclic depsipeptides were isolated (2–7).

The molecular formula of 1 was determined to be $C_{31}H_{47}N_3O_7$ based on HRESIMS analysis, which is the same as that of leualacin. It was further confirmed to be leualacin by a comparison of its NMR data (Figures S2 and S3; Tables 1 and

2) with those previously reported.¹¹ Compound 2 was obtained as an off-white powder. The molecular formula was elucidated as $C_{32}H_{49}N_3O_7$ based on HRESIMS analysis as well as its 1H and ^{13}C NMR data (Figures S4 and S5; Tables 1 and 2), which indicated that 2 has an additional CH_2 group compared to 1. By comparing the 1H NMR data of 2 with those of 1, one CH at δ_H 2.64–2.69 (1H, m) in 2 replaced the CH_2 at 2.54–2.61 (2H, m) in 1. Compound 2 has an additional methyl group. Analysis of its HSQC, HMBC, and 1H – 1H COSY spectra (Figure 1 and Figures S6–S8) revealed that 2 contains one unit of *N*-methyl-L-phenylalanine (*N*-Me-L-Phe), two units of 2-hydroxyisocaproic acid (2-HICA or leucic acid), and one unit of leucine (Leu) that are also present in 1. However, the β -alanine (β -Ala) moiety in 1 was replaced with 3-aminoisobutyric acid (3-AIB) based on the specific COSY signals of the spin system of $NH-CH_2-CH-CH_3$ and the HMBC correlations of CH_3 at δ_H 1.21 (3H, d, 7.1) and CH at δ_H 2.64–2.69 (1H, m) to the carbonyl group at δ_C 174.9. The connection order of the five moieties was determined by HMBC correlations of α -CH of the *N*-Me-L-Phe moiety at δ_H 4.63 (1H, dd, 2.5, 10.5), β - CH_2 at δ_H 3.60–3.71 (2H, m), and NH at δ_H 7.45 (1H, dd, 5.0, 5.5) of the 3-AIB unit to the carbonyl carbon of the *N*-Me-L-Phe moiety at δ_C 170.0, α -CH of the 3-AIB moiety at δ_H 2.64–2.69 (1H, m) and α -CH of the R-HICA moiety at δ_H 5.12 (1H, dd, 5.0, 7.6) to the carbonyl group of the 3-AIB moiety at δ_C 174.9, α -CH of R-HICA, α -CH at δ_H 4.50 (1H, ddd, 3.0, 9.7, 10.8) and NH at δ_H 6.32 (1H, d, 9.4) of the Leu moiety to the carbonyl group of the R-HICA unit at δ_C 171.0, α -CH of the Leu moiety and α -CH of the S-HICA moiety at δ_H 4.73 (1H, dd, 2.1, 10.5) to the carbonyl group of the Leu moiety at δ_C 174.1, and α -CH of the S-HICA moiety, α -CH of the *N*-Me-L-Phe moiety, and NCH_3 of the *N*-Me-L-Phe moiety at δ_H 2.87 (3H, s) to the carbonyl group of the S-HICA moiety at δ_C 170.1. Thus, 2 was identified as a new compound that is identical to 1 except for a 3-AIB moiety that replaces β -Ala in 1. It was named leualacin B.

Compounds 3 and 4 were obtained as an off-white powder. HRESIMS analysis revealed that 3 and 4 have the same molecular formula of $C_{30}H_{45}N_3O_7$. Thus, they have one CH_2 less than 1. A comparison of the NMR data of 3 (Figures S9 and S10) and 1 and analysis of the 2D NMR spectra (Figures S11–S13) revealed that 3 does not contain a methyl group at the amino group of the L-Phe moiety. Instead, an NH signal was found at δ_H 6.54 (1H, d, 8.6), which was confirmed by its 1H – 1H COSY correlation to the α -CH proton of the L-Phe moiety at δ_H 4.69–4.77 (1H, m) as well as its HMBC correlation to the carbonyl group of the S-HICA moiety at δ_C 170.1. Further analysis of the HSQC, HMBC, and 1H – 1H COSY spectra (Figure 1) indicated that 3 is also a cyclic depsipeptide composed of five units, including β -Ala, R-HICA, L-Leu, and S-HICA and L-Phe. Unlike 3, compound 4 was found to have the methyl group at the amino group of the L-Phe moiety (Figures S14–S18). Except for the leucine unit, the 1H and ^{13}C NMR data of the other four units in compound 4 were nearly identical to those of compound 1. Furthermore, the 1H – 1H COSY spectrum revealed a spin system of $NH-CH-CH-(CH_3)_2$. The HMBC correlations of the CH at δ_H 4.43 (1H, d, 4.1) and CH at δ_H 2.35 (1H, qdd, 4.1, 6.5, 6.9) of this spin system to the carbonyl group at δ_C 174.1 showed that the L-Leu moiety in 1 was replaced with valine (Val) in this compound. The connection of the five units was confirmed by

Table 1. ¹H NMR Data of Compounds 1–7 (1–6 Recorded in CDCl₃, 7 Recorded in CD₃OD, δ_H in ppm, J in Hz)

position	1	2	3	4	5	6	7
L-Phenylalanine				L-Tyrosine			
α	4.63–4.67, m	4.63, dd (2.5, 10.5)	4.69–4.77, m	4.65, dd (3.4, 11.4)	4.61, dd (3.0, 11.0)	4.61–4.67, m	4.52, dd (6.2, 8.3)
α-NH			6.54, d (8.6)			6.8, d (8.3)	N/A
N-CH ₃	2.87, s	2.87, s		2.87, s	2.87, s		
β	2.94 dd (10.5, 14.4)	2.95, dd (10.9, 12.4)	3.12–3.23, m	2.94, dd (11.7, 14.4)	2.88, dd (11.5, 14.5)	3.05–3.10, m	2.99 dd (8.6, 14.1)
	3.38, dd (2.7, 14.3)	3.30, dd (2.52, 12.2)		3.41, dd (3.4, 14.1)	3.31, dd (3.0, 14.5)		3.12, dd (6.2, 14.1)
o	7.10, d (7.5)	7.07, d (6.9)	7.17, d (7.1)	7.09, d (6.9)	6.91, d (8.5)	6.96, d (7.9)	6.99, d (8.6)
m	7.26–7.31, m	7.23–7.29, m	7.23–7.32, m	7.22–7.31, m	6.75, d (8.5)	6.74, d (7.9)	6.99, d (8.6)
p	7.23–7.25, m	7.23, t (6.9)	7.23, t (6.9)	7.29, t (7.6)			
β-Alanine		R-3-Aminoisobutyric acid (R-3-AIB)	β-Alanine				
α	2.54–2.61, m	2.64–2.69, m	2.53–2.67, m	2.57, dd (3.8, 8.3)	2.54–2.57, m	2.49–2.52, m	2.44–2.59, m
α-CH ₃		1.21, d (7.1)					
β	3.31–3.39, m	3.60–3.71, m	3.82–3.94, m	4.04, dt (3.8, 13.7)	3.97–4.10, m	3.80–3.84, m	3.73–3.80, m
	3.98–4.09, m		3.25–3.34, m	3.33–3.42, m	3.34–3.42, m	3.21–3.25, m	3.22–3.30, m
β-NH	7.45–7.48, m	7.45, dd (5.0, 5.5)	7.10–7.15, m	7.44–7.49, m	7.49, d (6.8)	7.55–7.57, m	N/A
(R)-2-hydroxyisocaproic acid (R-HICA)							
α	5.13–5.18, m	5.12, dd (5.0, 7.6)	5.19, dd (3.8, 9.9)	5.21, dd (6.2, 7.9)	5.17, dd (6.4, 7.6)	5.17–5.21, m	5.17–5.24, m
β	1.68–1.85, m	1.68–1.88, m	1.66–1.90, m	1.72–1.87, m	1.71–1.85, m	1.62–1.83, m	1.61–1.78, m
γ	1.59–1.71, m	1.58–1.73, m	1.60–1.77, m	1.59–1.70, m	1.60–1.69, m	1.54–1.68, m	1.64–1.74, m
L-Leucine				L-Valine	L-Leucine	L-Valine	
α	4.48–4.56, m	4.50, ddd (3.0, 9.7, 10.8)	4.08–4.18, m	4.43, d (4.1)	4.48–4.55, m	4.13–4.17, m	3.90, d (8.9)
α-NH	6.16, d (9.8)	6.32, d (9.4)	6.89, d (6.6)	6.41, d (9.6)	6.32, d (9.6)	7.41, d (6.6)	N/A
β	1.57–1.72, m	1.48–1.73, m	1.54–1.73, m	2.35, qqd (4.1, 6.5, 6.9)	1.50–1.81, m	1.50–1.70, m	2.08–2.16, m
γ	1.65–1.86, m	1.61–1.76, m	1.60–1.77, m		1.64–1.82, m	1.54–1.65, m	
(S)-2-Hydroxyisocaproic acid (S-HICA)							
α	4.65–4.70, m	4.73, dd (2.1, 10.5)	5.07, dd (3.6, 6.1)	4.66, dd (2.3, 11.3)	4.72, dd (1.8, 11.0)	5.15–5.17, m	5.17–5.24, m
β	1.35–1.59, m	1.36–1.48, m	1.35–1.54, m	1.37–1.52, m	1.41–1.50, m	1.45–1.57, m	1.29–1.54, m
	–0.38 - -0.47, m	–0.32 - -0.40, m		–0.38 to -0.47, m	–0.21 to -0.27, m		
γ	1.35–1.59, m	1.36–1.48, m	1.33–1.52, m	1.23–1.40, m	1.43–1.60, m	1.30–1.43, m	1.43–1.49, m
R/S-2-Hydroxyisocaproic acid (R/S-HICA) and δ-H of L-leucine (L-Leu)/γ-H of L-valine (L-Val)							
	R-HICA	R-HICA/L-Leu	S-HICA	R-HICA	R-HICA/L-Leu	S-HICA	R-HICA
	0.56, d (6.5)	0.92, d (5.5)	0.84, d (5.9)	0.91, d (6.5)		0.91, d (5.9)	0.95, d (6.5)
	0.67, d (6.5)	0.87, d (5.9)	0.86, d (6.8)	0.95, d (6.1)	0.90, d (6.2)	0.92, d (6.2)	0.99, d (6.2)
	S-HICA/	S-HICA	R-HICA/	L-Val	0.9, d (6.4)	L-Leu	L-Val
	L-Leu	0.66, d (6.2)	L-Leu	0.91, d (6.5)	S-HICA	0.91, d (5.9)	0.97, d (6.5)
	0.89, d (6.5)	0.56, d (6.4)	0.92, d (6.2)	1.01, d (6.9)	0.62, d (6.4)	0.97, d (5.5)	1.00, d (6.5)
	0.93, d (6.5)		0.93, d (6.2)	S-HICA	0.73, d (6.2)	R-HICA	S-HICA
			0.95, d (6.5)	0.56, d (6.5)		0.83, d (5.8)	0.86, d (5.8)
			1.00, d (5.9)	0.67, d (6.5)		0.87, d (5.2)	0.89, d (5.8)

its HMBC spectrum (Figure 1). Both 3 and 4 are new compounds and were named leualacins C and D, respectively.

The molecular formula of 5 was determined to be C₃₁H₄₇N₃O₈ according to the HRESIMS data, suggesting that it has one more oxygen atom than 1. The NMR spectra of 5 (Figures S19–S23) were similar to those of 1, except the difference in the N-Me-L-Phe moiety. The aromatic proton signals at δ_H 6.91 (2H, d, 8.5) and 6.75 (2H, d, 8.5) in the ¹H NMR spectrum and carbon signals at δ_C 127.6 (C), 130.4 (2C, CH), 115.9 (2C, CH), and 155.7 (C) in the ¹³C NMR spectrum (Tables 1 and 2) indicated that there is a 1,4-disubstituted phenyl ring, with a hydroxy as one of the substituents. The HMBC correlations (Figure 1) revealed the presence of an N-Me-L-Tyr moiety. According to the 1D and

2D NMR spectra, 5 was characterized as a new leualacin analogue that has the same structure as 1 except that the N-Me-L-Phe was replaced with N-methyl-L-tyrosine (N-Me-L-Tyr). This compound was named leualacin E. Compound 6 also has eight oxygen atoms in its molecular formula of C₃₀H₄₅N₃O₈ based on the HRESIMS, but it has one less CH₂ than 5. The NMR spectra (Figures S24–S28) also showed that it has a Tyr unit, and the detailed 2D NMR signals confirmed it contains the same units as those in 5 with the same connection order, except that there is no methyl group at the amino group of the L-Tyr moiety. This was confirmed by the ¹H–¹H COSY correlation (Figure 1) between NH at δ_H 6.83 (1H, d, 8.3) and the α-CH of L-Tyr at δ_H 4.61–4.67 (1H, m). 6 was named leualacin F.

Table 2. ¹³C NMR Data of Compounds 1–7 (1–6 Recorded in CDCl₃, 7 Recorded in CD₃OD, δ_C in ppm)

position	1	2	3	4	5	6	7
L-Phenylalanine				L-Tyrosine			
C=O	173.5 (s)	170.0 (s)	171.4 (s)	168.5 (s)	168.8 (s)	172.0 (s)	173.6 (s)
α	63.9 (d)	63.3 (d)	54.4 (d)	63.8 (d)	64.0 (d)	54.8 (d)	56.5 (d)
N-CH ₃	30.3 (q)	30.2 (q)		30.2 (q)	30.3 (q)		
β	34.4 (t)	34.4 (t)	37.8 (t)	34.3 (t)	33.3 (t)	37.2 (t)	37.6 (t)
γ	138.1 (s)	137.5 (s)	136.9 (s)	138.0 (s)	128.7 (s)	127.6 (s)	128.8 (s)
o	129.8 (2C, d)	129.7 (2C, d)	129.5 (2C, d)	129.7 (2C, d)	130.7 (2C, d)	130.4 (2C, d)	131.5 (2C, d)
m	129.5 (2C, d)	129.4 (2C, d)	128.9 (2C, d)	129.4 (2C, d)	116.4 (2C, d)	115.9 (2C, d)	116.4 (2C, d)
p	127.4 (d)	127.3 (d)	127.1 (d)	127.3 (d)	155.9 (s)	155.7 (s)	157.6 (s)
β-Alanine		R-3-Aminoisobutyric acid (R-3-AIB)	β-Alanine				
C=O	174.8 (s)	174.9 (s)	171.8 (s)	173.5 (s)	173.2 (s)	171.8 (s)	172.2 (s)
α	34.7 (t)	40.4 (d)	34.8 (t)	34.8 (t)	34.5 (t)	34.8 (t)	35.6 (t)
α-CH ₃		13.6 (q)					
β	34.9 (t)	41.4 (t)	36.1 (t)	34.3 (t)	34.7 (t)	36.1 (t)	37.1 (t)
(R)-2-Hydroxyisocaproic acid (R-HICA)							
C=O	170.6 (s)	171.0 (s)	171.0 (s)	170.7 (s)	171.0 (s)	171.2 (s)	173.0 (s)
α	70.9 (d)	72.8 (d)	71.7 (d)	72.9 (d)	72.9 (d)	71.9 (d)	73.9 (d)
β	37.9 (t)	37.8 (t)	39.0 (t)	37.4 (t)	37.5 (t)	39.5 (t)	41.3 (t)
γ	24.4 (d)	24.4 (d)	24.6 (d)	24.3 (d)	24.3 (d)	24.6 (d)	25.9 (d)
L-Leucine				L-Valine	L-Leucine	L-Valine	
C=O	171.6 (s)	174.1 (s)	171.2 (s)	174.1 (s)	174.6 (s)	170.6 (s)	171.0 (s)
α	51.5 (d)	51.6 (d)	53.2 (d)	57.8 (d)	51.6 (d)	53.2 (d)	61.9 (d)
β	38.9 (t)	38.5 (t)	39.2 (t)	29.0 (d)	38.6 (t)	39.1 (t)	30.5 (d)
γ	25.4 (d)	25.2 (d)	25.2 (d)		25.2 (d)	25.1 (d)	
(S)-2-Hydroxyisocaproic acid (S-HICA)							
C=O	168.6 (s)	171.7 (s)	170.1 (s)	171.2 (s)	172.1 (s)	170.8 (s)	173.3 (s)
α	72.9 (d)	70.6 (d)	74.5 (d)	70.8 (d)	71.0 (d)	74.1 (d)	74.3 (d)
β	37.6 (t)	37.9 (t)	40.4 (t)	37.9 (t)	37.7 (t)	40.6 (t)	42.4 (t)
γ	24.2 (d)	24.2 (d)	24.7 (d)	24.4 (d)	24.4 (d)	24.6 (d)	25.6 (d)
δ-C of L-Leucine and R/S-HICA, γ-C of L-valine							
	R-HICA	R-HICA	S-HICA	R-HICA	R-HICA	R-HICA	R-HICA
	20.9 (q)	22.2/22.9 (q)	21.7 (q)	22.4/23.0 (q)	22.3/23.0 (q)	21.7 (q)	22.7/23.5 (q)
	23.1(q)	L-Leu	23.1 (q)	L-Val	L-Leu	22.9 (q)	L-Val
	S-HICA/L-Leu	20.8/22.9 (q)	R-HICA/L-Leu	17.2/19.6 (q)	20.8/22.8 (q)	S-HICA/L-Leu	20.2/19.6 (q)
	23.2 (q)	S-HICA	23.3 (q)	S-HICA	S-HICA	23.1 (2C, q)	S-HICA
	23.1 (q)		23.2 (q)			21.7 (q)	
	22.5 (q)		21.8 (2C, q)				
	21.3 (q)						

The molecular formula of **7** was determined to be C₂₉H₄₃N₃O₈ by HRESIMS analysis, which is one CH₂ unit less than **6**. Analysis of the 2D NMR spectra of **7** (Figure 1 and Figures S29–S33; Tables 1 and 2) revealed that it contains the N-Me-L-Tyr, β-Ala, R-HICA, and S-HICA moieties that are also present in **6**. However, **7** has a Val moiety in the same position as the L-Leu moiety in **6**. It was named leualacin G.

The absolute configuration of the major product **1** has been previously determined¹¹ to consist of β-Ala, R-HICA, L-Leu, S-HICA, and N-Me-L-Phe. **2**–**7** are new compounds that have one or two new moieties that substitute the β-Ala, L-Leu, or N-Me-L-Phe moiety in **1**. In order to determine the absolute configurations of these new moieties, we hydrolyzed the representative compounds **2**, **4**, and **6** at the ester bonds using 1 N NaOH (Supporting Information) to get two fragments and analyzed the products by LC-MS. The two fragments were separated by HPLC and then treated with 6 N HCl to hydrolyze the amide bonds, from which individual units of these compounds were obtained (Figure S34). The optical rotations of these units were recorded (Table S1). Their

absolute configurations were determined by comparing the recorded optical rotations with those of standards or reported. The results showed the moieties in **2**–**7** are L-Phe¹¹ or L-Tyr,¹⁶ β-Ala or R-3-AIB,¹⁷ R-HICA, S-HICA,^{11,18} and L-Leu¹⁹ or L-Val.²⁰

Compound **1** blocks *H*-nitrendipine binding to porcine heart membranes *in vitro* and can lower the blood pressure of spontaneous hypertensive rats.⁵ Calcium is a versatile and critical second messenger in all cells. Stimulation and inhibition of changes in intracellular calcium content caused by **1**–**7** were assessed in lobar bronchial epithelial cells. TRP ion channels in lung epithelial cells are involved in innate immunity and various other processes. Calcium influx was elicited by **3** and **6** at 100 μM, which was correlated with activation of the human TRPA1 and to a lesser extent M8 channels, but not TRPV1, -V3, or -V4 (Table S2). Activation of TRPA1 is associated with pulmonary irritation, cough, and edema. Elsewhere, TRPA1 signals pain and is associated with allergic sensitization including asthma and dermatitis.^{21–23}




Figure 1. Selected ^1H – ^1H COSY and HMBC correlations for 2–7.




Figure 2. Proposed biosynthesis of 1 by a noniterative NRPS.

Compounds 1–7 belong to a large group of structurally diverse natural products, nonribosomal peptides (NRPs). They are synthesized by nonribosomal peptide synthetases (NRPSs). These synthetases are modular enzymes that consist of different catalytic domains, such as condensation (C), adenylation (A), thiolation (T), methylation (MT), epimerization (E), and oxidation (Ox). To understand how 1–7 are synthesized, the genome of *H. irregularis* was sequenced. Because the structure of 1 contains different amino acid/hydroxy acid units, it is proposed that this molecule is assembled by a noniterative NRPS that is often present in bacteria. Annotation of the sequenced genome revealed the presence of a large noniterative NRPS (Figure 2, GenBank accession number KU994894) that consists of five A domains, five T domains, and five C domains. Additionally, there is an E domain sitting between the T_2 and C_3 domains. An N-MT domain is embedded in the A_5 domain. The domain organization of this 6521-aa enzyme (717 kDa, named leualacin synthetase, or LACS) is colinear with the structure of 1, and thus it represents a possible NRPS involved

in leualacin biosynthesis. Although the signature sequences of the five A domains (Table S3) could be extracted from the amino acid sequence of LACS using NRPPSPredictor2²⁴ based on the reported bacterial NRPSs, it is hard to conclusively predict the substrate of each A domain due to the lack of information about genuine signature sequences in fungal NRPSs. In addition, there are several novel precursors used in the biosynthetic process of 1–7, such as HICA, which has not been reported as a direct precursor for NRP biosynthesis. Although cryptophycins also have an HICA unit, it was reported that the A_4 domain of the dedicated PKS/NRPS selects α -ketoisocaproic acid, which is reduced by a ketoreductase domain in the same module to yield the HICA unit.²⁵ We predicted the substrate of each A domain based on the domain organization of LACS and the structure of 1. A_1 – A_5 were proposed to select and activate β -Ala, S-HICA, L-Leu, S-HICA, and L-Phe. The E domain in the module converts S-HICA to R-HICA, while the MT domain in module 5 methylates the NH_2 group of the L-Phe moiety. The C-terminal




Figure 3. Disruption of *lacs* in *H. irregularis*. (A) Strategy for targeted disruption of *lacs*. (B) Verification of the correct mutant by PCR. (C) HPLC analysis of the products of the wild type and mutant on the oatmeal agar plates at 210 nm. (i) Wild type; (ii) mutant; (iii) standard of 1; (iv) standard of 2.

condensation-like (C_T) domain catalyzes the cyclization and concomitant release of the final products.

To find out whether this NRPS is involved in the biosynthesis of **1**, we conducted a gene disruption experiment. Two fragments (3.5 and 4.0 kb) of *lacs* were amplified from the genome of *H. irregularis* and ligated into the binary donor vector pAG1-H3 as the left and right arms, respectively. The resulting plasmid pAG1-H3-KO (Figure 3A) was introduced into *H. irregularis* through *Agrobacterium tumefaciens*-mediated transformation. Correct transformants were then subjected to PCR screening using the genome- and vector-specific primers A–D (Figure 3A). As shown in Figure 3B, a 4534-bp fragment was amplified from the correct double crossover mutant with primers A and B and a 5026-bp fragment with primers C and D. Because primers B and C were specifically from the hygromycin resistance gene (*HygR*) in the vector, no PCR products were obtained from the wild type. This correct mutant was then grown on the oatmeal agar modified from WSH agar⁵ for product analysis. As shown in Figure 3C, when grown in the oatmeal medium, the wild type produces **1** and **2** as the major metabolites (traces i–iii). In contrast, the mutant failed to produce **1** or any analogues (trace iv), indicating the essential role of LACS in leuolacin biosynthesis.

H. irregularis is a fungus of the family Pseudeurotiaceae.²⁶ In this work, six new cyclic depsipeptides were characterized, which significantly expanded the spectrum of natural products from this fungus. These structures consist of two HICA units with different configurations and three amino acid moieties. The main difference among the seven cyclic depsipeptides lies in the amino acid units, while the two HICA units remain the same. This indicated that A_2 and A_4 of LACS, which activate S-HICA, are more substrate specific. In contrast, A_1 , A_3 , and A_5 are more flexible. The structures of **5**–**7** showed that A_5 can use L-Tyr instead of L-Phe in **1**. Similarly, A_1 was found to take 3-AIB to yield **2**, while the discovery of **4** and **7** indicated that A_3 activates both L-Val and L-Leu. Thus, LACS is a highly flexible NRPS that synthesizes a variety of leuolacin analogues in the presence of different precursors.

EXPERIMENTAL SECTION

General Experimental Procedures. 1D NMR spectra were recorded in $CDCl_3$ or CD_3OD at 25 °C on a JEOL ECX-300 instrument (300 MHz for 1H NMR and 75 MHz for ^{13}C NMR). 2D NMR spectra were collected on a JEOL ECX-300 or a Bruker AvanceIII HD Ascend-500 (HMBC spectrum of compound **7**) instrument. The chemical shift (δ) values are given in parts per million (ppm). The coupling constants (J values) are reported in hertz (Hz). LRESIMS were acquired on an Agilent 6130 LC-MS, and

HRESIMS were measured on an Agilent G6224A TOF mass spectrometer.

Strains, Media, and Culture Conditions. *Hapsidospora irregularis* FERM BP-2511 was purchased from the International Patent Organism Depository, Japan. It was routinely grown in K2 agar medium consisting of potato dextrose agar (3.9%, w/v), sucrose (2%, w/v), casamino acid (1%, w/v), KH_2PO_4 (0.5%, w/v), and $MgSO_4$ (0.125%, w/v) at 30 °C. *E. coli* XL-1 Blue (Stratagene) was used for routine cloning and plasmid propagation. It was routinely grown at 37 °C on LB agar plates or in liquid LB medium supplemented with appropriate antibiotics (ampicillin, 50 μ g/mL; kanamycin, 50 μ g/mL). *Agrobacterium tumefaciens* LBA4404 was used for the introduction of the disruption plasmid into *H. irregularis* FERM BP-2511.

Production and Purification of Cyclic Depsipeptides. Single colonies were obtained by streaking the fungus on K2 agar medium at 30 °C for 8 days. A single colony was picked and precultured in 4 mL of K2 broth, then cultured in 50 mL of K2 broth consisting of potato dextrose broth (2.4%, w/v), sucrose (2%, w/v), casamino acid (1%, w/v), KH_2PO_4 (0.5%, w/v), and $MgSO_4$ (0.125%, w/v) at 250 rpm and 30 °C for 7 days. The culture was centrifuged at 3500 rpm for 5 min. The supernatant was extracted with 50 mL of ethyl acetate three times, and the cells were extracted three times with 50 mL of methanol. The ethyl acetate and methanol extracts were combined and dried under reduced pressure. The residue was redissolved in 1 mL of methanol for analysis on an Agilent 1200 HPLC (HPLC conditions are shown in Figure S1).

The strain was also grown in 3 L of K2 broth and 3 L of K2 plus 0.5% L-tyrosine (to enhance the production of **5**–**7**) for the isolation of **1**–**7**. After 8 days, the cultures were combined and centrifuged at 3500 rpm for 5 min. The supernatant was combined and loaded to an HP-20 column (10 cm \times 50 cm), and the column was eluted with 4 L of water. The cells were extracted with 500 mL of methanol three times with sonication for 40 min. After centrifugation at 3500 rpm for 5 min, the methanol extract was diluted with 3 volumes of water and the solution was loaded to the HP-20 column as the 25% eluant. The column was then successively eluted with 4 L of 50% aqueous methanol, 75% aqueous methanol, and 100% methanol. These fractions were analyzed by HPLC, and the target compounds were found in the 75% and 100% methanol fractions. These two fractions were separated on the Agilent 1220 HPLC with an Agilent ZORBAX SB-C₁₈ column (5 μ m, 21.2 mm \times 150 mm), eluted with a gradient of methanol–water (0–15 min, 70%; 15–18 min, 70–80%; 18–25 min, 80%; 25–27 min, 80–90%; 27–35 min, 90%, v/v) with 0.1% formic acid at a flow rate of 3 mL min^{−1} to obtain six fractions, A–F. LC-MS analysis indicated that fraction A contained **7**, fraction B contained **5** and **6**, fraction D contained **3** and **4**, fraction E contained **1**, and fraction F contained **1** and **2**. Then fractions A–F were repurified by HPLC with an Agilent ZORBAX SB-C₁₈ column (5 μ m, 4.6 mm \times 250 mm) at 1 mL min^{−1} and detected at 210 nm. Fraction A was eluted with 40% acetonitrile–water to get pure **7** (18 mg). Fraction B was eluted with 55% acetonitrile–water to get pure **5** (33 mg) and **6** (59 mg). Fraction D was eluted with 60% acetonitrile–water to get pure **3** (35 mg) and **4** (35 mg). Fraction E was eluted with 70%

methanol–water to get pure **1** (450 mg). Fraction F was eluted with 85% methanol–water to get pure **2** (39 mg). The purified compounds were subjected to spectral analyses for structure determination (Tables 1 and 2).

Compound 1: off-white powder; $[\alpha]_D^{19}$ –97.8 (*c* 1.0, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 574.3493 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{31}\text{H}_{48}\text{N}_3\text{O}_7$, 574.3487).

Compound 2: off-white powder; $[\alpha]_D^{19}$ –99.0 (*c* 0.3, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 588.3648 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{32}\text{H}_{50}\text{N}_3\text{O}_7$, 588.3643).

Compound 3: off-white powder; $[\alpha]_D^{19}$ –48.2 (*c* 1.17, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 560.3332 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{46}\text{N}_3\text{O}_7$, 560.3330).

Compound 4: off-white powder; $[\alpha]_D^{19}$ –68.0 (*c* 0.25, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 560.3333 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{46}\text{N}_3\text{O}_7$, 560.3330).

Compound 5: off-white powder; $[\alpha]_D^{19}$ –96.1 (*c* 0.62, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 590.3440 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{31}\text{H}_{48}\text{N}_3\text{O}_8$, 590.3436).

Compound 6: off-white powder; $[\alpha]_D^{19}$ –46.0 (*c* 0.50, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 576.3284 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{30}\text{H}_{46}\text{N}_3\text{O}_8$, 576.3279).

Compound 7: off-white powder; $[\alpha]_D^{19}$ –52.9 (*c* 0.34, MeOH); ^1H and ^{13}C NMR data, see Tables 1 and 2; HRESIMS m/z 562.3119 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{29}\text{H}_{44}\text{N}_3\text{O}_8$, 562.3123).

Determination of Absolute Configurations of the Five Units in 1–7 (Table S1 and Figure S34). A 25 mg amount of **6** was dissolved in 600 μL of 30% aqueous acetonitrile, and 100 μL of 1 N NaOH in water was added under stirring. The compound was hydrolyzed at 40 °C for 1 h. The reaction mixture was neutralized with 1 N HCl. The hydrolysis was confirmed by LC-MS, and two fragments (with a MW of 366 and 245, respectively) were purified on an Agilent 1220 HPLC with an Agilent ZORBAX SB-C₁₈ (5 μm , 4.6 mm \times 250 mm) at 1 mL min^{–1}, eluted with 35% acetonitrile–water and monitored at 210 nm. The two fragments were further hydrolyzed with 6 N HCl at 115 °C for 12 h. After neutralizing with NaOH, the products were analyzed by LC-MS and compared with the standards. Then the target products were purified on the same HPLC eluted with a gradient of acetonitrile–water from 0% to 100% in 30 min. The optical rotations of the purified amino acids and hydroxy acids were recorded on a Rudolph Autopol IV polarimeter with a 10 cm cell at 19 °C. Compounds **2** and **4** were hydrolyzed and purified according to the same procedure for **6**. The absolute configurations were determined by comparing the optical data with those reported in the literature. The optical rotations of compounds **1**–**7** were also recorded on the same equipment (Table S1).

Transient Receptor Potential Channel Assay. Fluorometric cell-based Ca²⁺ flux assays were performed using a BMG Labtech NOVOSTar fluorescence plate reader equipped with a plate-to-plate reagent delivery system. Human embryonic kidney-293 (HEK-293) cells that stably overexpress human TRPA1, M8, TRPV3, or V4 have been previously described²⁷ and were grown to confluence in 2% (w/v) gelatin-coated 96-well plates in DMEM [F12 media supplemented with 5% fetal bovine serum, 1 \times penicillin/streptomycin (Invitrogen), and geneticin (300 $\mu\text{g}/\text{mL}$)]. Assays using TRPV1 overexpressing and normal BEAS-2B cells have also been described.¹ Lobar bronchial epithelial cells (Lifeline Cell Technology, donor lot #1344) were grown to 80–90% confluence in BronchiaLife B/T medium (Lifeline Cell Technology).

Cells were prepared for assay by replacing the growth media with a 1:1 solution of LHC-9 and Fluo 4-Direct (Invitrogen) reagent containing Fluo 4-AM, pluronic F-127, probenecid, and a proprietary quencher dye. Cells were then incubated at 37 °C for 1 h in a cell culture incubator or at room temperature (BEAS-2B cells). Cells were subsequently washed by replacing the loading solution with LHC-9 containing 1 mM water-soluble probenecid (Invitrogen) and 750 μM Trypan Red (ATT Bioquest). All cell types were incubated for an additional 30 min at 37 °C to allow for Fluo 4-AM cleavage and activation as well as equilibration to both the test compounds and Fluo 4.

Treatment solutions were prepared in LHC9 (TRPV1, M8, and V4) or calcium assay buffer (TRPA1, V3, LOBAR) at 3 \times concentration, and 25 μL was added to 50 μL of media on the cells in 96-well plates. Changes in cellular fluorescence were monitored for 1 min at 37 °C. Data were quantified as the maximum rate of change in fluorescence intensity (max $\Delta F/s$) vs media-only treatment (negative control) and, in HEK-293 or normal BEAS-2B cells, were represented as the percentage of response relative to the specific TRP agonist control (TRPA1-2,4-di-*tert*-butylphenol at 250 μM , M8-icilin at 25 μM , V1-capsaicin at 10 μM , V3-carvacrol at 300 μM , and V4-GSK1016790A at 30 nM). A minimum of three replicates were used for all treatments.

Knockout of *lacs* in *H. irregularis* FERM BP-2511 through *Agrobacterium*-Mediated Transformation. *H. irregularis* FERM BP-2511 was cultured in 50 mL of K2 broth at 30 °C and 250 rpm for 6 days. The mycelia were collected by filtration and stored at –80 °C for 2 h. Then 200 mg of the mycelia was ground in liquid nitrogen. A 3 mL amount of the lysis buffer consisting of 0.5 M NaCl, 10 mM Tris-HCl (pH = 7.5), 10 mM EDTA, and 1% SDS was added into the mycelia at 65 °C. The extraction of genomic DNA was done with cetyltrimethylammonium bromide extraction buffer²⁸ followed by purification through phenol/chloroform extraction and precipitation with 2-propanol.²⁹ RNA was removed by adding RNase.

Two DNA fragments were amplified by PCR from the *lacs* gene using the genomic DNA of *H. irregularis* FERM BP-2511 as the template. A 3600-bp fragment upstream of the disruption region of the *lacs* gene was amplified using the primers BP2511-128-NRPS-L-AvrII-3 (5'-aaCCTAGGacagcctggttctgagaac-3', AvrII site in bold) and BP2511-128-NRPS-L-KpnI-5 (5'-aaGGTACCagagcctgtgactccagac-3', KpnI site in bold) and cloned into pJET1.2 for sequencing. After sequence validation, this left-arm gene fragment was excised with AvrII and KpnI and inserted into pAG1-H3 between the same sites. A 3976-bp fragment downstream of the disruption region of the *lacs* gene was amplified using the primers BP2511-128-NRPS-R-SbfI-3 (5'-aaCCTGCAGGgtgtcgtctttcgtcgtg-3', SbfI site in bold) and BP2511-128-NRPS-R-SpeI-5 (5'-aaACTAGTcgagcccgactgttgctga-3', SpeI site in bold) and cloned into pJET1.2 for sequencing. After sequence validation, the right-arm gene fragment was excised with SbfI and SpeI and inserted into pAG1-H3-right arm between the same sites. This disruption plasmid was designed to replace a portion of the *lacs* gene with the hygromycin phosphotransferase gene of pAG1-H3 upon double homologous recombination with the *H. irregularis* genome.

Agrobacterium-mediated transformation was performed according to the previously reported procedure.³⁰ We used the modified WSH agar (1.0% crushed oatmeal, 0.1% KH₂PO₄, 0.1% NaNO₃, 0.1% MgSO₄, 0.5% glycerol, 0.1% MES, 2.0% agar) plates supplemented with 200 $\mu\text{g}/\text{mL}$ hygromycin and 500 $\mu\text{g}/\text{mL}$ kanamycin (final concentrations) to select the correct transformants at 30 °C. The growing fungal spores were transferred onto new oatmeal plates with 500 $\mu\text{g}/\text{mL}$ carbenicillin and 200 $\mu\text{g}/\text{mL}$ hygromycin for the second-round selection.

The spores of the selected mutant were cultured in YM broth (yeast extract, 4 g/L; malt extract, 10 g/L; glucose, 4 g/L) for 7 days to extract DNA according to the extract protocol for the wild type. Four genome- and vector-specific primers (A–D) were designed to confirm the correct mutant. A 4534-bp fragment was amplified from the correct double crossover mutant with primers A (5'-ATGTATGACAATGGCCATCATAC-3') and B (5'-CGGAGACGCTGTGCAACTTT-3'), and a 5,026-bp fragment was amplified with primers C (5'-AGCTTGACTATGAAAATTCGTCAC-3') and D (5'-CACCAGTCATGTAGACCCTTCC-3'). The genomic DNA of wild-type *H. irregularis* FERM BP-2511 was used as the template to amplify the same fragments by the same primers, but no target bands were obtained.

The mutant and wild-type strains were cultured on the same oatmeal plate without any antibiotics at 30 °C for 7 days. The cultures were harvested and extracted three times with methanol. After drying, the extracts were redissolved in 1 mL of methanol for analysis on an Agilent 1200 HPLC with an Agilent XDB-C₁₈ column (5 μm , 4.6 mm \times 250 mm), eluted with a gradient of acetonitrile–water (0–10 min,

50%; 10–50 min, 50–90%) with 0.1% formic acid at a flow rate of 1 mL/min.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.jnatprod.6b00808](https://doi.org/10.1021/acs.jnatprod.6b00808).

Optical rotation values of the parent depsipeptides and hydrolyzed units (Table S1), Ca^{2+} flux data for 1–7 (Table S2), extracted signature sequences of the A domains of LACS (Table S3), HPLC analysis of the metabolites of *Hapsidospora irregularis* FERM BP-2S11 (Figure S1), NMR spectra of compounds 1–7 (Figures S2–S33), and the hydrolysis of compounds 2, 4, and 6 (Figure S34) (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*Tel: +1 435 797 8774. Fax: +1 435 797 2769. E-mail: wangwei402@hotmail.com.

*E-mail: jixun.zhan@usu.edu.

ORCID

Wei Wang: 0000-0003-0876-2205

Lijiang Xuan: 0000-0002-0593-3921

Jixun Zhan: 0000-0003-0200-9183

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by a Utah State University SPARC grant, a National Science Foundation Award CHE-1429195, and NIH grants ES017431 and GM107557. The authors thank Dr. S. T. Lee at the USDA Poisonous Plant Research Lab for helping measure the optical rotation data, and Dr. R. Shin at the University of Alabama at Birmingham for collecting the HMBC spectrum of 2.

■ REFERENCES

- (1) Chakraborty, T. K.; Ghosh, S.; Dutta, S. *Tetrahedron Lett.* **2001**, 42, 5085–5088.
- (2) Thell, K.; Hellinger, R.; Schabbauer, G.; Gruber, C. W. *Drug Discovery Today* **2014**, 19, 645–653.
- (3) Boros, C.; Smith, C. J.; Vasina, Y.; Che, Y.; Dix, A. B.; Darveaux, B.; Pearce, C. J. *Antibiot.* **2006**, 59, 486–494.
- (4) Lang, G.; Kalvelage, T.; Peters, A.; Wiese, J.; Imhoff, J. F. *J. Nat. Prod.* **2008**, 71, 1074–1077.
- (5) Chung, Y.-M.; El-Shazly, M.; Chuang, D.-W.; Hwang, T.-L.; Asai, T.; Oshima, Y.; Ashour, M. L.; Wu, Y.-C.; Chang, F.-R. *J. Nat. Prod.* **2013**, 76, 1260–1266.
- (6) Randazzo, A.; Bifulco, G.; Giannini, C.; Bucci, M.; Debitus, C.; Cirino, G.; Gomez-Paloma, L. *J. Am. Chem. Soc.* **2001**, 123, 10870–10876.
- (7) Kitagaki, J.; Shi, G.; Miyauchi, S.; Murakami, S.; Yang, Y. *Anti-Cancer Drugs* **2015**, 26, 259–271.
- (8) Stoianov, A. M.; Robson, D. L.; Hetherington, A. M.; Sawyez, C. G.; Borradaile, N. M. *PLoS One* **2015**, 10, e0131269.
- (9) Zhang, G.; Liu, S.; Liu, Y.; Wang, F.; Ren, J.; Gu, J.; Zhou, K.; Shan, B. *Oncol. Lett.* **2014**, 8, 248–252.
- (10) Hamano, K.; Kinoshita, M.; Furuya, K.; Miyamoto, M.; Takamatsu, Y.; Hemmi, A.; Tanzawa, K. *J. Antibiot.* **1992**, 45, 899–905.
- (11) Hamano, K.; Kinoshita, M.; Tanzawa, K.; Yoda, K.; Ohki, Y.; Nakamura, T.; Kinoshita, T. *J. Antibiot.* **1992**, 45, 906–913.
- (12) Abernethy, D. R.; Schwartz, J. B. *N. Engl. J. Med.* **1999**, 341, 1447–1457.
- (13) Grossman, E.; Messerli, F. H. *Prog. Cardiovasc. Dis.* **2004**, 47, 34–57.
- (14) Kakule, T. B.; Zhang, S.; Zhan, J.; Schmidt, E. W. *Org. Lett.* **2015**, 17, 2295–2297.
- (15) Zhang, S.; Wang, S.; Zhang, Q.; Chang, C.-W. T.; Zhan, J. *Bioorg. Med. Chem. Lett.* **2015**, 25, 1920–1924.
- (16) Katchalski, E.; Sela, M. *J. Am. Chem. Soc.* **1953**, 75, 5284–5289.
- (17) Beddow, J. E.; Davies, S. G.; Ling, K. B.; Roberts, P. M.; Russell, A. J.; Smith, A. D.; Thomson, J. E. *Org. Biomol. Chem.* **2007**, 5, 2812–2825.
- (18) Bauer, T.; Gajewiak, J. *Tetrahedron* **2004**, 60, 9163–9170.
- (19) Plaisance, G. P. *J. Am. Chem. Soc.* **1917**, 39, 2087–2088.
- (20) MacDonald, J. C. *Can. J. Chem.* **1969**, 47, 2739–2746.
- (21) Caceres, A. I.; Brackmann, M.; Elia, M. D.; Bessac, B. F.; del Camino, D.; D'Amours, M.; Witek, J. S.; Fanger, C. M.; Chong, J. A.; Hayward, N. J.; Homer, R. J.; Cohn, L.; Huang, X.; Moran, M. M.; Jordt, S. E. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, 106, 9099–9104.
- (22) Liu, B.; Escalera, J.; Balakrishna, S.; Fan, L.; Caceres, A. I.; Robinson, E.; Sui, A.; McKay, M. C.; McAlexander, M. A.; Herrick, C. A.; Jordt, S. E. *FASEB J.* **2013**, 27, 3549–3563.
- (23) Bessac, B. F.; Jordt, S. E. *Physiology* **2008**, 23, 360–370.
- (24) Rausch, C.; Weber, T.; Kohlbacher, O.; Wohlleben, W.; Huson, D. H. *Nucleic Acids Res.* **2005**, 33, 5799–5808.
- (25) Magarvey, N. A.; Beck, Z. Q.; Golakoti, T.; Ding, Y.; Huber, U.; Hemscheidt, T. K.; Abelson, D.; Moore, R. R.; Sherman, D. H. *ACS Chem. Biol.* **2006**, 1, 766–779.
- (26) Malloch, D.; Cain, R. F. *Can. J. Bot.* **1970**, 48, 1815–1825.
- (27) Deering-Rice, C. E.; Romero, E. G.; Shapiro, D.; Huguen, R. W.; Light, A. R.; Yost, G. S.; Veranth, J. M.; Reilly, C. A. *Chem. Res. Toxicol.* **2011**, 24, 950–959.
- (28) Doyle, J. J.; Doyle, J. L. *Phytochem. Bull.* **1987**, 19, 11–15.
- (29) Ashktorab, H.; Cohen, R. J. *Biotechniques* **1992**, 13, 198–200.
- (30) Zhang, A.; Lu, P.; Dahl-Roshak, A. M.; Paress, P. S.; Kennedy, S.; Tkacz, J. S.; An, Z. *Mol. Genet. Genomics* **2003**, 268, 645–655.