

Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

PYRIDINIUM SULFONATE CHLOROCHROMATE (VI), $C_5H_5NSO_3H$ [CrO₃CI] (PSCC): A NEW REAGENT FOR OXIDATION OF ORGANIC SUBSTRATES

M. Z. Kassaee^a & A. R. Bekhradnia^a ^a Department of Chemistry, Faculty of Science, Tarbiat Modarres University, Tehran, Iran

Version of record first published: 16 Aug 2010.

To cite this article: M. Z. Kassaee & A. R. Bekhradnia (2004): PYRIDINIUM SULFONATE CHLOROCHROMATE (VI), C₅H₅NSO₃H [CrO₃CI] (PSCC): A NEW REAGENT FOR OXIDATION OF ORGANIC SUBSTRATES, Phosphorus, Sulfur, and Silicon and the Related Elements, 179:10, 2025-2028

To link to this article: <u>http://dx.doi.org/10.1080/10426500490473555</u>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.tandfonline.com/page/terms-and-conditions

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



PYRIDINIUM SULFONATE CHLOROCHROMATE (VI), C₅H₅NSO₃H [CrO₃Cl] (PSCC): A NEW REAGENT FOR OXIDATION OF ORGANIC SUBSTRATES

M. Z. Kassaee and A. R. Bekhradnia Department of Chemistry, Faculty of Science, Tarbiat Modarres University, Tehran, Iran

(Received January 27, 2004; accepted March 16, 2004)

A fast, mild, and reasonable oxidizing agent, pyridinium sulfonate chlorochromate (VI), $C_5H_5NSO_3H[CrO_3Cl]$ (PSCC) is synthesized. Its reactions with primary, secondary, benzylic, and allylic alcohols under very mild conditions give the corresponding carbonyl compounds. These are obtained with relatively short reaction times. The oxidant/substrate ratios of 1:1 are employed.

Keywords: Carbonyl compounds; oxidation of alcohols

INTRODUCTION

The value of mild oxidation of alcohols to their corresponding aldehydes is evident.¹⁻⁴ Many reagents and methods have been reported that carry out this fundamental transformation. Among these, chromium (VI)-based oxidizing agents have become quite popular. Typical examples are: the Collins reagent,⁵ chromium trioxide-3,5-dimethyl-pyrazole complex,⁶ pyridinium chlorochromate,⁷ pyridinum dichromate,⁸ 2,2'-bipyridinium chlorochromate,⁹ and 3,5 dimethylpyrazolium fluorochromate (VI).¹⁰

In this article a novel, highly efficient, and widely applicable method for fast and mild oxidation of alcohols to their corresponding aldehydes is reported. First, pyridinium sulfonate chloride (I) is prepared via addition of chlorosulfonic acid to pyridine in chloroform.¹¹ Consequently, pyridinium sulfonate chlorochromate (VI), $C_5H_5NSO_3H$

Address correspondence to M. Z. Kassaee, Department of Chemistry, Faculty of Science, Tarbiat Modarres University, PO Box 14155-4838, Tehran, Iran. E-mail: kassaeem@modares.ac.ir

 $[CrO_3Cl]$ (PSCC) is synthesized through the reaction of CrO_3 with aqueous I (Scheme 1). This is done using a molar ratio of 1:1 An excellent yield of PSCC is obtained. PSCC is soluble in dichloromethane, acetonitril, and chloroform. Its melting point is 65°C.



SCHEME 1

RESULTS AND DISCUSSION

PSCC readily oxidizes primary, allylic, benzylic, and secondary alcohols to their corresponding carbonyls. Also, anthracene is oxidized to 9,10-anthraquinone (Table I).

TABLE I 1 H NMR and GC Analyses of Oxidation of Organic Substrates with Pyridinium-1-sulfonate Chlorochromate

Entry	Substrates	Products	Yield% (by ¹ H NMR and GC)
1	Allyl alcohol	Acrolein	80
2	Anthracene	9,10-Anthraquinone	57
3	p-Methoxybenzyl alcohol	p-Methoxybenzaldehyde	95
4	Benzyl alcohol	Benzaldehyde	91
5	Furfuryl alcohol	Furfural	55
6	1-Pentanol	1-Pentanal	65
7	1-Butanol	1-Butanal	60
8	Ethylene glycol	Glyoxal	50
9	Isobutanol	2-Methylpropanal	65
10	Isopropyl alcohol	Acetone	68
11	2-Butanol	2-Butanone	66
12	2-pentanol	2-Pentanone	69

EXPERIMENTAL

Preparation of PSCC

Chlorosulfonic acid (23 ml) is added dropwise to a stirred solution of pyridine (64 ml) in chloroform (100 ml) at -5° C. The reaction mixture is stirred at the same temperature for 20 min. Pyridinium sulfonate chloride is formed as a white solid and separated. This solid (4.25 g) is dissolved in 2 ml of water, and CrO₃ (2 g) is added and stirred for 30 min at room temperature. An orange-red solution is obtained. The reaction mixture is then cooled in an ice bath (0–5°C). The yield of C₅H₅NSO₃H [CrO₃Cl] is found to be 91%. The extend of the oxidation is determined via proton nuclear magnetic resonance (¹H NMR) and gas chromatography (GC) (Table I).

General Experimental Procedure for the Application of PSCC as an Oxidizing Agent

To a stirred solution of PSCC (5 mmole) in CH_2Cl_2 (10 ml), a solution of alcohol (5 mmole) in CH_2Cl_2 (10 ml) is added in one portion. Stirring is continued for 30 min to give a yellow solution. The progress of the reaction is monitored by thin layer chromatography (TLC) (ethyl acetate-hexane (1:9)). After completion of the reaction, dry diethyl ether (50 ml) is added to the reaction mixture.

The whole mass is shaken well. The organic layer is decanted, and the residue is washed with dry diethyl ether $(3 \times 20 \text{ ml})$. The combined organic layers are passed through a short pad of celite. This is to trap the reduced chromium. The product of oxidation is also washed thoroughly with dietyl ether $(3 \times 10 \text{ ml})$. The solvent is removed in a rotavapor under reduced pressure.*

REFERENCES

- [1] J.-D. Lou and Z.-N. Xu, Tetrahedron Lett., 43, 6095 (2002).
- [2] M. N. Bhattacharjee, M. K. Chaudhuri, and S. Purkayastha, *Tetrahedron*, 43, 5389 (1987).

^{*1}H NMR, ¹³C NMR, IR, and UV spectra were entirely consistent with the assigned structure of PSCC. Selected data are as follows: ¹H NMR (90 MHz, CH₃CN): δ = 7.9 (m, 2H, *m*-CH), 8.5 (m, 1H, *p*-CH), 8.8 (m, 2H, *o*-CH), ¹³C NMR (90 MHz, CH₃CN): δ = 127.33 (2C, *m*-CH), 141 (1C, *p*-CH), 178 (2C, *o*-CH); IR ν (KBr): 954 (s, Cr-O), 921 (s, Cr-O), 435 (m, Cr-Cl), 1150–1250 (s, $-SO_3$) 1440–1640 (originating due to the pyridinium solfonate cation); UV(CH³CN): 350–400 (CrCl), 256–260 (originating due to the pyridinium solfonate cation).

Elemental analysis data for the PSCC is found as follows: Cr, 17.6%; C, 20.5%; N, 4.80%; H, 1.95%. Calculated: Cr, 17.65%; C, 20.37%; N, 4.75%; H, 1.83%.

- [3] T. Mukaiyama, J.-I. Matsuo, D. Iada, and H. Kitagawa, Chem. Lett., 30, 864 (2001).
- [4] A. A. Wolf, J. Fluorine Chem., 27, 285 (1985).
- [5] J. C. Collins, W. W. Hes, and F. J. Frank, Tetrahedron Lett., 9, 3363 (1968).
- [6] E. J. Corey and G. W. J. Fleet, Tetrahedron Lett., 44, 4499 (1973).
- [7] E. J. Corey and J. W. Suggs, Tetrahedron Lett., 16, 2647 (1975).
- [8] E. J. Corey and G. Schmidt, Tetrahedron Lett., 20, 399 (1979).
- [9] F. S. Guziece and F. A. Luzzio, Synthesis, 691 (1980).
- [10] U. Bora, M. K. Chaudhuri, D. Dey, D. Kalita, W. Kharmawphlang, and G. C. Mandal, *Tetrahedron*, 57, 2445 (2001).
- [11] H. H. Silser and J. F. Andrieth, Inorg. Syn., 2, 173 (1946).