Laboratoire de Synthèse Organique, UA 40533 CNRS, Institut Curie, Section de Biologie, Bâtiment 110, Centre Universitaire, 91405 Orsay, France Received April 9, 1986

4-Methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (3) was synthetized from 2-acetylfuro[3,2-f]benzo[b]furan (4) or from 2-acetyl-5,6-dihydrofuro[3,2-f]benzo[b]furan (10). The key step involves a rearrangement-cyclization of azides 6 and 12 to form 4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridin-1(2H)-one (7) and 8,9-dihydro-4-methylfuro[3',2':5,6]benzofuro[3,2c]pyridin-1(2H)-one (13). Introduction of an aminoalkyl chain on carbon 1 was effected by substitution of 1-chloro-4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (8).

### J. Heterocyclic Chem., 23, 1637 (1986).

Recently we described the synthesis of 5*H*-furo-[3',2':5,6]benzopyrano[3,4-c]pyridin-5-one (1) and 6-methyl-8*H*-pyrano[3',2':5,6]benzofuro[3,2-c]pyridin-8-one (2) [1], a new class of psoralen derivatives behaving as monofunctional DNA photobinding agents [2]. These compounds in combination with 365 nm light exhibit high antiproliferative activity [3]. In addition, 1a and 1b have been shown to possess photochemotherapeutic effectiveness in the topical treatment of psoriasis [4].



As a further development, we intended to prepare 4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (3), a new planar tetracyclic heterocycle in which the  $\alpha$ -pyrone ring of the pyridopsoralen skeleton is replaced by a furan ring, with a view to evaluate the effect of this modification on photobiological activities.

The synthesis of the title compound has been accomplished by using 2-acetylfuro[3,2-f]benzo[b]furan (4) as starting material (Scheme 1). The sodium salt of triethylphosphonoacetate was condensed with ketone 4. Subsequent hydrolysis gave acid 5 in 71% yield as a  $\frac{1}{2}$  mixture



of Z and E isomers. This crude isomeric mixture was converted to the acid azide 6 by the mixed anhydride method [5]. Rearrangement of the azide 6 with concomitant cyclization was accomplished by a modified Eloy and Deryckere reaction [6,7] to afford compound 7 in 74% yield, which was transformed into the chloro derivative 8 by refluxing in phosphorus oxychloride. The target compound 3 was obtained from 8 by reduction with zinc and acetic acid in 57% yield.

Alternatively compound **3** could also be obtained by palladium-charcoal dehydrogenation of the dihydro derivative **9** which was prepared from 2-acetyl-5,6-dihydrofuro-[3,2-f]benzofuran (**10**) following the same general procedure employed for the synthesis of **3** (Scheme 2). However, the former route (Scheme 1) proved to be more convenient as it did not involve the dehydrogenation step required in the procedure outlined in Scheme 2. Furthermore this would allow the introduction of an aminoalkyl



side chain on carbon 1 which is expected to increase the affinity of the molecule toward DNA. This was illustrated by the nucleophilic substitution of the chloro derivative **8** with excess of 3-(diethylamino)propylamine affording the [3-(diethylamino)propyl]amino derivative **15**.

### **EXPERIMENTAL**

Melting points were determined with a Reichert hot-stage microscope and are uncorrected. The ir spectra were recorded on a Perkin-Elmer double beam spectrometer model 21. The nmr spectra were obtained with a Varian XL100 (100-MHz) spectrometer operating in the Fourier transform mode. Mass spectra were recorded on a MS30 AEI Kratos apparatus equiped with a computer Varian 100MS. Elemental analyses were performed by the Service Central de Microanalyses du C.N.R.S.

#### 6-Hydroxy-5-benzofurancarboxaldehyde [8].

This known compound was obtained from 2,3-dihydro-6-hydroxy-5-benzofurancarboxaldehyde [9] with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone in refluxing dioxane, as previously described [10] for oxidation of 2,3-dihydro-6-hydroxy-7-methoxy-5-benzofurancarboxaldehyde to 6hydroxy-7-methoxy-5-benzofurancarboxaldehyde.

# 2-Acetylfuro[3,2-f]benzo[b)furan (4).

6-hydroxy-5-benzofurancarboxaldehyde was converted to compound **4** as described in the literature [11] in 75% yield, mp 169-170° (lit mp [11] 169°).

### 2-Acetyl-5,6-dihydrofuro[3,2-f]benzo[b]furan (10).

Compound 10 was prepared from 2,3-dihydro-6-hydroxy-5-benzofurancarboxaldehyde [9] following the same procedure as described for 4 [11]. 2,3-Dihydro-6-hydroxy-5-benzofurancarboxaldehyde (6.56 g, 40 mmoles) was treated with chloroacetone (4.18 g, 45 mmoles) in 2-butanone (72 ml) in presence of potassium carbonate (12 g) at reflux for 30 minutes. Workup followed by recrystallization from ethanol gave 6.2 g (76%) of 10, mp 157°; nmr (deuteriochloroform):  $\delta$  2.54 (s, 3H, COCH<sub>3</sub>), 3.28 (t, 2H, J = 8.5 Hz, 5-CH<sub>2</sub>), 4.68 (t, 2H, J = 8.5 Hz, 6-CH<sub>2</sub>). 6.93 (1H, d, J = 0.4 Hz, 3-H), 7.4 (d, 1H, J = 0.8 Hz, 4-H), 7.43 (m, 1H, 8-H).

Anal. Calcd. for C<sub>12</sub>H<sub>10</sub>O<sub>3</sub>: C, 71.28; H, 4.99. Found: C, 70.94; H, 4.90.

### 3-[2'(Furo[3,2-f]benzo[b]furyl)]crotonic Acid (5).

Sodium hydride (50% dispersion in mineral oil) (1.2 g, 25 mmoles) was washed twice with dry hexane under argon. To the dry residue in dry tetrahydrofuran (10 ml) triethylphosphonoacetate (5.67 g, 25.3 mmoles) was added dropwise with stirring an cooling. The reaction mixture was stirred at room temperature for 30 minutes then compound 4 (5.06 g, 25.3 mmoles) in tetrahydrofuran (160 ml) was added over 15 minutes. The red solution was stirred at room temperature for 1 hour and refluxed for 45 minutes. The solvent was evaporated, to the residue was added ethanol (20 ml), 2N potassium hydroxide (50 ml) and the mixture was refluxed for 2.5 hours. The cooled solution was diluted with water and extracted with methylene chloride. The aqueous layer was acidified with N hydrochloric acid and the resulting precipitate was collected, washed with water, dried and recrystallized from toluene to yield 4.39 g (72%) of acid 5 as a  $\frac{1}{2}$  mixture of Z and E isomers (as indicated by the nmr). A portion of this mixture was recrystallized twice from ethanol to give pure E isomer as pale yellow needles, mp 238-240°; nmr (DMSO-d<sub>6</sub>):  $\delta$  2.52 (d,  $3H, J = 1.3 Hz, CH_3$ , 6.51 (d, 1H,  $J = 1.3 Hz, C = CH \cdot COOH$ ), 7.06 (dd, 1H, J = 2.3 Hz, J = 0.7 Hz, 5-H), 7.47 (br s, 1H, 8-H), 7.91 (br s, 2H, 4-H and 3-H), 8.04 (d, 1H, J = 2.3 Hz, 6-H).

Anal. Calcd. for C<sub>14</sub>H<sub>10</sub>O<sub>4</sub>; C, 69.42; H, 4.16. Found: C, 69.51; H, 4.14.

# 3-[2'(5',6'-dihydrofuro[3,2-f]benzo[b]furyl)]crotonic Acid (11).

Compound 11 was prepared following the procedure described above for 5. From the dihydro derivative 10 (5.59 g, 27.6 mmoles) there was obtained after recrystallization from toluene 5.1 g (75%) of acid 11 as a  $\frac{1}{2}$ mixture of Z and E isomers. A portion of this mixture was recrystallized twice from ethanol to give pure E isomer as pale yellow needles, mp 228-230°; nmr (DMSO-d<sub>6</sub>:  $\delta$  2.46 (d, 1H, J = 1.2 Hz, CH<sub>3</sub>), 3.26 (t, 2H, J = 8.5 Hz, 5-CH<sub>2</sub>), 4.64 (t, 2H, J = 8.5 Hz, 6-CH<sub>2</sub>), 6.38 (d, J = 1.2 Hz, C = CH-COOH), 7.02 (s, 1H, 3-H), 7.28 (s, 1H, 4-H), 7.49 (s, 1H, H-8). Anal. Calcd. for C<sub>14</sub>H<sub>12</sub>O<sub>4</sub>: C, 68.84; H, 4.95. Found: C, 69.14; H, 4.93.

### 4-Methylfuro[3',2':5,6]benzofuro[3,2-c]pyridin-1(2H)-one (7).

Ethyl chloroformate (2.20 g, 20 mmoles) in acetone (30 ml) was added dropwise to a stirred solution at  $0^{\circ}$  of the crude mixture of acids 5 (4.38 g, 18 mmoles) and triethylamine (2.3 ml) in acetone (15 ml). Stirring was continued for 1 hour at  $0^{\circ}$ , then a solution of sodium azide (1.8 g, 27 mmoles) in water (5 ml) was added. The solution was poured in ice-water (200 ml). The resulting precipitate formed was filtered off and dried to afford 4 g (83%) of azide 6; ir (nujol):  $2.150 \text{ cm}^{-1}$  (N<sub>3</sub>).

A solution of the crude azide **6** (3.97 g, 14.8 mmoles) in diphenyl ether at 30° (50 ml) was added dropwise to a rapidly stirred mixture of diphenyl ether (50 ml) and tributylamine (5 ml) at 250° over 20 minutes. Stirring was continued for 15 minutes at 250°, then the solution was concentrated under reduced pressure to *ca* 70 ml and after cooling hexane was added. The precipitate was filtered off, washed with hexane and dried to afford 3.24 g (91%) of crude 7. Recrystallization from toluene gave colourless crystals, mp > 290° dec; nmr (DMSO-d<sub>6</sub>):  $\delta$  2.32 (d, 3H, J = 1 Hz, 4-CH<sub>3</sub>), 7.15 (dd, 1H, J = 2.2 Hz, J = 1 Hz, 9-H), 7.39 (br s, 1H, 10-H), 8.10-8.07 (m, 2H, 6-H and 8-H), 8.26 (br s, 1H, 3-H), 8.87 (m, 1H, NH).

Anal. Calcd. for  $C_{14}H_9NO_3$ : C, 70.29; H, 3.79; N, 5.86. Found: C, 70.56; H, 4.13; N, 5.59.

# 8,9-Dihydro-4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridin-1(2H)-one (13).

Compound 13 was prepared following the procedure described above for compound 7. From the crude mixture of acids 11 (5.59 g, 19.26 mmoles) there was obtained 4.41 g (85%) of azide 12; ir (nujol): 2.150 cm<sup>-1</sup> N<sub>3</sub>).

The azide 12 (4.3 g, 16 mmoles) gave 3.20 g (84%) of compound 13. Recrystallization from toluene afforded colourless needles, mp > 290° dec; nmr (deuteriochloroform):  $\delta$  2.34 (d, 3H, J = 1 Hz, CH<sub>3</sub>), 3.35 (t, 2H, J = 8.5 Hz, 9-CH<sub>2</sub>), 4.71 (t, 2H, J = 8.5 Hz, 8-CH<sub>2</sub>), 7.01 (s, 1H, 6-H), 7.18 (d, 1H, J = 1 Hz, 3-H), 7.99 (s, 1H, 10-H).

Anal. Calcd. for C<sub>14</sub>H<sub>11</sub>NO<sub>3</sub>: C, 69.70; H, 4.59; N, 5.80. Found: C, 69.98; H, 4.55; N, 6.03.

# 1-Chloro-4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (8).

Compound 7 (3 g, 11.67 mmoles) was refluxed with phosphorus oxychloride (80 ml) for 6 hours. Excess of phosphorus oxychloride was removed under reduced pressure. The residue was taken up in methylene chloride, treated with aqueous solution was extracted with methylene chloride. Evaporation of the solvent gave a residue which was recrystallized from toluene to give 1.6 g of compound 8, mp 202-203°; nmr (deuteriochloroform):  $\delta$  2.56 (d, 3H, J = 1 Hz, 4-CH<sub>3</sub>), 6.96 (dd, 1H, J = 2.2 Hz, J = 1 Hz, 9-H), 7.73 (d, 1H, J = 1 Hz, 6-H), 7.74 (d, 1H, J = 2.2 Hz, 8-H), 8.22 (d, 1H, J = 0.7 Hz, 10-H), 8.46 (d, 1H, J = 1 Hz, 3-H). Anal. Calcd. for C<sub>14</sub>H<sub>8</sub>CINO<sub>2</sub>: C, 65.25; H, 3.13; Cl, 13.77; N, 5.44.

Found: C, 65.36; H, 3.09; Cl, 13.58; N, 5.21. Chromatography of the mother liquor on silicagel with methylene

chloride-ethanol (98/2) gave after recrystallization from toluene an additional crop of 0.36 g (overall yield 61%) of 8.

### 1-Chloro-8,9-dihydro-4-methylfuro[3',2':5,6]benzofuro[3,2-c]-pyridine (14).

Compound 14 was synthesized following the procedure described above for the preparation of 8. From 13 (3.1 g, 12.8 mmoles) there was obtained 1.45 g (43%) of compound 14 as colourless crystals (from toluene), mp 180°; nmr (deuteriochloroform):  $\delta$  2.52 (d, 3H, J = 0.8 Hz, 4-CH<sub>3</sub>), 3.38 (t, 2H, J = 8.5 Hz, 9-CH<sub>2</sub>), 4.74 (t, 2H, J = 8.5 Hz, 8-CH<sub>2</sub>), 7.02 (s, 1H, 6-H), 8.03 (1H, s, 10-H), 8.14 (d, 1H, J = 0.8 Hz, 3-H).

Anal. Calcd. for  $C_{14}H_{10}CINO_2$ : C, 64.75; H, 3.88; N, 5.39. Found: C, 64.68; H, 3.91; N, 5.71.

### 8,9-Dihydro-4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (9).

Compound 14 (0.772 g, 2.98 mmoles) in ethyl acetate (100 ml) was hydrogenated at room temperature and atmospheric pressure in presence of anhydrous sodium acetate (0.24 g) using 10% palladium on charcoal (0.080 g) as catalyst. After 7 hours the catalyst was filtered off and washed with ethyl acetate. The filtrate was evaporated to dryness and the residue recrystallized from ethanol to afford 0.483 g (72%) of the dihydro derivative 9 as colourless needles, mp 174-175°; nmr (deuteriochloroform):  $\delta$  2.54 (s, 3H, 4-CH<sub>3</sub>), 3.35 (t, 2H, J = 8.5 Hz, 9-CH<sub>3</sub>), 4.72 (t, 2H, J = 8.5 Hz, 8-CH<sub>2</sub>), 7.02 (s, 1H, 6-H), 7.75 (s, 1H, 10-H), 8.37 (s, 1H-3-H), 8.96 (br s, 1H, 1-H).

Anal. Calcd. for C14H11NO2: C, 74.65; H, 4.92; N, 6.22. Found: C, 74.60; H, 5.03; N, 6.33.

4-Methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (3).

### Method A.

The chloro derivative **8** (0.23 g, 0.89 mmole) in acetic acid (3 ml) was refluxed with powdered zinc (0.36 g) for 2 hours. After cooling, the reaction mixture was filtered off and the solid washed with acetic acid. The filtrate was evaporated under reduced pressure, the residue was taken up in water and made alkaline with 20% aqueous sodium hydroxide. The aqueous solution was extracted with methylene chloride. Evaporation of the solvent gave a residue which was recrystallized from aqueous methanol to afford 0.114 g (57%) of compound **3** as colourless needles, mp 167-169°; nmr (deuteriochloroform):  $\delta$  2.59 (s, 3H, 4-CH<sub>3</sub>), 6.92 (dd, 1H, J = 2 Hz, 9-H), 7.73 (m, 2H, 6-H and 8-H), 8.16 (s, 1H, 10-H), 8.46 (s, 1H, 3-H), 9.11 (s, 1H, 1-H): ms: M<sup>+</sup> (m/e) 233.

Anal. Calcd. for C14H, NO2: C, 75.32; H, 4.06; N, 6.28. Found: C, 74.88; H, 4.28; N, 5.94.

### Method B.

A mixture of compound 9 (0.24 g, 1.06 mmoles), 10% palladium on charcoal (0.22 g) and diphenyl ether (2.5 ml) was refluxed for 4 hours; it was then filtered whilst hot and the catalyst washed with hot ethanol. After removal of the ethanol the mixture was chromatographed on silica gel (10 g), eluting the diphenyl ether with methylene chloride. Fractions eluted with methylene chloride-ethanol (95/5) were recrystallized from ethanol to afford 0.12 g (50%) of compound 3 identical in all respects to the sample obtained in A.

1-[3-(Diethylamino)propylamino]-4-methylfuro[3',2':5,6]benzofuro[3,2-c]pyridine (15).

A mixture of compound **8** (0.247 g, 0.96 mmoles) and 3-(diethylamino)propylamine (5 ml) was refluxed for 6 hours. Excess of amine was removed under reduced pressure, the residue was taken up in 1N sodium hydroxide and extracted with methylene chloride. Evaporation of the solvent gave a residue which was chromatographed on neutral alumina (30 g) with methylene chloride to give 0.315 g (89%) of **15**; nmr (deuteriochloroform):  $\delta$  1.13 (t, 2 x 3H, CH<sub>3</sub>-CH<sub>2</sub>), 1.93 (m, 2H,  $\beta$ -CH<sub>2</sub>), 2.43 (d, 3H,  $J=0.7~Hz,\,4\text{-}CH_3),\,2.8\text{-}2.6~(m,\,3~x~2H,\,\gamma\text{-}CH_2~and~N\text{-}CH_2\text{-}CH_3),\,3.78~(m,\,2H,\,\alpha\text{-}CH_2),\,7.03~(br~s,\,1H,~NH),\,6.85~(dd,\,1H,~J=2.2~Hz,~J=0.7~Hz,\,9\text{-}H),\,7.68\text{-}7.69~(m,\,2H,~8\text{-}H~and~6\text{-}H),\,7.97~(d,\,1H,~J=0.9~Hz,\,10\text{-}H),\,8.02~(s,\,1H,~3\text{-}H).$ 

To a stirred solution of maleic acid (0.15 g, 1.29 mmoles) in acetone (1 ml) was added amine 15 (0.167 g, 0.475 mmole) in the minimum of acetone. The resulting precipitate was filtered off, washed with acetone and recrystallized from ethanol to yield 0.197 g of the dimaleate salt of amine 15, mp 176-177°.

Anal. Calcd. for  $C_{25}H_{33}N_3O_{16}$ : C, 59.68; H, 5.70; N, 7.20. Found: C, 59.96; H, 5.71; N, 7.35.

### Acknowledgements.

We acknowledge support of this work by the Institut National de la Santé et de la Recherche Médicale (INSERM Contract CRE 832005). We thank Mrs. C. Huel for the nmr spectra.

### **REFERENCES AND NOTES**

[1] J. Moron, C. H. Nguyen and E. Bisagni, J. Chem. Soc., Perkin Trans. 1, 225 (1983).

[2] J. Blais, P. Vigny, J. Moron and E. Bisagni, *Photochem.* Photobiol., **39**, 145 (1984).

- [3] D. Averbeck, S. Averbeck, E. Bisagni and J. Moron, *Mutat. Res.*, **148**, 45 (1985).
- [4] D. Averbeck, L. Dubertret, E. Bisagni, J. Moron, D. Papadopoulo,
  S. Nocentini, J. Blais and F. Zajdela, J. Invest. Dermatol., 80, 306 (1983).

[5] J. Weinstock, J. Org. Chem., 26, 3511 (1961).

[6] F. Eloy and A. Deryckere, Helv. Chim. Acta, 52, 1755 (1969).

[7] E. Bisagni, J. D. Bourzat, J. P. Marquet, C. Labrid, P. Delort and A. Leridant, Chim. Ther. Eur. J. Med. Chem., 8, 559 (1973).

[8] J. R. Merchant and S. R. Auraval, J. Indian Chem. Soc., 40, 472 (1963).

- [9] E. C. Horning and D. B. Reisner, J. Am. Chem. Soc., 70, 3619 (1948).
- [10] Y. Y. Lin, E. Thom and A. Liebman, J. Heterocyclic Chem., 16, 799 (1979).
- [11] L. René, J. P. Buisson and R. Royer, Eur. J. Med. Chem., 12, 31 (1977).