

Acta Cryst. (1999). **C55**, 1513–1514

1,2,3,4,5,6,7,8-Octahydro-3,3,6,6-tetramethylacridine-1,8-dione

R. SANKARANARAYANAN,^a S. SHANMUGA SUNDARA RAJ,^b
D. VELMURUGAN^a AND HOONG-KUN FUN^b

^a*Department of Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and*
^b*X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia. E-mail: crystal@giasmd01.vsnl.net.in*


(Received 20 November 1998; accepted 4 May 1999)

Abstract

In the title compound, C₁₇H₂₁NO₂, the central pyridine ring is almost planar and the outer rings are half-chairs. The O atoms of the keto groups deviate significantly from the mean plane passing through their corresponding rings. The molecule contains a pseudo-mirror plane which passes through the central N and C atoms and the molecule is slightly folded along an axis passing through the same atoms. The dihedral angle between the outer rings is 6.5 (2)°. The packing is stabilized by an intermolecular C—H···O hydrogen bond.

Comment

Acridines are found to have a wide range of biological activities, such as mutagenic, antitumour (Talacki *et al.*, 1974), antibacterial (Achenson, 1956), anti-amoebic (Prasad Krishna *et al.*, 1984), hypertensive, antiinflammatory and antiimplantation (Asthana *et al.*, 1991) activities. A drug containing the acridine moiety has been found to possess antiprotozoal activity (Karolak-Wojciechowska *et al.*, 1996). The ability of acridine to intercalate between the base-pairs of DNA is also well known (Neidle, 1979; Fan *et al.*, 1997). Acridine compounds are considered to be efficient drugs for the treatment of Alzheimer's disease (Bandoli *et al.*, 1994). The use of decahydroacridine-1,8-diones as photo-sensitizers is also well known (Timpe *et al.*, 1993). In addition, acridine diones act as laser dyes whose laser activity has been studied extensively (Murugan *et al.*, 1998). Because of these wide-ranging biological and photochemical properties, an X-ray diffraction study has been undertaken for the title compound, (I), so that its structure–activity correlations could be studied.



Interestingly, the molecule of (I) contains a pseudo-mirror plane which passes through N10 and C5. A comparison of equivalent bond lengths and torsion angles across this pseudo-mirror plane indicates the presence of this non-crystallographic symmetry element. The keto bond distances C4=O4 [1.218 (3) Å] and C6=O6 [1.225 (4) Å] are comparable with those in similar structures (Sankaranarayanan *et al.*, 1998; Ganesh *et al.*, 1998). The deviations of atoms O4 and O6 from the mean planes passing through rings A and C are −0.287 (3) and −0.299 (3) Å, respectively. The π conjugation along C5—C6a—C9a—N10—C10—C4a [C5—C6a = 1.383 (4), C6a—C9a = 1.400 (4), C9a—N10 = 1.347 (3), N10—C10 = 1.350 (3), C10—C4a = 1.400 (4) and C5—C4a = 1.396 (4) Å] indicates the strong aromaticity in the central ring B, which makes all the atoms of the ring lie almost in a plane, with the maximum deviation being 0.006 (3) Å for C6a and C9a. This planarity of the central ring B is further supported by the low value of the puckering amplitude (Cremer & Pople, 1975), $Q_T = 0.009 (3)$ Å, when compared with the values for the other rings, A and C, for which $Q_T = 0.470 (3)$ and $0.477 (3)$ Å, respectively. The asymmetry parameter (Nardelli, 1983a) for ring C is $\Delta C_2 = 0.057 (2)$, revealing a half-chair conformation for this ring with a local pseudo-twofold axis running through the midpoint of the C7—C8 bond. The asymmetry parameter for ring A, $\Delta C_2(C2—C3) = 0.058 (1)$, reveals the conformation of ring A as a half-chair with a local twofold axis running through the midpoint of the C2—C3 bond. The dihedral angle between the two mean planes passing through C5/C6a/C6/C7/C8/C9/C9a/N10 and C5/C4a/C4/C3/C2/C1/C10/N10 is 5.7 (1)°, indicating that the acridine moiety is slightly folded about a line passing through N10 and C5, as observed in related structures (Gunasekaran *et al.*, 1996; Sankaranarayanan *et al.*, 1998). The dihedral angle between rings A and C is 6.5 (2)°.

In addition to van der Waals interactions, an intermolecular C—H···O hydrogen bond stabilizes the molecular packing: C14···O6ⁱ = 3.519 (4) and




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom-numbering scheme. H atoms are drawn as spheres of arbitrary radii.

H14A···O6ⁱ = 2.641 Å, and C14—H14A···O6ⁱ = 152.2° [symmetry code: (i) $x - 1, y, z$].

Experimental

3,3,6,6-Tetramethyl-3,4,6,7,9,10-hexahydro-1,8(2*H*,5*H*)-acridinedione (10 mmol) was dissolved in chloroform (100 ml) and stirred at room temperature with active MnO₂ (5 g) for 8 h. The MnO₂ was filtered off and the filtrate concentrated; the solid obtained was filtered and recrystallized from methanol.

Crystal data

C ₁₇ H ₂₁ NO ₂	Mo K α radiation
$M_r = 271.35$	$\lambda = 0.71073$ Å
Orthorhombic	Cell parameters from 39 reflections
$P2_12_12_1$	$\theta = 5.38\text{--}12.51^\circ$
$a = 6.9887$ (2) Å	$\mu = 0.077$ mm ⁻¹
$b = 10.7123$ (2) Å	$T = 293$ (2) K
$c = 20.4198$ (4) Å	Rectangular prism
$V = 1528.73$ (6) Å ³	$0.45 \times 0.38 \times 0.29$ mm
$Z = 4$	Pale yellow
$D_x = 1.179$ Mg m ⁻³	
D_m not measured	

Data collection

Siemens P4 diffractometer	$\theta_{\max} = 27.49^\circ$
$\omega/2\theta$ scans	$h = -8 \rightarrow 9$
Absorption correction: none	$k = -13 \rightarrow 13$
9982 measured reflections	$l = -20 \rightarrow 26$
2024 independent reflections	3 standard reflections
1501 reflections with $I > 2\sigma(I)$	every 97 reflections
$R_{\text{int}} = 0.047$	frequency: 60 min
	intensity decay: <3%

Refinement

Refinement on F^2	$(\Delta/\sigma)_{\max} < 0.001$
$R[F^2 > 2\sigma(F^2)] = 0.051$	$\Delta\rho_{\max} = 0.14$ e Å ⁻³
$wR(F^2) = 0.118$	$\Delta\rho_{\min} = -0.17$ e Å ⁻³
$S = 1.094$	Extinction correction: none
2024 reflections	Scattering factors from
181 parameters	<i>International Tables for</i>
H atoms constrained	<i>Crystallography</i> (Vol. C)
$w = 1/[\sigma^2(F_o^2) + (0.0394P)^2 + 0.3749P]$	
where $P = (F_o^2 + 2F_c^2)/3$	

Table 1. Selected geometric parameters (Å, °)

C1—C10	1.502 (4)	C6—C6a	1.490 (4)
C4—C4a	1.488 (4)	C6a—C9a	1.400 (4)
C4a—C5	1.396 (4)	C9—C9a	1.503 (4)
C5—C6a	1.383 (4)	C9a—N10	1.347 (3)
C6a—C5—C4a	119.5 (3)	C9a—N10—C10	118.1 (2)
C12—C2—C3—C4	64.8 (3)	C6—C7—C8—C14	-63.6 (3)
C11—C2—C3—C4	-175.7 (2)	C6—C7—C8—C13	176.7 (3)

All H atoms were geometrically fixed and allowed to ride on the corresponding non-H atoms, with C—H = 0.96–0.97 Å, and $U_{\text{iso}} = 1.5U_{\text{eq}}$ of the attached C atom for methyl-H atoms and $1.2U_{\text{eq}}$ for other H atoms. Because of the presence of only very weak anomalous scatterers such as O, N and C, the absolute structure cannot be determined reliably [the $F(hkl)$ and $F(\bar{h}\bar{k}\bar{l})$ reflections have been merged using the command MERG3 in SHELXL97 (Sheldrick, 1997a)].

Data collection: XSCANS (Siemens, 1994). Cell refinement: XSCANS. Data reduction: XSCANS. Program(s) used to solve structure: SHELXS97 (Sheldrick, 1997b). Program(s) used to refine structure: SHELXL97 (Sheldrick, 1997a). Molecular graphics: ZORTEP (Zsolnai, 1997). Software used to prepare material for publication: SHELXL97 and PARST (Nardelli, 1983b, 1995).

The authors thank Professor V. T. Ramakrishnan for providing the title compound for X-ray study. RS and DV thank the DST, India, for providing financial assistance. SSSR thanks Universiti Sains Malaysia for a visiting Postdoctoral Research Fellowship and HKF would like to thank the Malaysian Government and Universiti Sains Malaysia for research grant R&D No. 190-9609-2801.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FR1186). Services for accessing these data are described at the back of the journal.

References

- Achenson, R. M. (1956). In *Acridines. The Chemistry of Heterocyclic Compounds*, Vol. 9, edited by A. Weissberger, pp. 339–361. New York: Interscience.
- Asthana, P., Rastogi, S., Ghose, S. & Das, S. R. (1991). *Indian. J. Chem. B*, **30**, 893–900.
- Bandoli, G., Dolmella, A., Gatto, S. & Nicolini, M. (1994). *J. Chem. Cryst.* **24**, 301–310.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Fan, J.-Y., Terrel, M. & Denny, W. A. (1997). *Anti-Cancer Drug Des.* **12**, 277–293.
- Ganesh, V. K., Banumathi, S., Velmurugan, D., Ramasubbu, N. & Ramakrishnan, V. T. (1998). *Acta Cryst.* **C54**, 633–635.
- Gunasekaran, K., Velmurugan, D., Murugan, P., Ramakrishnan, V. T., Panneerselvam, K. & Soriano-García, M. (1996). *Acta Cryst.* **C52**, 2609–2612.
- Karolak-Wojciechowska, J., Morzek, A., Amiel, P., Brouant, P. & Barbe, J. (1996). *Acta Cryst.* **C52**, 2939–2941.
- Murugan, P., Shanmugasundaram, P., Ramakrishnan, V. T., Venkatachalapathy, B., Srividya, N., Ramamurthy, P., Gunasekaran, K. & Velmurugan, D. (1998). *J. Chem. Soc. Perkin Trans. 2*, pp. 999–1003.
- Nardelli, M. (1983a). *Acta Cryst.* **C39**, 1141–1142.
- Nardelli, M. (1983b). *Comput. Chem.* **7**, 95–98.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Neidle, S. (1979). *Prog. Med. Chem.* **16**, 151–221.
- Prasad Krishna, B. N., Banasal, I., Das, P. & Srivastava, R. (1984). *Curr. Sci.* **53**, 778–780.
- Sankaranarayanan, R., Velmurugan, D., Murugan, P. & Ramasubbu, N. (1998). *Acta Cryst.* **C54**, 1534–1535.
- Sheldrick, G. M. (1997a). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997b). *SHELXS97. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
- Siemens (1994). *XSCANS. X-ray Single-Crystal Analysis System*. Version 2.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Talacki, R., Carrel, H. L. & Glusker, J. P. (1974). *Acta Cryst.* **B30**, 1044–1047.
- Timpe, H. J., Ulrich, S., Decker, C. & Fouassier, J. P. (1993). *Macromolecules*, **26**, 4560–4566.
- Zsolnai, L. (1997). *ZORTEP. An Interactive Molecular Graphics Program*. University of Heidelberg, Germany.