This article was downloaded by: [Gebze Yuksek Teknoloji Enstitïsu ] On: 22 December 2014, At: 15:43 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lcar20

# Synthesis of the Trisaccharide Repeat of the O-Antigen of Rhanella Aquatilis 1-95

Goutam Guchhait<sup>a</sup> & Anup Kumar Misra<sup>a</sup>

<sup>a</sup> Division of Molecular Medicine , Bose Institute , A.J.C. Bose Centenary Campus, P-1/12, C.I.T. Scheme VII-M, Kolkata, 700054, India

Published online: 04 Jan 2010.

To cite this article: Goutam Guchhait & Anup Kumar Misra (2010) Synthesis of the Trisaccharide Repeat of the O-Antigen of Rhanella Aquatilis 1-95, Journal of Carbohydrate Chemistry, 29:1, 1-9, DOI: <u>10.1080/07328300903428989</u>

To link to this article: http://dx.doi.org/10.1080/07328300903428989

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

Journal of Carbohydrate Chemistry, 29:1–9, 2010 Copyright © Taylor & Francis Group, LLC ISSN: 0732-8303 print / 1532-2327 online DOI: 10.1080/07328300903428989



# Synthesis of the Trisaccharide Repeat of the O-Antigen of *Rhanella Aquatilis* 1-95

Goutam Guchhait and Anup Kumar Misra

Division of Molecular Medicine, Bose Institute, A.J.C. Bose Centenary Campus, P-1/12, C.I.T. Scheme VII-M, Kolkata 700054, India

A branched trisaccharide containing D-fucose and D-galactofuranose moieties corresponding to the repeating unit of the O-antigen of *Rahnella aquatilis* 1–95 has been synthesized from the monosaccharide intermediates in a five-step synthesis in an overall 38% yield. All glycosylation steps and protecting group functionalization steps are high yielding.

**Keywords** Trisaccharide; *Rahnella aquatilis*; Glycosylation; D-fucose; 2-Aminoethyl; D-Galactofuranose.

### INTRODUCTION

Rahnella aquatilis belongs to the family of Enterobacteriaceae, which are found in soil and water.<sup>[1]</sup> Although formerly *R. aquatilis* was assigned to the *Enterobacter* agglomerans biogroup, later it was recognized as a distinct species.<sup>[2]</sup> *R. aquatilis* is an opportunistic pathogen responsible for a number of pathogenic infections in the gastrointestinal and urinary tracts and in the respiratory and cardiovascular systems.<sup>[3]</sup> In the pathogenesis of diseases caused by Gram-negative bacteria, endotoxin (lipopolysaccharide) plays a significant role.<sup>[4]</sup> Recently, for the first time, the structure of the *O*-antigen of *R. aquatilis* strain 1–95 was reported by Zdorovenko et al. (Fig. 1).<sup>[4]</sup> The trisaccharide repeating unit of the *O*-antigen is branched in nature and contains uncommon monosaccharide units such as D-fucose and D-galactofuranose moieties.

Preparation of the glycoconjugate vaccines against infectious diseases is considered an important area in medicinal research.<sup>[5]</sup> The prime requirement for designing a glycoconjugate vaccine candidate is to have pure

Received August 12, 2009; accepted October 20, 2009.

Address correspondence to Anup Kumar Misra, Division of Molecular Medicine, Bose Institute, A.J.C. Bose Centenary Campus, P-1/12, C.I.T. Scheme VII-M, Kolkata 700054, India. E-mail: akmisra69@rediffmail.com

$$\alpha$$
-D-Galp  
1  
↓  
2  
→3)-β-D-Galf-(1→3)-α-D-Fucp-(1→

Figure 1: Structure of the repeating unit of the O-antigen of Rahnella aquatilis 1-95.

oligosaccharides or polysaccharides at hand. The limited availability of pure oligosaccharide fragments of a particular polysaccharide isolated from the natural sources cannot meet the requirement for their extensive biological studies. Concise chemical synthesis can provide access to larger quantities of pure oligosaccharides and several analogs thereof. As part of our program directed toward the synthesis of bacterial *O*-antigen fragments for use in glycoconjugate preparation,<sup>[6]</sup> we describe a concise chemical synthesis of a branched trisaccharide repeating unit of the *O*-antigen of *R. aquatilis* 1–95 as its 2aminoethyl glycoside (1, Fig. 2). The 2-aminoethyl group serves as a linker for the conjugation to a carrier protein.

### **RESULTS AND DISCUSSION**

In order to synthesize the target trisaccharide **1** having a branched D-fucopyranosyl moiety, three suitably protected monosaccharide intermediates (**2**, **3**, and **4**, Fig. 2) have been prepared. Starting from 1,2:3,4-di-*O*isopropylidene- $\alpha$ -D-galactopyranose (**5**),<sup>[7]</sup> D-fucose derivative **6**<sup>[8]</sup> was prepared in 90% yield following Lerner's procedure.<sup>[8]</sup> Compound **6** was converted to 2-azidoethyl 3,4-*O*-isopropylidene- $\alpha$ -D-fucopyranoside (**2**) following a set of reactions involving hydrolytic removal of isopropylidene acetal, Fischer



Figure 2: Structure of the target trisaccharide 1 and of the monosaccharide building blocks 2, 3, and 4.

### Synthesis of Rhanella aquatilis 1–95 3

glycosylation with 2-azido ethanol in the presence of HClO<sub>4</sub>-SiO<sub>2</sub>, <sup>[9]</sup> and isopropylidenation in 80% overall yield. Compound  $3^{[10]}$  was synthesized following literature-reported reaction conditions. Ethyl 2,3,5,6-tetra-O-acetyl-1-thio- $\alpha$ -D-galactofuranoside (4)<sup>[11]</sup> was prepared from D-galactose diethyldithioacetal (7) following the report of Wolfrom et al.<sup>[11]</sup> Stereoselective glycosylation of compound 2 with thioethyl glycoside 3 in the presence of a combination of copper(II) bromide-tetrabutylammonium bromide (TBAB)<sup>[12]</sup> in a mixed solvent (1,2-dichloroethane-DMF; 5:1, v/v) at room temperature furnished disaccharide derivative 8 in 76% yield. Appearance of a signature peak at  $\delta$  5.02 (br s, H-1<sub>B</sub>) in the <sup>1</sup>H NMR spectrum confirmed the stereoselective formation of compound 8. Removal of the isopropylidene acetal followed by selective acetylation of the axial hydroxyl group through orthoester formation<sup>[13]</sup> gave disaccharide acceptor 9 in 88% yield. Iodonium ion promoted stereoselective glycosylation of compound 9 with D-galactofuranose thioglycoside donor 4 in the presence of N-iodosuccinimide, and trimethylsilyl trifluoromethanesulfonate (TMSOTf)<sup>[14,15]</sup> combination furnished trisaccharide derivative 10 in 82% yield. Appearance of signals at  $\delta$  5.34 (br s, H-1<sub>C</sub>), 5.07 (d, J = 3.2 Hz, H-1<sub>B</sub>), and 5.00 (d, J = 3.2 Hz, H-1<sub>A</sub>) in the <sup>1</sup>H NMR spectrum and  $\delta$  107.3  $(C-1_C)$ , 96.9  $(C-1_A)$ , and 95.3  $(C-1_B)$  in the <sup>13</sup>C NMR spectrum supported the formation of compound 10. As the value of 1,2-coupling constant of the  $\beta$ -Dgalactofuranosyl linkage is very low,<sup>[16]</sup> sometimes it appears as a broad singlet. The formation of  $\beta$ -D-galactofuranosyl linkage can be explained by considering the neighboring participating O-acetyl group at the C-2 position of thioglycoside donor 4. Saponification of compound 10 followed by hydrogenolysis<sup>[17]</sup> furnished target trisaccharide 1 as its 2-aminoethyl glycoside in 70%vield (Sch. 1). Appearance of anomeric signals at  $\delta$  5.16 (br s, 1 H, H-1<sub>c</sub>), 5.01 (br s, 1 H, H-1<sub>B</sub>), and 5.00 (br s, 1 H, H-1<sub>A</sub>) in the <sup>1</sup>H NMR and at  $\delta$  107.8  $(C-1_C)$ , 94.5  $(C-1_A)$ , and 94.0  $(C-1_B)$  in the <sup>13</sup>C NMR confirmed the formation of trisaccharide 1. Anomeric signals of 1 were compared with those of the native polysaccharide [ $\delta$  5.25 (br s,  $\alpha$ -D-Fucp), 5.11 (br s,  $\alpha$ -D-Galp), and 5.23 (br s,  $\beta$ -D-Galf) in <sup>1</sup>H NMR and  $\delta$  110.3, 97.4, and 96.8 in <sup>13</sup>C NMR spectra].<sup>[4]</sup> The signals of the native polysaccharide have been shifted from compound 1 and may be due to the polysaccharide structure.

In summary, a concise synthesis of a branched trisaccharide repeating unit of the O-antigen of R. aquatilis 1–95 as its 2-aminoethyl glycoside was achieved in 38% yield starting from the three monosaccharide building blocks **2**, **3**, and **4**. The key features of the synthesis are the preparation of the D-fucose precursor from D-galactose and its subsequent glycosylation with 2-azido ethanol, as well as glycosylation of the D-galactofuranose residue. It is noteworthy that Dfucose and D-galactose moieties are linked through 1,2-*cis* linkages. The trisaccharide **1** can be conjugated to a carrier protein using the amino function of its 2-aminoethyl aglycon.



**Scheme 1: Reagents**: (a) 80% aq. AcOH, 80°C, 6 h; (b) 2-azidoethanol, HClO<sub>4</sub>-SiO<sub>2</sub>, 80°C, 50 h; (c) 2,2-dimethoxypropane, acetone, rt, 2 h, 80%; (d) Cu(II)Br<sub>2</sub>, TBAB, MS 4Å, 1,2-(CH<sub>2</sub>Cl)<sub>2</sub>-DMF, rt, 48 h, 76%; (e) 80% aq. AcOH, 80°C, 2 h; (f) CH<sub>3</sub>C(OC<sub>2</sub>H<sub>5</sub>)<sub>3</sub>, *p*-TsOH, DMF, rt, 2 h; 80% aq. AcOH, rt, 30 min, 88%; (g) *N*-iodosuccinimide, TMSOTf, CH<sub>2</sub>Cl<sub>2</sub>, MS 4Å, -40°C, 1 h, 82%; (h) (i) CH<sub>3</sub>ONa, CH<sub>3</sub>OH, rt, 5 h; (ii) 20% Pd(OH)<sub>2</sub>-C, CH<sub>3</sub>OH-AcOH-H<sub>2</sub>O, rt, 12 h, 70%.

### **EXPERIMENTAL**

### **General Procedure**

All reactions were monitored by thin layer chromatography over silica gelcoated TLC plates. The spots on TLC were visualized by warming ceric sulphate (2% Ce(SO<sub>4</sub>)<sub>2</sub> in 2N H<sub>2</sub>SO<sub>4</sub>) sprayed plates on a hot plate. Silica gel 230–400 mesh was used for column chromatography. <sup>1</sup>H and <sup>13</sup>C NMR, 2D COSY, and HMQC spectra were recorded on Bruker Avance DRX 500 MHz using CDCl<sub>3</sub> and D<sub>2</sub>O as solvents and TMS as internal reference unless stated otherwise. Chemical shift value is expressed in  $\delta$  ppm. ESI-MS were recorded on a Micromass Quattro II triple quadrupole mass spectrometer. Elementary analysis was carried out on Carlo ERBA-1108 analyzer. Optical rotations were measured at 25°C on a Perkin Elmer 341 polarimeter. Commercially available grades of organic solvents of adequate purity are used in many reactions.

### 2-Azidoethyl 3,4-O-isopropylidene- $\alpha$ -D-fucopyranoside (2)

A solution of compound **6** (4 g, 16.4 mmol) in 80% aq. AcOH (100 mL) was allowed to stir at  $80^{\circ}$ C for 6 h and evaporated and coevaporated with toluene

 $(3 \times 50 \text{ mL})$  under reduced pressure. To a solution of the crude product in 2-azidoethanol (15 mL) was added HClO<sub>4</sub>-SiO<sub>2</sub> (500 mg) and the reaction mixture was allowed to stir at 80°C for 50 h. The reaction mixture was cooled to rt and the solvents were removed under reduced pressure. To a solution of the crude reaction mixture in anhydrous acetone (30 mL) was added 2,2dimethoxypropane (5 mL, 40.6 mmol) and it was allowed to stir at rt for 2 h. The reaction mixture was neutralized with  $Et_3N$  (1 mL), filtered, and concentrated under reduced pressure. The crude product was purified over  $m SiO_2$  using hexane-EtOAc (4:1) as eluant to give pure compound 2 (3.6 g, 80%). Yellow oil;  $[\alpha]_{D}^{25}$  +5 (c 1.6, CHCl<sub>3</sub>); IR (neat): 3450, 2989, 2933, 2877, 2107, 1738, 1635, 1456, 1381, 1345, 1292, 1245, 1218, 1180, 1155, 1130, 1072, 992, 921, 800, 756,  $687 \text{ cm}^{-1}$ ; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  4.86 (d, J = 3.8 Hz, 1 H, H-1), 4.24 (t, J = 6.3 Hz, 1 H, H-2), 4.19–4.17 (m, 1 H, H-5), 4.08 (dd, J = 2.3, 6.1 Hz, 1 H, H-3), 4.00-3.96 (m, 2 H, OCH<sub>2</sub>), 3.83 (dd, J = 3.9, 6.1 Hz, 1 H, H-4), 3.72-3.68(m, 2 H, OCH<sub>2</sub>), 3.50–3.39 (m, 2 H, N-CH<sub>2</sub>), 1.51 (s, 3 H, CH<sub>3</sub>), 1.35 (s, 3 H,  $CCH_3$ ), 1.31 (d, J = 6.0 Hz, 3 H,  $CCH_3$ ); <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ ):  $\delta$  109.7 (C(CH<sub>3</sub>)<sub>2</sub>), 98.1 (C-1), 76.2 (C-5), 75.9 (C-3), 69.5 (C-4), 67.6 (C-2), 64.8 (OCH<sub>2</sub>), 51.1 (NCH<sub>2</sub>), 28.1 (CH<sub>3</sub>), 26.2 (CH<sub>3</sub>), 16.7 (CCH<sub>3</sub>); ESI-MS: 296.1 [M+Na]<sup>+</sup>; Anal. Calcd. for C<sub>11</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub> (273.13): C, 48.34; H, 7.01; Found: C, 48.15; H, 7.25.

# 2-Azidoethyl (2,3,4,6-tetra-O-benzyl- $\alpha$ -D-galactopyranosyl)-(1 $\rightarrow$ 2)-3,4-O-isopropylidene- $\alpha$ -D-fucopyranoside (8)

To a solution of compound 2 (2.5 g, 9.15 mmol) and compound 3 (6.4 g, 10.94 mmol) in anhydrous 1,2-(CH<sub>2</sub>Cl)<sub>2</sub>-DMF (30 mL, 5:1 v/v) were added MS 4A (5 g), CuBr<sub>2</sub> (3.6 g, 16.12 mmol), and TBAB (700 mg, 2.17 mmol) and the reaction mixture was allowed to stir at rt under argon for 48 h. The reaction mixture was filtered through a Celite bed and washed with CH<sub>2</sub>Cl<sub>2</sub> (150 mL). The organic layer was successively washed with satd. NaHCO<sub>3</sub> and H<sub>2</sub>O, dried  $(Na_2SO_4)$ , and concentrated under reduced pressure. The crude product was purified over  $SiO_2$  using hexane-EtOAc (5:1) as eluant to give pure compound 8 (5.5 g, 76%). Yellow oil;  $[\alpha]_D^{25} + 2.28$  (c 1.6, CHCl<sub>3</sub>); IR (neat): 3062, 3030, 2979, 2927, 2873, 2144, 2105, 1606, 1480, 1454, 1355, 1339, 1299, 1209, 1158, 1109, 1089, 1068, 1049, 989, 967, 863, 751, 727, 702, 632  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.36–7.22 (m, 20 H, Ar-H), 5.02 (br s, 1 H, H- $1_{\rm B}$ ), 4.94 (d, J = 11.3 Hz, 1 H, PhCH<sub>2</sub>), 4.82–4.78 (m, 3 H, PhCH<sub>2</sub>, H-1<sub>A</sub>), 4.74  $(d, J = 11.7 Hz, 1 H, PhCH_2), 4.64 (d, J = 11.7 Hz, 1 H, PhCH_2), 4.57 (d, J = 11.7 Hz, 1 Hz,$ J = 11.3 Hz, 1 H, PhCH<sub>2</sub>), 4.43 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.40 (d, J = 11.9 Hz, 1 11.9 Hz, 1 H, PhC $H_2$ ), 4.30 (dd, J = 5.6, 7.7 Hz, 1 H, H-5<sub>B</sub>), 4.26–4.23 (m, 1 H, H-3<sub>A</sub>), 4.19–4.18 (m, 1 H, H-5<sub>A</sub>), 4.09–4.07 (m, 3 H, H-2<sub>B</sub>, H-3<sub>B</sub>, H-4<sub>B</sub>), 4.03 (dd, J = 2.4, 5.4 Hz, 1 H, H-4<sub>A</sub>), 3.77–3.71 (m, 2 H, H-2<sub>A</sub>, OCH<sub>2a</sub>), 3.6

### 6 G. Guchhait & A.K. Misra

(t, J = 8.6 Hz, 1 H, OCH<sub>2b</sub>), 3.48–3.43 (m, 2 H, H-6<sub>abB</sub>), 3.31–3.26 (m, 1 H, NCH<sub>2a</sub>), 3.05–3.00 (m, 1 H, NCH<sub>2b</sub>), 1.39 (s, 3 H, CH<sub>3</sub>), 1.32 (d, J = 6.6 Hz, 3 H, CCH<sub>3</sub>), 1.27 (s, 3 H, CCH<sub>3</sub>); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>):  $\delta$  128.8–127.8 (Ar-C), 109.1 (C(CH<sub>3</sub>)<sub>2</sub>), 97.1 (C-1<sub>A</sub>), 96.5 (C-1<sub>B</sub>), 79.2 (C-3<sub>B</sub>), 76.6 (C-2<sub>B</sub>), 76.4 (C-4<sub>B</sub>), 75.3 (PhCH<sub>2</sub>), 75.2 (C-2<sub>A</sub>), 74.8 (C-3<sub>A</sub>), 74.5 (C-4<sub>A</sub>), 73.8 (PhCH<sub>2</sub>), 73.4 (PhCH<sub>2</sub>), 73.0 (PhCH<sub>2</sub>), 69.4 (C-5<sub>B</sub>), 68.5 (C-6<sub>B</sub>), 68.0 (OCH<sub>2</sub>), 64.2 (C-5<sub>A</sub>), 50.8 (N-CH<sub>2</sub>), 28.5, 26.8 (2 CH<sub>3</sub>), 16.7 (CCH<sub>3</sub>); ESI-MS: 818.3 [M+Na]<sup>+</sup>; Anal. Calcd. for C<sub>45</sub>H<sub>53</sub>N<sub>3</sub>O<sub>10</sub> (795.37): C, 67.91; H, 6.71; Found: C, 67.73; H, 6.95.

# 2-Azidoethyl (2,3,4,6-tetra-O-benzyl- $\alpha$ -D-galactopyranosyl)-(1 $\rightarrow$ 2)-4-O-acetyl- $\alpha$ -D-fucopyranoside (9)

A solution of compound 8 (5 g, 6.28 mmol) in 80% aq. AcOH (100 mL) was allowed to stir at  $80^{\circ}$ C for 2 h and the solvents were evaporated and coevaporated under reduced pressure. To a solution of the crude product in anhydrous DMF (10 mL) was added CH<sub>3</sub>C(OC<sub>2</sub>H<sub>5</sub>)<sub>3</sub> (7 mL, 38.18 mmol) followed by p-TsOH (200 mg) and the reaction mixture was allowed to stir at rt for 2 h concentrated under reduced pressure. The crude product was dissolved in 80% aq. AcOH (50 mL) and the reaction mixture was allowed to stir at rt for 30 min. The solvents were removed under reduced pressure and the crude product was purified over  $SiO_2$  using hexane-EtOAc (5:1) as eluant to give pure compound **9** (4.4 g, 88%). Yellow oil;  $[\alpha]_D^{25}$  +8.5 (c 1.6, CHCl<sub>3</sub>); IR (neat): 3414, 3019, 2923, 2400, 2109, 1740, 1454, 1373, 1215, 1094, 1047, 928, 760, 699, 669, cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  7.33–7.25 (m, 20 H, Ar-H), 5.25 (d, J = 2.8 Hz, 1 H, H-4<sub>A</sub>), 4.99 (d, J = 3.7 Hz, 1 H, H-1<sub>B</sub>), 4.91 (d, J =11.6 Hz, 1 H, PhCH<sub>2</sub>), 4.88 (d, J = 3.6 Hz, 1 H, H-1<sub>A</sub>), 4.83 (d, J = 11.7 Hz, 1 H, PhC $H_2$ ), 4.79 (d, J = 11.9 Hz, 1 H, PhC $H_2$ ), 4.74 (d, J = 11.9 Hz, 1 H,  $PhCH_2$ , 4.64 (d, J = 11.9 Hz, 1 H,  $PhCH_2$ ), 4.54 (d, J = 11.6 Hz, 1 H,  $PhCH_2$ ), 4.47 (d, J = 11.9 Hz, 1 H, PhCH<sub>2</sub>), 4.37 (d, J = 12.0 Hz, 1 H, PhCH<sub>2</sub>), 4.21  $(dd, J = 6.5, 12.1 Hz, 1 H, H-5_B), 4.11-4.05 (m, 3 H, H-3_A, H-2_B, H-5_A), 3.99$  $(dd, J = 2.7, 10.1 Hz, 1 H, H-3_B), 3.8 (d, J = 1.8 Hz, 1 H, H-4_B), 3.77-3.73$ (m, 2 H, H-2<sub>A</sub>, H-6<sub>aB</sub>), 3.55–3.51 (m, 3 H, H-6<sub>bB</sub>, OCH<sub>2a</sub>, OH), 3.36–3.30 (m, 1 H, OCH<sub>2b</sub>), 3.28–3.26 (m, 1 H, NCH<sub>2a</sub>), 3.19–3.17 (m, 1 H, NCH<sub>2b</sub>), 2.06 (s, 3 H, COCH<sub>3</sub>), 1.11 (d, J = 6.6 Hz, 3 H, CCH<sub>3</sub>); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>):  $\delta$  171.2 (COCH<sub>3</sub>), 139.0–127.9 (Ar-C), 98.7 (2 C, C-1<sub>A</sub>, C-1<sub>B</sub>), 79.0 (C-2<sub>A</sub>), 78.8 (C-3<sub>A</sub>), 76.7 (C-5<sub>A</sub>), 75.2 (C-4<sub>B</sub>), 75.0 (PhCH<sub>2</sub>), 73.9 (PhCH<sub>2</sub>), 73.8 (PhCH<sub>2</sub>),  $73.5 (C-4_A), 73.2 (PhCH_2), 70.7 (C-5_B), 70.0 (C-6_B), 68.2 (OCH_2), 67.3 (C-3_B), 68.2 (OCH_2), 67.3 (C-3_B), 68.2 (OCH_2), 67.3 (C-3_B), 68.2 (OCH_2), 67.3 (C-3_B), 68.2 (OCH_2), 68.2 (OCH_2),$  $65.8 (C-2_B), 50.9 (NCH_2), 21.2 (COCH_3), 16.6 (CCH_3); ESI-MS: 820.3 [M+Na]^+;$ Anal. Calcd. for C<sub>44</sub>H<sub>51</sub>N<sub>3</sub>O<sub>11</sub> (797.35): C, 66.23; H, 6.44; Found: C, 66.05; H, 6.68.

# 2-Azidoethyl (2,3,5,6-tetra-O-acetyl-β-D-galactofuranosyl)-(1→3)-[(2,3,4,6-tetra-O-benzyl-α-D-galactopyranosyl)-(1→2)]-4-O-acetyl-α-D-fucopyranoside (10)

To a solution of compound 9 (2 g, 2.5 mmol) and compound 4 (1.2 g, 3 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (15 mL) was added MS 4Å (2 g) and the reaction mixture was allowed to stir under argon at rt for 1 h and then cooled to  $-40^{\circ}$ C. To the cold reaction mixture were added *N*-iodosuccinimide (800 mg, 3.55 mmol) and TMSOTf (15  $\mu$ L) and the reaction mixture was allowed to stir at the same temperature for 1 h. The reaction mixture was filtered through a Celite bed and washed with CH<sub>2</sub>Cl<sub>2</sub> (100 mL). The organic layer was successively washed with 5% aq.  $Na_2S_2O_3$ ,  $H_2O$ , dried ( $Na_2SO_4$ ), and evaporated to dryness. The crude product was purified over  $SiO_2$  using hexane-EtOAc (4:1) as eluant to give pure compound **10** (2.3 g, 82%). Yellow oil;  $[\alpha]_D^{25}$  +5.85 (*c* 1.6, CHCl<sub>3</sub>); IR (neat): 3020, 2926, 2857, 2401, 2108, 1744, 1496, 1454, 1372, 1217, 1054, 980, 758, 698, 668 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.33–7.25 (m, 20 H, Ar-H).  $5.43 (m, 1 H, H-5_C), 5.34 (br s, 1 H, H-1_C), 5.18 (d, J = 2.3 Hz, 1 H, H-4_A), 5.09$  $(d, J = 1.4 Hz, 1 H, H-2_C), 5.07 (d, J = 3.2 Hz, 1 H, H-1_B), 5.0 (d, J = 3.2 Hz)$ 1 H, H-1<sub>A</sub>), 4.92 (d, J = 11.4 Hz, 1 H, PhCH<sub>2</sub>), 4.91 (s, 1 H, H-3<sub>C</sub>), 4.80–4.78 (m, 3 H, 3 PhC $H_2$ ), 4.68 (d, J = 11.7 Hz, 1 H, PhC $H_2$ ), 4.53 (d, J = 11.4 Hz, 1 H, PhC $H_2$ ), 4.50 (d, J = 12.1 Hz, 1 H, PhC $H_2$ ), 4.45 (d, J = 12.1 Hz, 1 H, PhCH<sub>2</sub>), 4.39–4.35 (m, 2 H, H-4<sub>B</sub>, OCH<sub>2a</sub>), 4.28–4.21 (m, 3 H, OCH<sub>2b</sub>, H-3<sub>B</sub>, H-3<sub>A</sub>), 4.17–4.14 (m, 3 H, H-2<sub>A</sub>, H-5<sub>A</sub>, H-5<sub>B</sub>), 4.06–4.04 (m, 2 H, H-2<sub>B</sub>, H-4<sub>C</sub>), 3.74–3.72 (m, 1 H, H-6<sub>aC</sub>), 3.56–3.54 (m, 1 H, H-6<sub>abB</sub>), 3.49–3.47 (m, 1 H, H-6<sub>bC</sub>), 3.29–3.28 (m, 1 H, NCH<sub>2a</sub>), 3.14–3.11 (m, 1 H, NCH<sub>2b</sub>), 2.11, 2.06, 2.01, 1.98, 1.96 (5 s, 15 H, 5 COCH<sub>3</sub>), 1.1 (d, J = 6.6 Hz, 3 H, CCH<sub>3</sub>); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 171.0, 170.8, 170.4 (2 C), 169.7 (5 COCH<sub>3</sub>), 139.4–127.6 (Ar-C),  $107.3 (C-1_C), 96.9 (C-1_A), 95.3 (C-1_B), 81.4 (C-2_C), 81.2 (C-4_B), 79.5 (C-2_B), 77.5 (C-2_B),$ (C-3<sub>C</sub>), 76.5 (C-4<sub>C</sub>), 75.7 (C-2<sub>A</sub>), 75.2 (PhCH<sub>2</sub>), 73.9 (C-5<sub>A</sub>), 73.5 (2 C, 2 PhCH<sub>2</sub>), 73.0 (PhCH<sub>2</sub>), 72.9 (C-4<sub>A</sub>), 71.3 (C-5<sub>B</sub>), 70.2 (C-3<sub>B</sub>), 69.8 (2 C, C-3<sub>A</sub>, C-5<sub>C</sub>), 67.5 (C-6<sub>B</sub>), 65.6 (C-6<sub>C</sub>), 62.8 (OCH<sub>2</sub>), 21.2, 21.1, 21.0 (2 C), 20.9 (5 COCH<sub>3</sub>), 16.3 (CCH<sub>3</sub>); ESI-MS: 1150.5 [M+Na]<sup>+</sup>; Anal. Calcd. for C<sub>58</sub>H<sub>69</sub>N<sub>3</sub>O<sub>20</sub> (1127.45): C. 61.75; H, 6.16; Found: C, 61.57; H, 6.40.

# 2-Aminoethyl ( $\beta$ -D-galactofuranosyl)-(1 $\rightarrow$ 3) -[( $\alpha$ -D-galactopyranosyl)-(1 $\rightarrow$ 2)]- $\alpha$ -D-fucopyranoside (1)

A solution of compound **10** (1.5 g, 1.33 mmol) in 0.1 M CH<sub>3</sub>ONa in CH<sub>3</sub>OH (50 mL) was allowed to stir at rt for 5 h and neutralized with Dowex 50W X8 (H<sup>+</sup>) resin. The reaction mixture was filtered and concentrated under reduced pressure. To a solution of the crude product in CH<sub>3</sub>OH-H<sub>2</sub>O-AcOH (20 mL, 10:1:0.1, v/v) was added 20% Pd(OH)<sub>2</sub>-C (100 mg) and the reaction mixture was allowed to stir under a positive pressure of hydrogen at rt for 12 h. The

#### 8 G. Guchhait & A.K. Misra

reaction mixture was filtered through a Celite bed and evaporated to dryness to give compound 1, which was further purified over Sephadex LH-20 column using CH<sub>3</sub>OH-H<sub>2</sub>O (5:1) as eluant (490 mg, 70%). White powder;  $[\alpha]_D^{25}$  +6.24 (c 1.6, CH<sub>3</sub>OH); IR (KBr): 3406, 2936, 2107, 1639, 1433, 1380, 1244, 1073, 1032, 977, cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  5.16 (br s, 1 H, H-1<sub>C</sub>), 5.01 (br s, 1 H, H-1<sub>B</sub>), 5.00 (br s, 1 H, H-1<sub>A</sub>), 4.12–4.04 (m, 4 H, H-2<sub>A</sub>, H-2<sub>C</sub>, H-3<sub>B</sub>, H-3<sub>C</sub>), 4.01–3.97 (m, 3 H, H-2<sub>B</sub>, H-3<sub>A</sub>, H-2<sub>A</sub>), 3.94 (br s, 1 H, H-4<sub>C</sub>), 3.84 (d, *J* = 2.6 Hz, 1 H, H-5<sub>B</sub>), 3.80–3.78 (m, 2 H, H-4<sub>B</sub>, H-5<sub>A</sub>), 3.77–3.70 (m, 4 H, H-5<sub>C</sub>, H-6<sub>aC</sub>, OCH<sub>2</sub>), 3.64–3.60 (m, 2 H, H-6<sub>abB</sub>), 3.52–3.47 (m, 2 H, NCH<sub>2a</sub>, H-6<sub>bc</sub>), 3.33–3.28 (m, 1 H, NCH<sub>2b</sub>), 1.21 (d, *J* = 6.1 Hz, 3 H, CCH<sub>3</sub>); <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD):  $\delta$  107.8 (C-1<sub>C</sub>), 94.5 (C-1<sub>A</sub>), 94.0 (C-1<sub>B</sub>), 81.9 (C-3<sub>C</sub>), 80.0 (C-2<sub>C</sub>), 76.1 (C-3<sub>A</sub>), 73.6 (C-4<sub>C</sub>), 70.3 (C-5<sub>B</sub>), 69.3 (2 C, C-2<sub>A</sub>, C-5<sub>A</sub>), 69.0 (C-3<sub>B</sub>), 68.3 (C-4<sub>B</sub>), 67.7 (C-4<sub>A</sub>), 67.0 (C-5<sub>C</sub>), 64.4 (C-6<sub>C</sub>), 64.0 (C-2<sub>B</sub>), 61.1 (C-6<sub>B</sub>), 59.4 (OCH<sub>2</sub>), 45.3 (NCH<sub>2</sub>), 13.3 (CCH<sub>3</sub>); Anal. Calcd. for C<sub>20</sub>H<sub>37</sub>NO<sub>15</sub> (531.22): C, 45.20; H, 7.02; found: C, 44.95; H, 7.33.

### ACKNOWLEDGEMENTS

G.G. thanks CSIR, New Delhi, for providing a Junior Research Fellowship. This work was supported by Ramanna Fellowship (AKM), Department of Science and Technology, New Delhi (SR/S1/RFPC-06/2006).

### REFERENCES

1. Farmer, J.J. III; Davis, B.R.; Hickman-Brenner, F.W.; McWorther, A.; Huntley-Carter, G.P.; Asbury, M.A.; Riddle, C.; Wathen-Grady, H.G.; Elias, C.; Fanning, G.R.; Steigerwalt, A.G.; O'Hara, C.M.; Morris, G.K.; Smith, P.B.; Brenner, D.J. Biochemical identification of new species and biogroups of Enterobacteriaceae isolated from clinical specimens. J. Clin. Microbiol. **1985**, 21, 46–76.

2. (a) Gavini, F.; Ferragut, B.; Lefebvre, B.; Leclerc, H. Etude taxonomique d' enterobacteries appartenant ou apparentees au genre Enterobacter. *Ann. Microbial. Inst. Pasteur*, **1976**, *127B*, 317–335. (b) Izard, D.; Gavini, F.; Trinel, P.A.; Leclerc, H. Rahnella aqualitis, nouveau membre de la famille des Enterobacteriaceae. *Ann. Microbiol. Inst. Pasteur*. **1979**, *130A*, 163–167.

3. (a) Caroff, N.; Chamoux, C.; Le Gallou, F.; Gavini, F.; Gautreau, D.; Richet, H.; Reynaud, A. Two epidemiologically related cases of *Rahnella aquatilis* bacteremia. *Eur. J. Clin. Microbiol. Infect. Dis.* **1998**, *17*, 349–352. (b) Reina, J.; Lopez, A. Clinical and microbiological characteristics of *Rahnella aquatilis* strains isolated from children. *J. Infect.* **1996**, *33*, 135–137. (c) Maraki, S.; Samonis, G.; Marnelakis, E.; Tselentis, Y. Surgical wound infection caused by *Rahnella aquatilis. J. Clin. Microbiol.* **1994**, *32*, 2706–2708; (d) Stock, I.; Gruger, T.; Wiedemann, B. Natural antibiotic susceptibility of *Rahnella aquatilis* and *R. aquatilis* related strains. *J. Chemother.* **2000**, *12*, 30–39.

4. Zdorovenko, E.L.; Varbanets, L.D.; Zatonsky, G.V.; Ostapehuk, A.N. Structure of the O-polysaccharide of the lipopolysaccharide of *Rahnella aquatilis* 1–95. *Carbohydr*. *Res.* **2004**, *339*, 1809–1812.

5. (a) Roy, R. Ed. *Carbohydrate Based Vaccines*, Oxford University Press, U.S.A. **1998**. (b) Girard, M.P.; Preziosi, M.-P.; Aguado, M.-T.; Kieny, M.P. A review of vaccine

#### Synthesis of Rhanella aquatilis 1–95 9

research and development: meningococcal disease. *Vaccine* **2006**, *24*, 4692–4700; (c) Jansen, W.T.M.; Hogenboom, S.; Thijssen, M.J.L.; Kamerling, J.P.; Vliegenthart, J.F.G.; Verhoef, J.; Snippe, H.; Verheul, A.F. Synthetic 6B di-, tri-, and tetrasaccharideprotein conjugates contain pneumococcal type 6A and 6B common and 6B-specific epitopes that elicit protective antibodies in mice. *Infect. Immun.* **2001**, *69*, 787–793; (d) Pozsgay, V. Synthetic *Shigella* vaccines: a carbohydrate-protein conjugate with totally synthetic hexadecasaccharide haptens. *Angew. Chem. Int. Ed. Engl.* **1998**, *37*, 138–142.

6. (a) Mukherjee, C.; Misra, A.K. Total synthesis of an antigenic heptasaccharide motif found in the cell-wall lipooligosaccharide of *Mycobacterium gordonae* strain 989. *Gly*coconjugate J. **2008**, 25, 611–624. (b) Mandal, P.K.; Misra, A.K. Synthesis of oligosaccharides corresponding to the polysaccharides of *Lactobacillus* and *Thermophilus* strains. *Synthesis* **2007**, 2660–2666; (c) Mandal, P.K.; Misra, A.K. Concise synthesis of two pentasaccharides corresponding to the  $\alpha$ -chain oligosaccharides of *Neisseria gonorrhoeae* and *Neisseria meningitidis*. *Tetrahedron* **2008**, 64, 8685–8691; (d) Mukherjee, C.; Misra, A.K. Total synthesis of a unique tetrasaccharide present in the human clotting factor IX and mammalian Notch 1 receptor. *Tetrahedron Asymm*. **2009**, 20, 473–477; (e) Mukherjee, C.; Misra, A.K. Synthesis of a unique trisaccharide having an acetal linkage between open chain and cyclic sugar found in the cell wall of *Proteus*. *Tetrahedron Asymm*. **2008**, *19*, 2746–2751.

7. Schmidt, O.T. Acetalation. Methods Carbohydr. Chem. 1963, 2, 318–325.

8. Lerner, L.M. A convenient synthesis of 1,2,3,4-tetra-O-acetyl- $\alpha$ -D-fucopyranose from D-galactose. *Carbohydr. Res.* **1993**, 241, 291–294.

9. (a) Chakraborty, A.K.; Gulhane, R. Perchloric acid adsorbed on silica gel as a new, highly efficient, and versatile catalyst for acetylation of phenols, thiols, alcohols and amines. *Chem. Commun.* **2003**, 1896–1897. (b) Misra, A.K.; Tiwari, P.; Madhusudan, S.K. HClO<sub>4</sub>-SiO<sub>2</sub> catalyzed per-O-acetylation of carbohydrates. *Carbohydr. Res.* **2005**, 340, 325–329.

10. Basu, S.; Pal, J.N. Synthesis of methyl 2-O- $\alpha$ -D-galactopyranosyl-3-O- $\beta$ -D-glucopyranosyl- $\alpha$ -D-glucopyranoside, a trisaccharide of the R-1 core antigen of *Enter*obacteriaceae. Carbohydr. Res. **1990**, 208, 241–245.

11. Wolfrom, M.L.; Yosizawa, Z.; Juliano, B.O. Ethyl 1-thio- $\alpha$ -D-galactofuranoside. J. Org. Chem. **1959**, 24, 1529–1530.

12. Sato, S.; Mori, M.; Ito, Y.; Ogawa, T. An efficient approach to O-glycosides through CuBr<sub>2</sub>-Bu<sub>4</sub>NBr mediated activation of glycosides. *Carbohydr. Res.* **1986**, *155*, C6–C10.

13. Garegg, P.J.; Oscarson, S. A synthesis of 8-methoxycarbonyloct-1-yl O- $\alpha$ -D-galactopyranosyl- $(1\rightarrow 3)$ -O- $\beta$ -D-galactopyranosyl- $(1\rightarrow 4)$ -2-acetamido-2-deoxy- $\beta$ -D-glucopyranoside. *Carbohydr. Res.* **1985**, *136*, 207–213.

14. Veeneman, G.H.; van Leeuwen, S.H.; van Boom, J.H. Iodonium ion promoted reactions at the anomeric centre. II. An efficient thioglycoside mediated approach toward the formation of 1,2-*trans* linked glycosides and glycosidic esters. *Tetrahedron Lett.* **1990**, *31*, 1331–1334.

15. Konradsson, P.; Udodong, U.E.; Fraser-Reid, B. Iodonium promoted reactions of disarmed thioglycosides. *Tetrahedron Lett.* **1990**, *31*, 4313–4316.

16. Marino, K.; Baldoni, L.; Marino, C. Facile synthesis of benzyl- $\beta$ -D-galactofuranoside. A convenient intermediate for the synthesis of D-galactofuranose containing molecules. *Carbohydr. Res.* **2006**, *341*, 2286–2289.

17. Pearlman, W.M. Noble metal hydroxides on carbon nonpyrophoric dry catalysts. *Tetrahedron Lett.* **1967**, *8*, 1663–1664.