

Facile new method for synthesizing *N*-polyfluoroalkylated heterocycles – molecular structure of *N*-(bromodifluoromethyl)-4-dimethylaminopyridinium bromide

Alexander Kolomeitsev,^a Ralph-Matthias Schoth,^b Enno Lork^b and Gerd-Volker Rösenthaller^{*b}

^a Institute of Organic Chemistry, Ukrainian Academy of Sciences, Murmanskaya 6, 253660 Kiev-94, Ukraine

^b Institut für Anorganische und Physikalische Chemie, Universität Bremen, Leobener Strasse, D-28334 Bremen, Germany

Carbon–bromine bond cleavage is observed when 4-dimethylaminopyridine is treated with CF₂Br₂ and BrCF₂CF₂Br to yield *N*-(bromodifluoromethyl) and *N*-(2-bromo-1,1,2,2-tetrafluoromethyl)pyridinium bromides, **1-Br** and **2-Br**, which are reductively debrominated using Bu₃SnH and fluorinated by anhydrous Me₄N⁺F[−]; the molecular structure of **1-Br** is determined by single crystal X-ray crystallography to reveal a partial quinoidal character in the pyridine system and a hypervalent Br[−]⋯BrCF₂ pairing.

The introduction of polyfluoroalkyl groups into organic molecules is of considerable importance in medicine and agrochemistry.¹ To the best of our knowledge neither the polyfluoroalkylation of nitrogen-containing heterocycles nor a successful displacement of bromine in CF₂Br₂ using a nitrogen nucleophile has been reported.^{2,3,4} The reaction of 4-(dimethylamino)pyridine (DMAP) towards fluorinated organyls has not been studied to a great extent, only hexafluorobenzene and 1,2-dichloro-tetrafluorocyclobutene fluorine and chlorine have been substituted to form hexakis- or bispyridinium salts.^{5,6} Here we report some preliminary results on a new versatile synthesis of the first *N*-polyfluoroalkylated *N*-heterocycle starting from dibromodifluoromethane, 1,2-dibromotetrafluoromethane and DMAP.

In polar solvents, like DMF and acetonitrile the reaction (Scheme 1) was completed within 1 h after adding small amounts of activated zinc or copper forming the colourless pyridinium bromides, **1-Br** and **2-Br**.^{†‡} When *p*-dinitrobenzene was added to the BrCF₂CF₂Br/DMAP mixture no product was observed. This was probably due to a single-electron transfer (SET) process initiating an S_{RN}1 sequence,⁷ whereas in the case of CF₂Br₂/DMAP no influence due to a possible difluorocarbene-mediated ionic chain mechanism was observed.⁸ No reaction was observed in diethylether, THF or diglyme.

Using tributylstannane,⁹ compounds **1-Br** and **2-Br** were hydrogenated to give the colourless *N*-difluoromethylated and *N*-(1,1,2,2-tetrafluoro)ethylated pyridinium salts,[§] **3-Br** and **4-Br** respectively (Scheme 1). The BrCF₂ group was successfully fluorinated with anhydrous tetramethylammonium fluoride¹⁰ to furnish *N*-trifluoromethyl-4-dimethylamino-pyri-

dium bromide which was converted to the respective tetrafluoroborate **5-BF₄**.¶

The single crystal X-ray structure determination of **1-Br** (Fig. 1)¶ showed an almost planar six-membered ring with an 8.9° deviation of the exocyclic C(1)–N(1) bond. An angle of 10.6° was observed between the planes through C(3)–C(4)–C(5) and N(2)–C(7)–C(8). There was a substantial degree of quinoidal character^{5,6} in the pyridinium systems, since the bond length C(2)–C(3) and C(5)–C(6) was significantly shorter (134 pm) than the C(3)–C(4) or C(4)–C(5) distance (143 pm). The bond length C(4)–N(2) (134 pm) was between a carbon–nitrogen single and double bond (147 and 127 pm respectively),⁶ indicating significant conjugation. There was not much evidence for an sp³ lone pair neither at N(1) (sum of the angles 359.2°) nor at N(2) (sum of the angles 358.9°).





Fig. 1 Crystal structure of **1** with thermal ellipsoids. Selected bond distances (pm) and angles (°): C(1)–Br(1) 191.5(7), C(1)–F(1) 135.6(8), C(1)–N(1) 145.4(9), C(2)–N(1) 137.0(8), C(2)–C(3) 134.3(10), C(3)–C(4) 143.5(9), C(4)–N(2) 133.7(8), C(7)–N(2) 145.9(7); Br(1)–C(1)–F(1) 110.0(4), F(1)–C(1)–F(2) 107.3(5), Br(1)–C(1)–N(1) 112.3(4), C(2)–N(1)–C(6) 119.7, C(2)–C(3)–C(4) 121.5(6), C(3)–C(4)–C(5) 116.3(5), C(7)–N(2)–C(8) 117.2(5).



Scheme 1

Surprisingly the Br[−](2)⋯Br(1a)CF₂ distance was 315.7 pm, much less than the sum of the Van der Waals radii (370 pm) accounting for a hypervalent ion-pairing,¹¹ whereas the distance Br[−](2)⋯N(2)Me₂ was 361.2 pm. The Br(1)–C bond in **1-Br** seems to be unaffected by the close distance of the Br[−] anion since similar values were found in CF₂BrC(O)NH₂ and BrCF₂CF₂Br, 190(2) and 192.0 pm respectively.^{12,13}

Footnotes

† All reactions were performed under nitrogen in carefully dried solvents. All new compounds gave satisfactory elemental analyses. Mass spectra were recorded under DCI-positive conditions (reactand gas NH₃), NMR spectra at 80.13 (¹H, TMS) and 75.39 MHz (¹⁹F, CClF₃).

‡ *Synthesis of 1-Br and 2-Br*: To a solution of DMAP (10 mmol) in MeCN (10 ml) and Cu powder (5 μm), CF₂Br₂ (3.14 g, 15 mmol) or C₂F₄Br₂ (3.90 g, 15 mmol) were added. After 1 h at ambient temperature the solid formed was filtered, washed with diethylether (2 × 10 ml) and recrystallized from MeCN. Yields: **1-Br** 2.50 g (75%) (mp 233 °C), **2-Br** 3.00 g (80%) (mp 261 °C). *Selected data for 1*: ¹H NMR δ_H 3.2 (CH₃, 6 H), 6.34, 7.78 (AB system, CH, 4 H, J_{AB} 7.7 Hz); ¹⁹F NMR δ_F −36.8. IR ν/cm^{−1} (KBr): 3450, 3035 and 1649. MS (%): 253 (⁸¹Br, M⁺, 20), 239 (⁸¹Br, M⁺ CH₂, 50), 123 (C₇H₁₁N₂⁺, 100). For **2**: ¹⁹F NMR δ_F −68.6 (CF₂Br), −98.7 (CF₂, ³J_{FF} 4.2 Hz). MS (%): 303 (⁸¹Br, M⁺, 30), 123 (C₇H₁₁N₂⁺, 100).

§ *Synthesis of 3-Br and 4-Br*: To a solution of **1-Br** (1.60 g, 5 mmol) or **2-Br** (1.90 g, 5 mmol) in THF (10 ml), BuSnH (1.70 g, 6 mmol) was added and the solution stirred for 2 h at 50 °C. The solution was then filtrated and the remaining solid washed with diethylether (2 × 10 ml) and recrystallized from MeCN. Yields: **3-Br** 1.12 g (95%) (mp 140 °C); **4-Br** 1.32 g (88%) (mp 171 °C). *Selected data for 3*: ¹H NMR δ_H 8.4 (CF₂H, ²J_{FH} 59.3 Hz); ¹⁹F NMR δ_F −98.1. MS (%): 173 (M⁺, 60), 123 (C₇H₁₁N₂⁺, 100). For **4**: ¹H NMR δ 7.2 (CF₂H, ²J_{FH} 52.4, ³J_{FH} 3.1 Hz); ¹⁹F NMR δ_F −101.8 (CF₂, ³J_{FF} 6.3 Hz), −137.2 (CF₂H). MS (%): 223 (M⁺, 100).

¶ *Synthesis of 5-BF₄*: To a solution of **1-Br** (3.32 g, 10 mmol) in CH₂Cl₂ (15 ml), Me₄N⁺F[−] (2.00 g, 21.4 mmol) was added and stirred for 12 h at ambient temperature. The solvent was pumped off and the residue dissolved in MeOH (10 ml) and NH₄BF₄ (20 mmol) added. After filtration the solvent was removed and the solid recrystallized from THF–Et₂O (1 : 1). Yield **5-BF₄** 1.12 g (40%) (mp 210 °C). *Selected data for 5*: ¹⁹F NMR δ_F −61.0. MS (%): 191 (M⁺, 100).

|| *Crystal data for 1-Br*: C₈H₁₀Br₂F₂N₂, *M* = 332.0 orthorhombic, space group *P*2₁2₁2₁, *a* = 575.2(2), *b* = 1097.3(2), *c* = 1752.7(2) pm, *V* = 1106.3(3) Å³, *Z* = 4, *D_c* = 1.993 g cm^{−3}, λ(Mo–Kα) = 0.71073 Å,

Siemens R3 m/v diffractometer 2θ-scan type, 5.0° ≤ 2θ ≤ 55.0°, 173 K; 10357 reflection collected, 2560 independent reflection (*R*_{int} = 1.86%), observed reflections 1875 [*F* > 4.0σ(*F*)]; full-matrix least-squares refinement [Siemens SHELXTL PLUS (VMS)], direct methods, hydrogen atoms were treated as riding models *wR* = 3.27% (observed data). Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Information for Authors, Issue No. 1.

References

- 1 J. T. Welch and S. Eswarakrishnan, *Fluorine in Bioorganic Chemistry*, Wiley, New York, 1991; *Organofluorine Compounds in Medicinal Chemistry and Biomedical Applications*, ed. R. Filler, Y. Kobayashi and L. Yagulpolskii, Elsevier, Amsterdam, 1993.
- 2 M. J. Silvester, *Adv. Heterocycl. Chem.*, 1995, **59**, 1.
- 3 For C-perfluoroalkylated pyridines, see e.g. L. Strekowski, M. Hojjat, S. E. Patterson and A. S. Kiselyov, *J. Heterocycl. Chem.*, 1994, **31**, 1413; W.-Y. Huang, J.-Lin and J. Li, *J. Fluorine Chem.*, 1995, **71**, 51.
- 4 C. Wakselman and A. Lantz, *Organo-fluorine Chemistry (Principles and Commercial Applications)*, ed. R. E. Banks, B. E. Smart and J. C. Tatlow, Plenum Press, New York 1994, pp. 177.
- 5 R. Weiß, B. Pomrehn, F. Hampel and W. Bauer, *Angew. Chem.*, 1995, **107**, 1446.
- 6 A. S. Koch, A. S. Feng, T. A. Hopkins and A. Streitwieser, *J. Org. Chem.*, 1993, **58**, 1409.
- 7 C. Wakselman, *J. Fluorine Chem.*, 1992, **59**, 367.
- 8 C. Wakselman and C. Kaziz, *J. Fluorine Chem.*, 1986, **33**, 347.
- 9 M. Van der Puy, R. K. Belter, R. J. Borowski, L. A. S. Ellis, P. J. Persichimi, A. J. Poss, T. P. Rygas and H. S. Tung, *J. Fluorine Chem.*, 1995, **71**, 59.
- 10 A. A. Kolomeitsev, F. U. Seifert and G.-V. Röschenthaler, *J. Fluorine Chem.*, 1995, **71**, 47.
- 11 R. Weiß, M. Rechinger and F. Hampel, *Angew. Chem.*, 1994, **106**, 901.
- 12 B. Kalyanaraman, L. D. Kispert and J. L. Atwood, *Acta Crystallogr., Sect. B.*, 1978, **34**, 1131.
- 13 G. S. Pawley and E. Whitley, *Acta Crystallogr., Sect. C*, 1988, **44**, 1249.

Received, 5th October 1995; Com. 5/06530E