

Article

Subscriber access provided by Fudan University

Exploring Site Selectivity of Iridium Hydride Insertion into Allylic Alcohols: Serendipitous Discovery and Comparative Study of an Organic and an Organometallic Catalysts for the Vinylogous Peterson Elimination

Houhua Li, Daniele Fiorito, and Clément Mazet

ACS Catal., Just Accepted Manuscript • DOI: 10.1021/acscatal.6b03376 • Publication Date (Web): 17 Jan 2017 Downloaded from http://pubs.acs.org on January 17, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Catalysis is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Exploring Site Selectivity of Iridium Hydride Insertion into Allylic Alcohols: Serendipitous Discovery and Comparative Study of an Organic and an Organometallic Catalysts for the Vinylogous Peterson Elimination

Houhua Li[‡], Daniele Fiorito[‡], Clément Mazet*

Department of Organic Chemistry, University of Geneva, 30 quai Ernest Ansermet, 1211 Geneva, Switzerland.

ABSTRACT: The vinylogous Peterson elimination of a broad range of primary, secondary and tertiary silylated allylic alcohols by two distinct and complementary catalytic systems - a cationic iridium complex and a Brønsted acid - is reported. These results are unexpected. Non-silylated substrates are typically isomerized into aldehydes and silylated allylic alcohols into homoallylic alcohols with structurally related iridium complexes. Although several organic acids and bases are known to promote the vinylogous Peterson elimination, the practicality, mildness, functional group tolerance and generality of both catalysts are simply unprecedented. Highly substituted C=C bonds, stereochemically complex scaffolds, vicinal tertiary and quaternary (stereo)centers are also compatible with the two methods. Both systems are stereospecific and enantiospecific. After optimization, a vast number of dienes with substitution patterns that would be difficult to generate by established strategies are readily accessible. Importantly, control experiments secured that traces of acid that may be generated upon decomposition of the in situ generated iridium hydride are not responsible for the activity observed with the organometallic species. Upon inspection of the reaction scope and on the basis of preliminary investigations, a mechanism involving iridium-hydride and iridium-allyl intermediates is proposed to account for the elimination reaction. Overall, this study confirms that site selectivity for [Ir–H] insertion across the C=C bond of allylic alcohols is a key parameter for the reaction outcome. **KEYWORDS:** *iridium catalysis, Bronsted acid catalysis, vinylogous Peterson elimination, dienes, selective catalysis.*

■ INTRODUCTION

The 1,2-insertion of an olefin into a transition metal hydride (i.e. migratory insertion) is a fundamental elementary step in organometallic chemistry that constitutes the basis of a plethora of catalytic processes.¹ Siteselectivity of insertion is primarily dictated by the nature and the size of the substituents of the C=C bond. Modification of the catalyst structure allows to either alter or best override the inherent electronic and steric biases imposed by the olefin substituents and, consequently, to change the outcome of the transformation. Therefore, gaining understanding on the parameters that control site selectivity of [M-H] insertion across a C=C bond provides a means to elaborate improved catalyst structures (more reactive, more selective) or to access unconventional reactivity patterns by deviating transient intermediates towards novel catalytic manifolds.

In recent years, we and others have pursued the development of late transition metal catalysts for the selective isomerization of allylic and alkenyl alcohols into the corresponding carbonyl derivatives.²⁻⁴ Specifically, we have shown that iridium complexes of general formula [(P,N)Ir(cod)]BAr_F were competent candidates for the isomerization of primary allylic alcohols. Once activated by molecular hydrogen and after degassing of the solution, these typical hydrogenation catalysts can be diverted from their initial task and favor exclusive isomerization into aldehydes instead.⁵ The Pfaltz modified version of Crabtree's catalyst 1 was first established as a very general catalyst for the non-asymmetric version of the reaction, operating under unusually mild reaction conditions.^{5a,6,7} Subsequently, highly enantioselective variants of this reaction using prochiral 3,3-disubstituted allylic alcohols and catalyst-controlled diastereoselective isomerizations of stereochemically complex steroid derivatives have been developed using chiral catalysts such as 2 (Figure 1).⁵

Isotopic labeling experiments have shed light on an unconventional intermolecular hydride-type mechanism involving migratory insertion of the in situ generated [Ir–H] intermediates across the C=C bond of the substrate. Productive isomerization proceeds via insertion of iridium at C₂, followed by β -hydride elimination and tautomerization to deliver the carbonyl compound. The collective results gathered over several years of investigation indicate that substrate polarity clearly influences site se-



Figure 1. (A) Prototypical iridium precatalysts for the selective isomerization of primary allylic alcohols. (B) Influence of site-selectivity of [Ir-H] insertion in allylic alcohols isomerizations. (C) Vinylogous Peterson elimination.

lectivity of migratory insertion.^{2g,5} Formation of homoallylic alcohols - which implies iridium insertion at C₃ - has been observed only with alkyl/alkyl 3,3-disubstituted allylic alcohols and not with aryl/alkyl3,3-disubstituted substrates. Adventitious traces of water were also found to potentially influence site selectivity of migratory insertion and lead to the formation of homoallylic alcohols. Finally, competing E/Z isomerization – which also operates by insertion at C₃ – has been observed in several occasions (Figure 1, B).

In this report, we describe how attempts to tune siteselectivity of migratory insertion by introduction of a silyl substituent in the vicinity of the C=C bond of the allylic alcohol led to the serendipitous discovery of a cationic iridium catalyst for a vinylogous Peterson elimination (Figure 1, C). In addition, control experiments also showed that $[H(OEt_2)_2]BAr_F$ (noted $HBAr_F$), a Brønsted acid, effects this transformation in a complementary manner.⁸ Optimization of this rather underdeveloped reaction was pursued with both catalytic systems to furnish a variety of 1,3-dienes in excellent yields under particularly mild reaction conditions.9 Overall, olefinic substitution patterns that would be otherwise difficult to prepare and compatibility with functional groups that would not be tolerated by the more conventional reagents used for this transformation are important assets of these methods. Finally, preliminary investigations suggest that the iridium catalyst operates via an unorthodox mechanism, distinct from those previously reported for Peterson 1,4-elimination, while HBAr_F likely functions via a more traditional carbocationic pathway.^{9b,e,10}

RESULTS AND DISCUSSION

Exploring site selectivity of [Ir-H] insertion in the isomerization of allylic alcohols. The *β*-silicon effect refers to the ability of a silvl group to stabilize a developing positive charge at a carbon atom in the beta position by hyperconjugation; a property that has been judiciously employed in a variety of transformations in organic synthesis.^{10,11} At the outset of our investigations, we were intrigued by the potential influence a β -silvl group may exert on site-selectivity of [Ir-H] insertion in the C=C bond of allylic alcohols. Therefore, the representative silylated substrates 4a and 4b (which possess an aryl and an alkyl group at C₃ respectively) were subjected to the typical reaction conditions for isomerization of allylic alcohols using achiral complexes 1 and 3.^{5a,12,13} While isomerization of 4a with 1 produced aldehyde 5a exclusively (32% yield), isomerization of (*Z*)-3-cyclopentyl-4-(trimethylsilyl)but-2enol 4b afforded homoallylic alcohol 7b as the sole product of the reaction (eqs (1) and (3), Figure 2). Unexpectedly, when subjecting 4a and 4b to the same experimental protocol using 3, exclusive formation of 1,3-dienes 6a and 6b by a formal 1,4 Peterson elimination was observed (>99% conv., 91% and 82% yield respectively) (eqs (2) and (4), Figure 2). Consistent with our previous investigations, isomerization of 4c and 4d – two closely related allylic alcohols devoid of a β -silyl group – furnished quasiquantitatively aldehydes 5c and 5d – both with complex 1 and 3 (eqs (5) and (6), Figure 2). Collectively, these results indicate that if indeed the presence of a silyl group in the vicinity of the C=C bond of the substrate strongly influences site selectivity of [Ir-H] insertion; the outcome of the reaction is also heavily depending on the structure of the catalyst employed. This latter observation is more surprising as both 1 and 3 behave similarly for nonsilylated 3,3-disubstituted alkyl/alkyl and alkyl/aryl primary allylic alcohols.

Interrogating the specificity of 3 for the vinylogous Peterson elimination. Intrigued by the mildness of the reaction conditions and the selectivity with which 3 converted **4a-b** into **6a-b**, we set out to test the potential of this precatalyst with a substrate that in principle should not be compatible with the typical Brønsted and Lewis acid catalysts/reagents known to effect Peterson 1,4eliminations. Therefore, both geometrical isomers of an N-Boc-protected phenylalanine derived silvlated primary allylic alcohol (*E*)-**4e** and (*Z*)-**4e** were evaluated with a set

1

2

3

4

5

6

7

8 9

10

11

12

13 14

15

16

ACS Catalysis



Figure 2. Comparative reactivity of achiral catalysts 1 and 3 for the isomerization of aryl and alkyl containing silylated primary allylic alcohols (eqs. (1)-(4)). Comparison between 1 and 3 for the isomerization of model aryl and alkyl-containing primary allylic alcohols (eqs. (5) and (6)).

Table 1. Reaction optimization^{*a*}

Me₃Si				II
	OH -	THF, 23 °C, 24 h	► ┌ू	
	NHBoc	,,		NHBoc
	rac-(E)- 4e rac-(Z)- 4e			rac-6e
	. 1 .	1		$(\alpha, \lambda)^{k}$
entry	catalyst	substrate	mol %	conv. (%)*
1	ZnCl ₂	rac-(E)- 4e	5	nr
2	ZnCl ₂	rac-(E)- 4e	100	dec. ^c
3	HCl	rac-(E)- 4e	5	nr
4	HOTs	rac-(E)- 4e	5	nr
5	HCl	rac-(E)- 4e	100	~35 ^d
6	$[Ir(cod)_2]BAr_F$	rac-(E)- 4e	5	~40 ^d
7	$[Ir(cod)_2]BAr_F$	rac-(Z)- 4e	5	dec. ^c
8	3	rac-(E)- 4e	5	>99 (92)
9	3	rac-(E)- 4e	5	nr ^e
10	3	rac-(Z)- 4e	5	12
11	3	rac-(Z)- 4e	10	>99 (81)
12	$[H(OEt_2)_2BAr_F]$	rac-(E)- 4e	5	>99 (82) ^f
13	$[H(OEt_2)_2BAr_F]$	rac-(Z)- 4e	5	>99 (82) ^f

^{*a*} Reaction conditions: **4e** (0.1 mmol). Catalyst **3** was activated by molecular hydrogen unless otherwise noted. ^{*b*} Conversion into **6e** determined by ¹H NMR of the crude reaction mixture. In parenthesis, yield of isolated product after purification by column chromatography. ^{*c*} Decomposition of the substrate. ^{*d*} Along with an intractable mixture of degradation products. ^{*e*} Without activation by molecular hydrogen. ^{*f*} After **4** h.

of representative catalysts and reagents (Table 1). When used in catalytic amount ZnCl₂, HCl and HOTs did not display any reactivity (entry 1-5). A stoichiometric amount of ZnCl₂ only led to decomposition of (*E*)-**4e**, while a 100 mol% of HCl gave ca. 35% of **6e** along with an intractable mixture of side-products. A commercially available cationic iridium source ([Ir(cod)2]BAr_F) was also tested. Product formation was observed with (*E*)-**4e** (ca. 40% along with degradation products), and only decomposition of the substrate was noted with (*Z*)-**4e** (entry 6-7). Much to our satisfaction, with complex **3**, both geometrical isomers of allylic alcohol **4e** were quantitatively converted into **6e**, thus underlying the unique character of this catalyst to effect a 1,4 Peterson elimination on a sensitive substrate. Noticeably, while the use of only 5 mol% of **3** afforded **6e** in 92% yield, 10 mol% were required to observe a similar outcome (81% yield). This different reactivity between (*E*)-**4e** and (*Z*)-**4e** highlights the stereospecific nature of the elimination reaction when catalyzed by complex **3** (entry 8-12).

The decomposition of the catalytically competent iridium hydrides $[(P,N)Ir(H)_2(solv)_2]BAr_F$ generated upon activation of 1 (or related structures) by molecular hydrogen has been studied in details over the last 40 years.^{14,15} Typically, di-, tri- and tetranuclear polyhydrido-iridium clusters are generated upon aggregation of the coordinatively unsaturated cationic dihydride intermediates. As this process is formally accompanied by the liberation of one equivalent of HBAr_F, the behavior of a catalytic amount (5 mol%) of this peculiar Brønsted acid was also evaluated using allylic alcohol **4e** (Table 1, entry 12-13).¹⁶

In contrast to HCl or HOTs but also to **3**, after only 4 h quantitative conversion into **6e** was observed starting indifferently from (*E*)-**4e** or (*Z*)-**4e** (8_2 % yield in both cases).



Figure 3. Isomerization of silylated allylic alcohol **4e** catalyzed by **1**.



Figure 4. Substrates syntheses. (A) General route to geometrically pure silylated primary allylic alcohols. (B) Substrate diversity for primary allylic alcohols. ^{*a*} Catalyst used for the Negishi cross-coupling: [(PPh₃)4Pd] (1-5 mol%). ^{*b*} Catalyst used for the Negishi cross-coupling: [(PPh₃)2PdCl₂] (1-5 mol%). ^{*c*} Enantiopurity not determined. ^{*d*} Catalytic system used for the Negishi cross-coupling: [Pd(OAc)₂] (5 mol%)/CPhos (10 mol%). ^{*e*} Prepared via the corresponding enol triflate according to Frantz's protocol (see ref 19). (C) Synthesis of silylated secondary and tertiary allylic alcohols.

Of important note, at this stage of our studies, the distinct behaviors of 3 and HBAr_F toward both geometrical isomers of substrate 4e already point to potential mechanistic differences between the two catalytic systems. It also suggests that liberated HBAr_F upon catalyst decomposition is not responsible for the catalytic activity observed with 3 (vide infra), and clearly eliminates the possibility of hidden Brønsted acid catalysis.¹⁶ In line with the isomerization of 4b (Figure 2 eq. (3)), when rac-(E)-4ewas subjected to catalysis with 1 after activation by molecular hydrogen, homoallylic alcohol (Z)-7e was isolated as the sole product of the reaction. No reaction was observed with rac-(Z)-4e (Figure 3). These results further underscore the distinct reactivity profile of both iridium catalysts with silvlated allylic alcohols despite their seemingly related structures.

Modular synthesis of silylated allylic alcohols. Having established the ability of 3 and HBAr_F in effecting vinylogous Peterson elimination on a particularly sensitive substrate, we sought to explore and delineate the scope and limitations of both catalysts in a systematic comparative study. To ensure maximum efficiency and modularity for substrates synthesis, we adopted a strategy similar to the one recently followed in our laboratories for the preparation of steroidal allylic alcohols and which takes inspiration from protocols developed by Tanabe and co-workers on simple precursors (Figure 4).^{5h,17} The pivotal 1,3ketoesters ga-r were either commercially available or prepared in one step from the corresponding carboxylic acid.¹⁸ The (*E*)-configured enol tosylates (*E*)-10a-b,e-f,k-o were obtained after treatment with triethylamine, Nmethylimidazole and tosyl chloride (1.5 equiv. each). Ketoesters **9a-e**, **j-g**,**l-m**,**o-r** were stereoselectively converted

ACS Catalysis



Figure 5. Substrate scope (0.1-0.2 mmol scale). Isolated yields after column chromatography. ^{*a*} 1 mol% of **3**. ^{*b*} 10 mol% of **3**. ^{*c*} 4 h. ^{*d*} Enantiospecificity not determined.

to (*Z*)-10a-e,j-g,l-m,o-r in moderate to good yields using 5.0 equiv. of LiCl and otherwise identical reaction conditions. Pd-catalyzed stereo-retentive Negishi crosscouplings using in situ generated (trimethylsilyl)methylzinc chloride followed by enoate reduction with di-isobutyl aluminum hydride delivered the primary allylic alcohols in geometrically pure form and good yields over these two steps.¹⁹ Overall, our synthetic strategy enabled to rapidly assemble a collection of 24 different silylated primary allylic alcohols with a high level of structural diversity and molecular complexity (polycyclic, tertiary and quaternary α -stereocenters) along with an array of functional groups (aryl, perfluoroaryl, alcohol, ether, thioether, silyl ether, acetal, N- heterocycles, protected amine, alkene). Substrate 4s which features a tetrasubstituted C=C bond was prepared according to the same sequence after simple C-alkylation of 9g into 9s. Both geometrical isomers of secondary allylic alcohol 11m were obtained starting from (*E*)-**4m** and (*Z*)-**4m** by oxidation to the corresponding α , β -unsaturated aldehydes followed by 1,2-addition of methylmagnesium bromide. The corresponding tertiary allylic alcohol (*E*)-**13m** was obtained in 71% yield by treatment of enoate (*E*)-**12m** with 3 equivalents of MeLi. Similar sequences gave access to (*Z*)-**11s** and (*Z*)-**13s** a secondary and a tertiary allylic alcohol respectively, both featuring a tetrasubstituted C=C bond.

Comparative study between 3 and $HBAr_F$ for the vinylogous Peterson elimination. The complete collection of primary, secondary and tertiary allylic alcohols was subsequently subjected to the prototypical reaction conditions developed for the vinylogous Peterson elimination catalyzed by 3 (5 mol%, activation with molecular hydrogen, room temperature, 24 h) and $HBAr_F$ (5 mol%, room temperature, 4 h). The vast majority of silylated primary allylic alcohols tested proved competent in undergoing a 1,4 elimination reaction with catalyst 3 and

HBAr_F but some important differences were noted for several substrates (Figure 5). Specifically, substrates with either electron-neutral (4a), electron-rich (4g) or electron-poor (4h-i) aryl substituents at C3 were converted quantitatively and isolated in excellent yield with HBAr_F. In contrast, 4h required increased catalyst loading of 3 (10 mol%) to isolate **6h** in an acceptable 59% yield while no reaction was observed with 4i. Similarly, 6i which is characterized by a N-methyl protected indole motif, was isolated in 81% using 10 mol% of 3 while nor modifications of protocol B were necessary to achieve similar performances with $HBAr_F$ (5 mol%, 4 h). The stereospecific nature of the reaction with the iridium precatalyst was again noted in eliminations using (Z)- and (E)-4b, as the latter was reacted with a loading as low as 1 mol%. Interestingly, the optically active silvlated allylic alcohols 4e and 4k - which both possess a tertiary stereocenter adjacent to C3 - underwent the 1,4-Peterson elimination with virtually perfect enantiospecificity and furnished the corresponding chiral products in excellent yields with both catalytic systems. The configurational stability of these stereocenters using 3 is particularly remarkable because isomerization of 4e and 4k with catalyst 1 led to the exclusive formation of the corresponding tetrasubstituted homoallylic alcohols, implying migratory insertion at C₃ and β-H elimination at C4.20 Noticeably, a methionine-derived silvlated allylic alcohol was successfully engaged in the elimination reaction, affording 6f in 44% yield with 3 and 91% with the Brønsted catalyst.²¹ The compatibility of both methods with isolated C=C bonds was demonstrated with substrates 41-n, leading to 61-n in satisfactory to excellent yields (45-95% with 3; 76-97% with $HBAr_F$). Of note, no isomerization of the remote (Z)-configured 1,2disubstituted double bond was noted in the reaction with (Z,Z)- and (E,Z)-4m.²² The presence of a tertiary and more remarkably - of a quaternary stereocenter adjacent to the allylic system does not affect the efficiency of both processes as the stereochemically complex scaffolds 60, 6p and 6q were obtained in excellent yield upon vinylogous elimination of the appropriate allylic alcohols. These results also highlight the compatibility of the metal-based catalyst and the Brønsted acid catalyst with an endocyclic homoallylic alcohol (40), an endocyclic diene system (4p) and a galactose acetonide derivative (4g).^{23,24} The presence of a diene at these strategic positions of the terpene derivatives bodes well for rapid installation and diversification of heterocyclic ring systems via cycloaddition reactions. The ability of complex 3 to perform a high-yielding vinylogous Peterson elimination on a particularly acidsensitive silyl protected derivative such as 4r is simply remarkable (6r: 84% yield) especially because HBAr_F led to the corresponding deprotected secondary alcohol 6r' (78% yield). These contrasted outcomes reinforce the notion that 3 and HBAr_F certainly operate via distinct reaction mechanism and that the catalytic activity of 3 does not result from traces of acid that might have been generated upon catalyst decomposition.¹⁴⁻¹⁶ It is also worth mentioning that a silyl-protected alcohol would certainly not be compatible with the classical basic fluo-

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43 44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60 rine reagents used for vinylogous Peterson eliminations.⁹ Interestingly, (*Z*)-**4s**, a primary allylic alcohol with a tetrasubstituted C=C bond, delivered **6s** in 80-83% yield with **3** and HBAr_F, thus offering a potential complement to Heck-type cross-coupling with allenes or thermolysis of sulfones, two methods which typically deliver related 2,3-disubstituted diene motifs.²⁵

The scope of 1,3-dienes accessible with both protocols was further explored by subjecting secondary and tertiary allylic alcohols (11m,s and 13m,s) to the optimized reaction conditions for the two catalytic systems. All five substrates underwent vinylogous Peterson elimination with the iridium catalyst and afforded the corresponding dienes in excellent yields (82-92%). Noticeably, (*E*,*Z*)-14m was obtained in identical yields starting indifferently from (Z,Z)-11m or (E,Z)-11m, indicating that the metalcatalyzed process is stereoconvergent in nature. The elimination reactions of 11m and 13m led to comparable results when conducted with HBAr_F. The vinylogous Peterson elimination of 115 and 135 shed additional light on the striking differences between the organometallic and the organic catalysts. While with 3, 15s and 18s were isolated in excellent yield (15s: 82%; 18s: 86%), with HBAr_F generation of these 1,2,3- and 1,1',2,3-dienes was accompanied by the formation of the inseparable indenes 16s and 19s. Indene 195 was obtained exclusively when the reaction time was extended, indicating that it is generated by a formal hydroarylation of transient diene 18s.^{16e,1}

Overall, from a synthetic point of view, the diversity of substitution patterns available with both protocols is simply remarkable as it provides direct access to 2-, 2,3-, 1,3-, 1,2,3-, 1,1',3- and 1,1',2,3-substituted dienes. Existing alternatives are at best scarce, limited in scope and often require harsh reaction conditions.²⁵⁻²⁷

Preliminary mechanistic insights. On the basis of the control experiments conducted on substrates (*E*)-**4e** and (*Z*)-**4e** (Table 1) and the results of the vinylogous Peterson elimination conducted on the silylated primary allylic alcohols (*E*)-**4f**, (*Z*)-**4h**, (*Z*)-**4i**, (*Z*)-**40**, (*E*/*Z*)-**4q**, (*Z*)-**4r** and on the silylated secondary allylic alcohols (*Z*)-**11s** and (*Z*)-**13s** (Figure 5), it appears clearly that **3** and HBAr_F operate via different mechanisms.



Figure 6. Proposed mechanism for the vinylogous Peter-

Page 7 of 10

son elimination catalyzed by $HBAr_F$ exemplified with (*Z*)-**135**.

An intuitive proposal that accounts for diene formation when $HBAr_F$ is employed is depicted on Figure 6. It is represented using (*Z*)-**13s** as it also allows to propose a rationale for the obtention of indene **19s** resulting from the hydroarylation reaction.

In contrast, the mode of action of the dihydridoiridium complexes generated upon activation of $\mathbf{3}$ by molecular hydrogen appears less obvious. In an effort to gather preliminary information on the mechanism by which catalyst $\mathbf{3}$ may operate to effect the vinylogous Peterson elimination, a series of additional control experiments was conducted using (*E*)-**4b** as model substrate (Table 2).

Table 2. Control experiments^a



^{*a*} Reaction conditions: (*E*)-**4b** (0.1 mmol), 1 min. activation with H₂ followed by degassing. ^{*b*} Conversion into **6b** determined by ¹H NMR of the crude reaction mixture. In parenthesis, yield of isolated product after purification by column chromatography. ^{*c*} DTBMP = 2,6-di-*tert*-butyl-4-methylpyridine. ^{*d*} TEMPO = (2,2,6,6-Tetramethylpiperidin-1-yl)oxyl.



Figure 7. Tentative mechanism for the vinylogous Peterson elimination catalyzed by **3**.

When the reaction was run in presence of 10 mol% of 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP), a bulky

non-coordinating base, quantitative formation of diene 6b was observed after 24 h, suggesting that potential traces of acid that may be liberated upon decomposition of the iridium dihydride generated by activation with molecular hydrogen are not responsible for catalytic activity.¹⁴⁻¹⁶ In contrast, in the presence of 10 mol% of TEMPO – a notorious trap for transition metal hydrides - no product formation was observed, thus advocating the direct involvement of [Ir-H] intermediates in the operating mechanism for the vinylogous Peterson elimination.^{4c,28} Of additional note, the absence of ring-opened product in the reaction with the cyclopropyl-containing substrate (E)-4n rules out a potential cage-escaped radical mechanism.²⁸ Finally, the complete loss of catalytic activity observed in the presence of 4Å molecular sieves supports the notion that liberated water plays an important role in the overall catalytic process. Conversely, when (*E*)-4b was subjected to catalysis with HBAr_F in the presence of TEMPO (10 mol%) or 4Å molecular sieves, reactivity was still observed (33% conversion and >99% conversion respectively; see SI for details).

Although additional experiments are needed, based on these results and on observations made while investigating the scope of the reaction, we display on Figure 7 a tentative rationale that may account for the vinylogous Peterson elimination catalyzed by 3. Activation of complex 3 by molecular hydrogen generates the catalytically active cationic Ir(III) dihydride A.^{12,13} Upon reaction with the silvlated allylic alcohols, production of the cationic iridium-allyl complexes (B and/or C) accompanied by concomitant formation of one molecule of water seems reasonable. The decreased or absence of reactivity for arylated substrates with a para-electron-withdrawing substituent at C₃ may be explained by the difficulty in accessing or stabilizing intermediate C. Because of the stereospecific nature of the reaction, we believe that iridiumallyl formation is the rate determining step of the catalytic process and that one of the hydride ligands is responsible for activation of the hydroxy functionality of the substrate. The water molecule generated (which may or may not leave the first coordination sphere of the iridium atom) subsequently participates in elimination of the silvl fragment. Intermediate D is postulated to precede a related 6-membered pericyclic transition state leading to the formation of the diene and simultaneous regeneration of Α.

CONCLUSION

In this article, we have reported the serendipitous discovery of two distinct catalysts for the vinylogous Peterson elimination of a variety of silylated allylic alcohols leading to valuable dienes. The first catalyst, a cationic iridium complex supported by a chelating (P,N) ligand, was identified while investigating the site selectivity of [Ir–H] insertion across the C=C bond of silylated allylic alcohols. The second catalytic system, a Brønsted acid (HBAr_F), was discovered when assessing whether traces of acid that might be generated upon decomposition of the active form of the iridium complex were responsible for

catalytic activity. To delineate the synthetic potential of both entities a systematic comparative study was pursued on a vast number of silvlated allylic alcohols. This required the design of a modular sequence based on some of the most recent cross-coupling methods to access the substrates in geometrically pure form with the possibility to introduce up to four substituents on the C=C bond and a representative diversity of functional groups. Overall, both systems were found to react with a variety of primary silvlated allylic alcohols to afford the expected 2- and 2,3-substituted dienes in usually good to excellent yields independently of the extent of substitution of the olefinic moiety, the stereochemical complexity and the congested nature of proximal stereocenters. The two catalysts were found to be both stereo- and enantiospecific. Several sensitive functional groups that would not be compatible with more conventional reagents were also perfectly tolerated by the organic and the organometallic catalyst. Nonetheless, for some specific substrates, clear reactivity differences were noted. For instance, the Brønsted acid catalyst appeared more appropriate to effect the elimination of a methionine derived substrate but it proved not compatible with silyl-protected alcohol functionalities. In the latter case, the iridium catalyst led to the expected diene without cleavage of the acid-sensitive siliconoxygen bond. While all substrates with an aromatic substituent at C₃ were perfectly engaged in the elimination reaction with HBAr_F, a net decrease in reactivity was observed with the iridium complex upon para-substitution with electron-withdrawing substituents. Alkyl containing secondary and tertiary silvlated allylic alcohols were equally reactive with both systems and the corresponding 1,3- and 1,1',3-substituted dienes were isolated in usually high yields. Analogues with an aromatic ring at C₃ were converted smoothly to dienes with the organometallic species while a competing hydroarylation leading to intractable mixtures of products occurred with the organic catalyst.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60 Overall, both synthetic methodologies give access to a broad set of dienes which would be otherwise difficult to prepare with a single and unified experimental protocol. From a practical point of view, the two catalysts effect the vinylogous Peterson elimination under mild reaction conditions and at relatively low loadings. The experimental setup with the Brønsted acid is simpler but the use of the iridium catalyst is clearly preferred for some specific substrates and functional groups.

Aside from practical and synthetic considerations, collectively our results support the notion that both systems follow different catalytic manifolds. First, they clearly demonstrate that traces of acid that may be generated upon catalyst decomposition are not responsible for the activity observed with the iridium catalyst. Additional investigations underscored the key role of the in situ generated iridium-hydride and the liberated water. These control experiments, taken together with the stereospecific nature of the iridium-catalyzed reaction and the lack of reactivity towards substrates with electron-deficient aromatic substituents provided support for a mechanistic scenario involving iridium-allyl intermediates. Comparative study with other iridium catalysts clearly suggest that subtle variations in ligand design strongly influence siteselectivity for [Ir–H] insertion across the C=C bond of silylated allylic alcohols. Further experimental and computational studies into the origin of the reactivity and selectivity of these different catalytic systems are underway in our laboratories.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, characterization of all new compounds and spectral data. This material is available free of charge via the Internet at <u>http://pubs.acs.org</u>.

AUTHOR INFORMATION

Corresponding Author

Prof. Clément Mazet. University of Geneva, Organic Chemistry Department. Quai Ernest Ansermet 30, Geneva 1211 -Switzerland. <u>clement.mazet@unige.ch</u>

Notes

The authors declare no competing financial interests.

Author Contributions

*‡*These authors contributed equally.

ACKNOWLEDGMENT

We thank the University of Geneva and the Swiss National Science Foundation (PPooP2_133482) for financial support and Johnson Matthey for a generous gift of iridium precursors. This article is dedicated to Prof. Alexandre Alexakis for his constant support and friendship.

REFERENCES

(1) (a) Organotransition Metal Chemistry: from Bonding to Catalysis; Hartwig, J. F., Eds; University Science Books: New York, 2010; Chapter 9. (b) The Organometallic Chemistry of the Transition Metals; Crabtree, R. H., Eds; Wiley–VCH: Weinheim, 2014; Chapter 7.

(2) (a) van der Drift, R. C.; Bouwman, E.; Drent, E. J. Organomet. Chem. 2002, 650, 1-24. (b) Uma, R.; Crévisy, C.; Grée, R. Chem. Rev. 2003, 103, 27-52. (c) Fu, G. C. Modern Rhodium-Catalyzed Organic Reactions; Evans, P. A., Ed.; Wiley-VCH: Weinheim, 2005; Chapter 4. (d) Mantilli, L.; Mazet, C. Chem. Lett. 2011, 40, 341-344. (e) Ahlsten, N.; Bartoszewicz, A.; Martin-Matute, B. Dalton Trans. 2012, 41, 1660-1670. (f) Cahard, D.; Gaillard, S.; Renaud, J.-L. Tetrahedron Lett. 2015, 56, 6159-6169. (g) Li, H.; Mazet, C. Acc. Chem. Res. 2016, 49, 1232-1241.

(3) For selected examples of enantioselective isomerization of primary allylic alcohols, see: (a) Botteghi, C.; Giacomelli, G. *Gazz. Chim. Ital.* **1976**, *106*, 1131–1134. (b) Tani, K. *Pure Appl. Chem.* **1985**, *57*, 1845–1854. (c) Tanaka, K.; Qiao, S.; Tobisu, M.; Lo, M. M.-C.; Fu, G. C. *J. Am. Chem. Soc.* **2000**, *122*, 9870–9871. (d) Tanaka, K.; Fu, G. C. *J. Org. Chem.* **2001**, *66*, 8177–8186. (e) Chapuis, C.; Barthe, M.; de Saint Laumer, J.-Y. *Helv. Chim. Acta* **2001**, *84*, 230–242. (f) Li, J.-Q.; Peters, B.; Andersson, P. G. *Chem.-Eur. J.* **2011**, *17*, 1143–11145; (g) Arai, N.; Sato, K.; Azuma, K.; Ohkuma, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 7500–7504. For selected examples of enantioselective isomerization of secondary allylic alcohols, see: (h) Ito, M.; Kitahara, S.; Ikariya, T. *J. Am. Chem. Soc.* **2005**, *127*, 6172–6173. (i) Bizet, V.; Pannecoucke, X.;

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Renaud, J.-L.; Cahard, D. Angew. Chem., Int. Ed. 2012, 51, 6467–6470.

(4) For selected examples of selective isomerization of alkenyl alcohols, see: (a) Ishibashi, K.; Takahashi, M.; Yokota, Y.; Oshima, K.; Matsubara, S. *Chem. Lett.* **2005**, *34*, 664–665. (b) Grotjahn, D. B.; Larsen, C. R.; Gustafson, J. L.; Nair, R.; Sharma, A. J. *Am. Chem. Soc.* **2007**, *129*, 9592–9593. (c) Larionov, E.; Lin, L.; Guénée, L.; Mazet, C. J. Am. Chem. Soc. **2014**, *136*, 16882–16894. (d) Lin, L.; Romano, C; Mazet, C. J. Am. Chem. Soc. **2016**, *138*, 10344–10350. For a relevant review, see: (e) Vasseur, A.; Bruffaerts, J.; Marek, I. Nat. Chem. **2016**, *8*, 209–219.

(5) (a) Mantilli, L.; Mazet, C. *Tetrahedron Lett.* 2009, 50, 4141–4144. (b) Mantilli, L.; Gérard, D.; Torche, S.; Besnard, C.; Mazet, C. *Angew. Chem., Int. Ed.* 2009, 48, 5143–5147. (c) Mantilli, L.; Mazet, C. *Chem. Commun.* 2010, 46, 445–447. (d) Mantilli, L.; Gérard, D.; Torche, S.; Besnard, C.; Mazet, C. *Chem.-Eur. J.* 2010, 16, 12736–12745. (e) Quintard, A.; Alexakis, A.; Mazet, C. *Angew. Chem., Int. Ed.* 2011, 50, 2354–2358. (f) Mantilli, L.; Gérard, D.; Besnard, C.; Mazet, C. *Eur. J. Inorg. Chem.* 2012, 20, 3320–3330. (g) Li, H.; Mazet, C. *Org. Lett.* 2013, 15, 6170–6173. (h) Li, H.; Mazet, C. *J. Am. Chem. Soc.* 2015, 137, 10720–10727.

(6) (a) Crabtree, R. H.; Felkin, H.; Morris, G. E. J. Organomet. Chem. 1977, 141, 205–215. (b) Crabtree, R. H. Acc. Chem. Res. 1979, 12, 331–337.

(7) Wüstenberg, B.; Pfaltz, A. Adv. Synth. Catal. 2008, 350, 174–178.

(8) Brookhart, M.; Grant, B.; Volpe, A. F., Jr. *Organometallics* **1992**, 11, 3920–3922. For a recent use of HBAr_F in Brønsted acid catalysis, see: Chen, Q.-A.; Klare, H. F. T.; Oestreich, M. *J. Am. Chem. Soc.* **2016**, *138*, 7868–7871.

(9) (a) Angoh, A. G.; Clive, D. L. J. Chem. Soc. Chem. Commun.
1984, 534-536. (b) Fleming, I.; Morgan, I. T.; Sarkar, A. K. J. Chem. Soc., Chem. Commun. 1990, 1575-1577. (c) Angell, R.; Parsons, P. J.; Naylor, A.; Tyrrell, E. Synlett 1992, 599-600. (d) Maeta, H.; Suzuki, K. Tetrahedron Lett. 1992, 33, 5969-5972. (e) Fleming, I.; Morgan, I. T.; Sarkar, A. K. J. Chem. Soc., Perkin Trans. 1 1998, 17, 2749-2764. (f) Harmata, M.; Bohnert, G. J. Org. Lett. 2003, 5, 59-61. (g) Ahmed, M.; Atkinson, C. E.; Barrett, A. G. M.; Malagu, K.; Procopiou, A. P. Org. Lett. 2003, 5, 669-672. (h) Borg, T.; Tuzina, P.; Somfai, P. J. Org. Chem. 2011, 76, 8070-8075. (i) Wender, P. A.; Fournogerakis, D. N.; Jeffreys, M. S.; Quiroz, R. V.; Inagaki, F.; Pfaffenbach, M. Nat. Chem. 2014, 6, 448-452. (l) Wender, P. A.; Jeffreys, M. S.; Raub, A. G. J. Am. Chem. Soc. 2015, 137, 9088-9093.

(10) Fleming, I.; Barbero, A.; Walter, D. Chem. Rev. 1997, 97, 2063-2192.

(11) (a) Lambert, J. B. *Tetrahedron* **1990**, *46*, 2677–2689. (b) Lambert, J. B.; Zhao, Y.; Emblidge, R. W.; Salvador, L. A.; Liu, X.; So, J.-H.; Chelius, E. C. *Acc. Chem. Res.* **1999**, *32*, 183–190.

(12) It is now well-established that upon activation by molecular hydrogen of [(P,N)Ir(cod)]BAr_F precatalysts, iridium dihydrides of general formula [(P,N)Ir(H)2(solv)2]BAr_F are generated. For relevant studies, see: (a) Crabtree, R. H.; Davis, M. W. Organometallics 1983, 2, 681–682. (b) Crabtree, R. H.; Davis, M. W. J. Org. Chem. 1986, 51, 2655–2661. (c) Mazet, C.; Smidt, S. P.; Meuwly, M.; Pfaltz, A. J. Am. Chem. Soc. 2004, 126, 14176–14181. (d) Zhu, Y.; Fan, Y.; Burgess, K. J. Am. Chem. Soc. 2010, 132, 6249–6253. (e) Gruber, S.; Neuburger, M.; Pfaltz, A. Organometallics 2013, 32, 4702–4711. (f) Gruber, S.; Pfaltz, A. Angew. Chem., Int. Ed. 2014, 53, 1896–1900. (g) Gruber, S. Organometallics 2016, 35, 699–705. See also ref. 5a and 5b.

(13) During finalization of this study, a related precatalyst was reported. See: Celaje, J. J. A.; Lu, Z.; Kedzie, E. A.; Terrile, N. J.; Lo, J. N.; Williams, T. J. *Nat. Commun.* **2016**, *7*, 11308.

(14) (a) Zhu, Y.; Fan, Y.; Burgess, K. J. Am. Chem. Soc. 2010, 132, 6249–6253. (b) Morris, R. H. J. Am. Chem. Soc. 2014, 136, 1948–1959. (c) Morris, R. H. Chem. Rev. 2016, 116, 8588–8654. (15) (a) Chodosh, D. F.; Crabtree, R. H.; Felkin, H.; Morris, G. E. *J. Organomet. Chem.* **1978**, *161*, C67–C70. (b) Chodosh, D. F.; Crabtree, R. H.; Felkin, H.; Morehouse, S.; Morris, G. E. *Inorg. Chem.* **1982**, *21*, 1307–1311. (c) Smidt, S. P.; Pfaltz, A.; Martínez-Viviente, E.; Pregosin, P. S.; Albinati, A. *Organometallics* **2003**, *22*, 1000–1009. (d) Xu, Y.; Celik, M. A.; Thompson, A. L.; Cai, H.; Yurtsever, M.; Odell, B.; Green, J. C.; Mingos, D. M. P.; Brown, J. M. *Angew. Chem., Int. Ed.* **2009**, *48*, 582–585.

(16) For a short review on hidden Brønsted acid catalysis, see: (a) Taylor, J. G.; Adrio, L. A.; Hii, K. K. Dalton Trans. 2010, 39, 1171-1175. For selected examples, see: (b) Fürstner, A.; Szillat, H.; Gabor, B.; Mynott, R. J. Am. Chem. Soc. 1998, 120, 8305-8314. (c) Schlummer, B.; Hartwig, J. F. Org. Lett. 2002, 4, 1471-1474. (d) Wabnitz, T. C.; Yu, J.-Q.; Spencer, J. B. Chem.-Eur. J. 2004, 10, 484-493. (e) Rhee, J. U.; Krische, M. J. Org. Lett. 2005, 7, 2493-2495. (f) Anderson, L. L.; Arnold, J.; Bergman, R. G. J. Am. Chem. Soc. 2005, 127, 14542-14543. (g) Rosenfeld, D. C.; Shekhar, S.; Takemiya, A.; Utsunomiya, M.; Hartwig, J. F. Org. Lett. 2006, 8, 4179-4182. (h) McBee, J. L.; Bell, A. T.; Tilley, T. D. J. Am. Chem. Soc. 2008, 130, 16562-16571. (i) Kanno, O.; Kuriyama, W.; Wang, Z. J.; Toste, F. D. Angew. Chem., Int. Ed. 2011, 50, 9919-9922. (j) Bolte, B.; Gagosz, F. J. Am. Chem. Soc. 2011, 133, 7696-7699 (k) Dang, T. T.; Boeck, F.; Hintermann, L. J. Org. Chem. 2011, 76, 9353-9361. (1) Bowring, M. A.; Bergman, R. G.; Tilley, T. D. Organometallics 2011, 30, 1295–1298. (m) Schmidt, R. K.; Müther, K.; Mück-Lichtenfeld, C.; Grimme, S.; Oestreich, M. J. Am. Chem. Soc. 2012, 134, 4421-4428.

(17) (a) Nakatsuji, H.; Ueno, K.; Misaki, T.; Tanabe, Y. *Org. Lett.* **2008**, 10, 2131–2134. (b) Manabe, A.; Ohfune, Y.; Shinada, T. *Synlett* **2012**, 23, 1213–1216.

(18) Allan, K. M.; Hong, B. D.; Stoltz, B. M. Org. Biomol. Chem. 2009, 7, 4960–4964.

(19) Allylic alcohols (*Z*)-**40** and (*Z*)-**4p** were prepared via the corresponding (*Z*)-enol triflates according a protocol developed by Frantz and co-workers: Babinski, D.; Soltani, O.; Frantz, D. E. *Org. Lett.* **2008**, *10*, 2901–2904. See Supporting Information for details.

(20) Isomerization of *rac-*(E)-**4k** with complex **1** led to exclusive formation of the corresponding (Z)-homoallylic alcohol ((Z)-**7k**: 70% yield). See Supporting Information for details.

(21) Enantiospecificity not determined.

(22) Larionov, E.; Li, H.; Mazet, C. Chem. Commun. 2014, 50, 9816–9826.

(23) The endocyclic 1,3-diene system was found to be incompatible during the isomerization of steroidal allylic alcohols using complex 1. For more details see ref 5h. For a relevant ruthenium hydride-promoted isomerization of 1,3-dienes, see: Clark, J. R.; Griffiths, J. R.; Diver, S. T. J. Am. Chem. Soc. 2013, 135, 3327–3330.

(24) Liu, H.-X.; Wu, Q.-P.; Shu, Y.-N.; Chen, X.; Xi, X.-D.; Du, T.-J.; Zhang, Q.-S. *Carbohydr. Res.* **2009**, 344, 2342–2348.

(25) For approaches based on 2,3-dimethylenebutadiene dianion, see: (a) Bates, R. B.; Gordon III, B.; Highsmith, T. K.; White, J. J. J. Org. Chem. **1984**, 49, 2981–2987. For approaches based on thermolysis of sulfones, see: (b) Nakayama, J.; Machida, H.; Saito, R.; Akimoto, K.; Hoshino M. Chem. Lett. **1985**, *14*, 1173–1176. (c) Trost, B. M.; Huang, X Org. Lett. **2005**, *7*, 2097–2099. For approaches based on cross-coupling with allenes, see: (d) Chang, H.-M.; Cheng, C.-H. J. Org. Chem. **2000**, *65*, 1767–1773. For approaches based on cross-enyne metathesis with alkynes, see: (e) Kinoshita, A.; Sakakibara, N.; Mori, M. J. Am. Chem. Soc. **1997**, *119*, 12388–12389. (f) Tonogaki, K.; Mori, M. Tetrahedron Lett. **2002**, *43*, 2235–2238. (g) Diver, S. T.; Giessert, A. J. Chem. Rev. **2004**, *104*, 1317–1382.

(26) (a) *Science of Synthesis*; Trost, B. M.; Rawal, V. H.; Kozmin, S. A., Eds.; Thieme: Stuttgart, 2009; Vol. 46. (b) De

Paolis, M.; Chataigner, I.; Maddaluno, J. *Top. Curr. Chem.* 2012, 327, 87–146.

(27) For selected contributions, see: (a) Aufdermarsh, C. A. J. Org. Chem. 1964, 29, 1994–1996. (b) Nunomoto, S.; Yamashita, Y. J. Org. Chem. 1979, 44, 4788-4791. (c) Nunomoto, S.; Kawakami, Y.; Yamashita, Y. Bull. Chem. Soc. Jpn. 1981, 54, 2831-2832. (d) Brown, P. A.; Bonnert, R. V.; Jenkins, P. R.; Lawrence, N. J.; Selim M. R. J. Chem. Soc., Perkin Trans. 1 1991, 8, 1893-900. (e) Sofia, A.; Karlström, E.; Itami, K.; Bäckvall, J.-E. J. Org. Chem. 1999, 64, 1745-1749. (f) Katritzky, A. R.; Serdyuk, L.; Toader, D.; Wang, X. J. Org. Chem. 1999, 64, 1888-1892. (g) Trost, B. M.; Pinkerton, A. B.; Seidel, M. J. Am. Chem. Soc. 2001, 123, 12466-12476. (h) Shen, Q.; Hammond, G. B. Org. Lett. 2001, 3, 2213-2215. (i) Oh, C. H.; Jung, S. H.; Bang, S. Y.; Park, D. I. Org. Lett. 2002, 4, 3325-3327. (j) Alcaraz, L.; Cox, K.; Cridland, A. P.; Kinchin, E.; Morris, J.; Thompson, S. P. Org. Lett. 2005, 7, 1399-1401. (k) Ebran, J.-P.; Hansen, A. L.; Gøgsig, T. M.; Skrydstrup, T. J. Am. Chem. Soc. 2007, 129, 6931-6942. (1) Cahiez, G.; Habiak, V.; Gager, O. Org. Lett. 2008, 10, 2389–2392. (m) Naodovic, M.; Xia, G.; Yamamoto, H. Org. Lett. 2008, 10, 4053-4055 (n) Sen, S.; Singh, S.; Sieburth, S. McN. J. Org. Chem. 2009, 74, 2884-2886. (o) Kim, S.; Seomoon, D.; Lee, P. H. Chem. Commun. 2009, 1873-1875. (p) Zheng, C.; Wang, D.; Stahl, S. S. J. Am. Chem. Soc. 2012, 134, 16496-16499. (q) Ting, C.-M.; Hsu, Y.-L.; Liu, R.-S. Chem. Commun. 2012, 48, 6577-6579. (r) Campbell, N. E; Sammis, G. N. Angew. Chem., Int. Ed. 2014, 53, 6228–6231. (s) Wang, Z.; Wang, Y.; Zhang, L. J. Am. Chem. Soc. 2014, 136, 8887-8890.

(28) (a) Albéniz, A. C.; Espinet, P.; López-Fernández, R.; Sen, A. J. Am. Chem. Soc. 2002, 124, 11278-11279. (b) Guan, H. ; limura, M.; Magee, M. P.; Norton, J. R.; Janak, K. E. Organometallics 2003, 22, 4084-4089. (c) Tan, E. H. P.; Lloyd-Jones, G. C.; Harvey, J. N.; Lennox, A. J. J.; Mills, B. M. Angew. Chem., Int. Ed. 2011, 50, 9602-9606. (d) Li, G.; Han, A.; Pulling, M. E.; Estes, D. P.; Norton, J. R. J. Am. Chem. Soc. 2012, 134, 14662-14665. (e) Ventre, S.; Derat, E.; Amatore, M.; Aubert, C.; Petit, M. Adv. Synth. Catal. 2013, 355, 2584-2590. (f) Hu, Y.; Norton, J. R. J. Am. Chem. Soc. 2014, 136, 5938-5948.

SYNOPSIS TOC



1

2

3

4