Acta Crystallographica Section C Crystal Structure Communications

ISSN 0108-2701

Two germatranes with bulky substituents

Alexander A. Korlyukov,^a* Eugene A. Komissarov,^a Nikolay V. Alekseev,^b Konstantin V. Pavlov,^b Olga V. Krivolapova^b and Valentin G. Lachtin^b

^aA. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Science, 28 Vavilov Street, Moscow 119991, Russian Federation, and ^bState Research Institute of Chemistry and Technology of Organoelement Compounds, 38 Entuziastov Sh., Moscow 111123, Russian Federation Correspondence e-mail: alex@xrlab.ineos.ac.ru

Received 28 April 2006 Accepted 23 May 2006 Online 23 June 2006

In 1-adamantyl-2,8,9-trioxa-5-aza-1-germabicyclo[3.3.3]undecane or 1-adamantylgermatrane, $[Ge(C_{10}H_{15})(C_6H_{12}NO_3)]$, (I), and (2,8,9-trioxa-5-aza-1-germabicyclo[3.3.3]undecan-1-yl)methyl *N*-cyclohexylcarbamate or [(germatran-1-yl)methyl] -cyclohexylcarbamate, $[Ge(C_6H_{12}NO_3)(C_8H_{14}NO_2)]$, (II), the Ge atoms are characterized by trigonal–bypiramidal configurations. The Ge···N distances [2.266 (3) and 2.206 (3) Å in (I) and (II), respectively] are among the longest observed in germatranes. The significant distortion of the apical N–Ge–C angle in (II) is caused by crystal packing effects.

Comment

Organogermanic compounds with expanded coordination spheres around the metal, such as substituted germatranes, have been of interest since 1970. Most probably, this is due to their potential use as biologically active substances (Lukevics *et al.*, 1990, 1992). The wide spectrum of pharmacological action of organogermanic compounds, their low toxicity and other useful properties have created favourable conditions



for the development of bioorganogermanic chemistry. Successful progress in this area is impossible without a wide search for the most active substances and discovering the correlation between their structure and biological activity. As a continuation and development of our work in this area (Gurkova *et al.*, 1984; Knyazev *et al.*, 2000; Korlyukov *et al.*, 2003), the crystal and molecular structures of 1-germanyla-damantane, (I), and [1-(germathranyl)methyl]cyclohexyl-

carbamate, (II), have been studied by X-ray diffraction and the results are presented here (Figs. 1 and 2).

The coordination around atom Ge1 in (I) and (II) is trigonal-bipyramidal. Atom N1 of the atrane framework and the organic R group occupy the axial positions, while three O atoms occupy the equatorial positions. The Ge1 atoms deviate from the plane of the equatorial O atoms towards R by 0.28 Å in (I) and 0.24 Å in (II). Analysis of the Cambridge Structural Database (CSD, Version 5.27; Allen, 2002) reveals that the interatomic Ge1...N1 distance in (I) is among the longest observed for this class of compounds. According to the CSD, the Ge1...N1 distance varies in the range 2.01–2.32 Å (for 85 examined structures). The longest Ge1...N1 distances are observed in 1-(1-trimethylsylilcyclopropyl)germatrane (Korlyukov et al., 2003) and 1-(germatranylmethyl)germatrane (Gurkova et al., 1982) (2.300 and 2.315 Å). The presence of the electron-withdrawing carbamate group in (II) leads to a decrease in this interatomic distance by 0.06 Å compared with the corresponding value in (I). The absence of a greater shortening of the Ge1 \cdots N1 interatomic distance in (II) can be explained by a significant delocalization of electron density in the N2/C8/O4/O5 carbamate fragment. Indeed, the N2-C8 and C8-O4 bonds are shorter than the respective standard single bonds, but they are noticeably longer than the corresponding double bonds (Allen et al., 1987). Thus, in both (I)



Figure 1

The molecular structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms have been omitted for clarity.



Figure 2

The molecular structure of (II), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms have been omitted for clarity.

and (II), the interatomic Ge1···N1 distances are typical for germatranes with donor substituents (for instance, in 1-ethylgermatrane, the Ge1···N1 distance is 2.24 Å; Atovmyan *et al.*, 1970).

In the present compounds, the Ge atoms are bonded to bulky carbamate and adamantane substituents. In (II), the N1-Ge1-C7 angle is 173.73 (13)°, while in the majority of germatranes, the relative values are in the range 175-179°. In the case of the adamantane group in (I), the N1-Ge1-C7 angle remains almost undistorted [178.69 (11)°]. It is of interest to study the reason for the decrease in the N1-Ge1-C7 angle in (II).

We have found that the distortion of the N1-Ge1-C7 angle is related to the formation of intra- and intermolecular contacts between the O atoms of the atrane framework and the substituents (Allen et al., 1987). For a more reliable analysis of the intramolecular contacts, we normalize all C-H bonds to 1.08 Å according to single-crystal neutron diffraction data. The germatranes with $R_1C = CR_2R_3$ substituents ($R_1 = H$, I or Br, $R_2 = H$, Br or Cl and $R_3 = Ph$; N1-Ge1-C7 angles in the range 173.5–178.1°; Karlov et al., 2001; Faller & Kultyshev, 2003; Selina, Karlov et al., 2004; Selina, Zhachkina et al., 2004) may serve as examples of the influence of intramolecular contacts on the linearity of the N1-Ge1-C7 angle. In these compounds, short $C-H \cdots O$ contacts between an O atom and a phenyl group are always observed. In addition, the Br and I atoms also form short intermolecular contacts with O atoms of the atrane framework (torsion angle $O-Si-C-Hal \simeq 22^\circ$, where Hal = Br or I).

The influence of the intermolecular contacts can be demonstrated by methyl (germatranyl)trimethylsilylacetate (Zaitseva *et al.*, 1997) and (pentafluorophenyl)germatrane (Kultyshev *et al.*, 2004) (N1-Ge1-C7 = 175.73 and 176.29°, respectively). In the case of the former compound, the acetate group forms four C-H···O contacts (C···O $\simeq 3.5$ Å, H··· O $\simeq 2.6$ Å and C-H···O = 107-146°) as well as three H···H and C···H contacts (H···H $\simeq 2.2$ Å and C···H $\simeq 3.1$ Å). In (pentafluorophenyl)germatrane, four C-H···F contacts



Figure 3

The crystal packing of (II). H atoms have been omitted for clarity, with the exception of atom H2C. The N2-H2C \cdots O1 hydrogen bonds are shown as dashed lines.

 $(C \cdots F \simeq 3.4 \text{ Å}, H \cdots F \simeq 2.5 \text{ Å} \text{ and } C - H \cdots F = 138 - 164^{\circ})$ and one C-H··· π contact (H···Cg $\simeq 2.8$ Å and C-H··· $\pi = 116^{\circ}$, where Cg is the centroid of the C₆F₅ ring) are observed. Of course, all the above contacts are weak, but their total energy is sufficient for the distortion of the N1-Ge1-C7 angle. In (II), one may see that the N1-Ge1-C7 angle deviates further from the ideal value of 180° than in other germatranes. Such a deviation is related to the influence of crystal packing. Actually, atom N2 of the carbamate group participates in an N2-H2···O1 hydrogen bond [H2C···O1 = 2.077 Å, N2··· O1 = 2.951 (4) Å and $N2 - H2C - O1 = 172^{\circ}$; symmetry code: $x, \frac{3}{2} - y, z - \frac{1}{2}$, while atom O5 forms a contact with the atrane framework (Fig. 3). Taking into account the noticeable flexibility of the Ge coordination polyhedron, one may conclude that the formation of the N-H···O interaction leads to distortion of the N1–Ge1–C7 angle.

In (I), the presence of the adamantane fragment does not lead to a significant deviation of N1–Ge1–C7 from ideal linearity. The adamantane hydrocarbon does not afford the O atoms of the atrane cage the opportunity to form intermolecular contacts. In turn, the intramolecular contacts of the methylene groups of the adamantane fragment to the O atoms of the atrane cage are unfavourable, due to large O···H distances (2.7–2.9 Å) and small C–H···O angles (106–110°). So one may therefore conclude that the bulky adamantane group preserves the N1–Ge1–C7 angle from distortion due to crystal packing.

Experimental

The synthesis of (I) was carried out according to the previously published method of Gar *et al.* (1985). The compound was recrystallized from hexane. Compound (II) was prepared from (trichlorogermyl)methanol (21.0 g, 0.1 mol), isocyanocyclohexane (12.5 g, 0.1 mol) and triethanolamine (14.0 g, 0.9 mol). The reaction mixture was heated until gas evolution ceased. After cooling to room temperature, crystals of (II) appeared (yield 17.6 g, 46.9%; m.p. 421–425 K). ¹H NMR (Bruker AM-360 spectrometer; 360 MHz, DMSO- d_6): δ 1.53–1.92 (*m*, 11H, C₆H₁₁), 1.96 (*t*, 6H, CH₂), 3.50 (*t*, 6H, CH₂O), 4.42 (*s*, 1H, NH), 4.64 (*s*, 2H, OCH₂Ge); IR (Bruker IFS-113V spectrometer; KBr tablet, *v*, cm⁻¹): 3283 (NH), 1707 (C=O), 1103, 1074, 1045, 1022 [Ge(OCH₂CH₂)₃N].

Compound (I)

Crystal data

 $\begin{bmatrix} Ge(C_{10}H_{15})(C_{6}H_{12}NO_{3}) \end{bmatrix} \\ M_{r} = 353.98 \\ Monoclinic, P2_{1}/c \\ a = 10.641 (6) Å \\ b = 7.292 (4) Å \\ c = 19.943 (11) Å \\ \beta = 90.001 (12)^{\circ} \\ V = 1547.3 (14) Å^{3} \end{bmatrix}$

Data collection

- Bruker SMART 1000 CCD areadetector diffractometer
- φ and ω scans
- Absorption correction: multi-scan (SADABS; Sheldrick, 1998) $T_{min} = 0.587, T_{max} = 0.826$

Z = 4 D_x = 1.520 Mg m⁻³ Mo K α radiation μ = 1.99 mm⁻¹ T = 120 (2) K Prism, colourless 0.30 × 0.10 × 0.10 mm

15300 measured reflections 4396 independent reflections 3193 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.059$ $\theta_{\text{max}} = 30.0^{\circ}$ Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.038$ $wR(F^2) = 0.108$ S = 0.974396 reflections 191 parameters

Compound (II)

Crystal data

[Ge(C₆H₁₂NO₃)(C₈H₁₄NO₂)] $M_r=374.96$ Monoclinic, $P2_1/c$ a = 9.4398 (13) Å b = 16.690 (2) Å c = 10.3361 (16) Å $\beta = 97.740 \ (3)^{\circ}$ V = 1613.6 (4) Å³

Data collection

Bruker SMART CCD 1000 areadetector diffractometer φ and ω scans Absorption correction: multi-scan (SADABS; Sheldrick, 1998) $T_{\rm min}=0.700,\ T_{\rm max}=0.945$

Refinement

Refinement on F^2	H-atom parameters constrained		
$R[F^2 > 2\sigma(F^2)] = 0.047$	$w = 1/[\sigma^2(F_o^2) + (0.0462P)^2]$		
$wR(F^2) = 0.108$	where $P = (F_0^2 + 2F_c^2)/3$		
S = 1.05	$(\Delta/\sigma)_{\rm max} = 0.001$		
3487 reflections	$\Delta \rho_{\rm max} = 1.53 \text{ e} \text{ Å}^{-3}$		
199 parameters	$\Delta \rho_{\rm min} = -0.39 \text{ e} \text{ Å}^{-3}$		

Table 1

Selected bond lengths (Å) in (I) and (II).

Bond	(I)	(II)
Ge1···N1	2.206 (3)	2.266 (3)
Ge1-O1	1.817 (2)	1.822 (3)
Ge1-O2	1.794 (2)	1.825 (3)
Ge1-O3	1.791 (2)	1.820 (3)
Ge1-C7	1.957 (4)	1.966 (3)
N-C (mean)	1.471 (4)	1.473 (5)
O-C (mean)	1.415 (4)	1.416 (3)

Table 2

Selected bond angles (°) in (I) and (II).

Angle	(I)	(II)
N1-Ge1-C7	104.5 (2)	104.4 (3)
O-Ge-O (mean)	118.26 (12)	117.65 (14)
C-N-C (mean)	114.0 (3)	114.0 (3)
C7-O4-C8	114.3 (3)	

The H atoms were calculated geometrically and refined using a rigid-body model, with C-H = 0.99–1.00 Å and $U_{iso}(H) = 1.2U_{eq}(C)$. The crystals of (I) were found to be twinned. The structure was refined in XL using the method of Pratt et al. (1971) and Jameson

H-atom parameters constrained $w = 1/[\sigma^2(F_o^2) + (0.0517P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\rm max} < 0.001$ $\Delta \rho_{\rm max} = 1.02 \text{ e} \text{ } \text{\AA}^{-3}$ $\Delta \rho_{\rm min} = -0.56$ e Å⁻³

Z = 4 $D_x = 1.543 \text{ Mg m}^{-3}$ Mo Ka radiation $\mu = 1.92 \text{ mm}^{-1}$ T = 120 (2) K Plate, colourless $0.20 \times 0.10 \times 0.03~\text{mm}$

8957 measured reflections 3487 independent reflections 2310 reflections with $I > 2\sigma(I)$ $R_{\rm int} = 0.048$ $\theta_{\rm max} = 27.1^{\circ}$

-atom parameters constrained
=
$$1/[\sigma^2(F_o^2) + (0.0462P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $\sqrt{\sigma} = -0.001$

metal-organic compounds

(1982) with a TWIN matrix defined as TWIN $1000\overline{10}00\overline{1}$, which is the default for a monoclinic type with β approximately 90°. All calculations were carried out using SHELXTL (Version 5.10; Sheldrick, 1998).

For both compounds, data collection: SMART (Bruker, 1998); cell refinement: SAINT-Plus (Bruker, 1998); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXTL (Sheldrick, 1998); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This work was supported by the Russian Foundation for Basic Research (grant No. 04-03-32662) and the Programme for State Support of Young Scientists (grant No. MK-3578.2005.3).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SQ3019). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. (2002). Acta Cryst. B58, 380-388.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). J. Chem. Soc. Perkin Trans. 2, pp. S1-19.
- Atovmyan, L. O., Bleidelis, Ya. Ya., Kemme, A. A. & Shibaeva, R. P. (1970). Zh. Strukt. Khim. (Russ. J. Struct. Chem.), 11, 318-324. (In Russian.)
- Bruker (1998). SAINT-Plus (Version 6.01) and SMART (Version 5.059). Bruker AXS Inc., Madison, Wisconsin, USA.
- Faller, J. W. & Kultyshev, R. G. (2003). Organometallics, 22, 199-202.
- Gar, T. K., Chernyshev, O. N., Bochkarev, V. N., Chernyshev, E. A., Tandura, S. N., Kisin, A. V. & Mironov, V. F. (1985). Zh. Obshch. Khim. 55, 1057–1063.
- Gurkova, S. N., Gusev, A. I., Sharapov, V. A., Alekseev, N. V., Gar, T. K. & Chromova, N. J. (1984). J. Organomet. Chem. 268, 119-124.
- Gurkova, S. N., Tandura, S. N., Kisin, A. V., Gusev, A. I., Alekseev, N. V., Gar, T. K., Khromova, N. Yu. & Segel'man, I. R. (1982). Zh. Strukt. Khim. (Russ. J. Struct. Chem.), 23, 104-109.
- Jameson, G. B. (1982). Acta Cryst. A38, 817-820.
- Karlov, S. S., Shutov, P. L., Churakov, A. V., Lorberth, J. & Zaitseva, G. S. (2001). J. Organomet. Chem. 627, 1-5.
- Knyazev, S. P., Kirin, V. N., Alekseev, N. V., Korlyukov, A. A., Varnavskaya, I. A., Chernyshev, E. A., Lysenko, K. A. & Antipin, M. Yu. (2000). Dokl. Akad. Nauk, 371, 50-54.
- Korlyukov, A. A., Lyssenko, K. A., Antipin, M. Yu., Alekseev, N. V., Knyazev, S. P. & Chernyshev, E. A. (2003). J. Mol. Struct. 655, 215-220.
- Kultyshev, R. G., Prakash, G. K. S., Olah, G. A., Faller, J. W. & Parr, J. (2004). Organometallics, 23, 3184-3188.
- Lukevics, E. Ya., Gar, T. K., Ignatovich, L. V. & Mironov, V. F. (1990). Biologicheskaja aktivnost' soedinenii germaniya (Biological Activity of Germanium Compounds). Riga: Zinatne. (In Russian.)
- Lukevics, E. Ya., Germane, S. & Ignatovich, L. V. (1992). Appl. Organomet. Chem. 6, 543-564.
- Pratt, C. S., Coyle, B. A. & Ibers, J. A. (1971). J. Chem. Soc. pp. 2146-2151.
- Selina, A. A., Karlov, S. S., Gauchenova, E. V., Churakov, A. V., Kuz'mina, L. G., Howard, J. A. K., Lorberth, J. & Zaitseva, G. S. (2004). Heteroatom Chem. 15, 43-56.
- Selina, A. A., Zhachkina, A. E., Karlov, S. S., Churakov, A. V. & Zaitseva, G. S. (2004). Heteroatom Chem. 15, 169-174.
- Sheldrick, G. M. (1998). SADABS (Version 2.01) and SHELXTL (Version 5.10). Bruker AXS Inc., Madison, Wisconsin, USA.
- Zaitseva, G. S., Livantsova, L. I., Nasim, M., Karlov, S. S., Churakov, A. V., Howard, J. A. K., Avtomonov, E. V. & Lorberth, J. (1997). Chem. Ber. 130, 739-796