

A Highly Selective Fluorescence Turn-on Probe for Zn²⁺ Based on New Diaryloxadiazole Chelate

Sabir H. Mashraqui,^{*1} Mukesh A. Chandiramani,¹ Sushil S. Ghorpade,¹ Carolina Estarellas,² and Antonio Frontera²

¹Department of Chemistry, University of Mumbai, Vidyanaigari, Santacruz (E), Mumbai-400098, India

²Department of Química, Universitat de les Illes Balears, Crta. De Valldemossa km 7.5, 07122 Palma (Balears), Spain

(Received July 8, 2011; CL-110585; E-mail: sh_mashraqui@mu.ac.in)


A new chelating diaryloxadiazole probe, oxazid displayed emission red shift and 16-fold emission enhancement upon binding with Zn²⁺ in buffered condition, while competing Cd²⁺, Ca²⁺, Mg²⁺, and several potentially quenching metal ions posed no significant optical interferences. Experimental evidences indicated 1:1 complexation, and the computational work revealed formation of a stable octahedral geometry.

Metal-ion-triggered optical modulation of photoemittive receptors constitutes one of the key approaches for the recognition of chemically and biologically significant metal ions.¹ Zinc, an essential micronutrient, regulates many cellular and enzymatic processes.² However, Zn²⁺ metabolic disorder is associated with many debilitating neurological conditions.³ Consequently, a heightened interest persists in designing Zn²⁺-selective optical probes.⁴ Among the known Zn²⁺ luminophores, some prominent examples include TSQ, RhodaZin-3, ZinPyr-1, Rhodafluor-2, and ZnAF-1.⁵ These probes are constructed by hosting nitrogenous zinc chelators, such as dipicolylamine, iminodiacid or cyclen onto the photoemittive fluorescein, rhodamine, or quinoline platforms.⁵ Despite continuing advances, the cross affinities, especially from biologically abundant Ca²⁺ and Mg²⁺ and toxic Cd²⁺, frequent emission quenching observed from Co²⁺, Ni²⁺, and Cu²⁺, and the need for nonaqueous measurements forms limitations in many cases. Thus, it was of interest to explore new class of fluorescent chelators which might exhibit selective discrimination of Zn²⁺ over competing metal ions, thereby enabling useful sensing properties.

Diaryloxadiazole motif is a stable fluorophore, which others⁶ and we⁷ have previously incorporated as a reporter group to design certain luminescent metal-ion sensors. With intent to extending the range of the reporter ligands for Zn²⁺, we now describe a new diaryloxadiazole chemoreceptor, oxazid, carrying oxyacetic acid chelates for the selective “off-on” signaling for Zn²⁺ under semiaqueous buffer conditions.

Synthesis of oxazid involved O-alkylation of 2,5-di(2-hydroxyphenyl)-1,3,4-oxadiazole (**1**) with *t*-butyl bromoacetate in K₂CO₃/DMF to form diester **2**, which upon acid-catalyzed hydrolysis afforded the target, oxazid⁸ (Scheme 1). With regard to the sensing mechanism, we presumed that metal-ion binding might induce the chelation-enhanced fluorescence due to the conversion of the conformationally flexible probe into the conformationally restricted metal ion complex.⁹

The fluorescence sensitivities of oxazid toward selected metal ions of biological and/or environmental relevance were evaluated in MeCN–H₂O (7:3 v/v, buffered by 10 mM Tris–HCl, pH 7.2)¹⁰ Excitation of oxazid at 314 nm resulted in a weak emission band at 352 nm with quantum yield Φ_f of 0.017



Scheme 1. Synthetic sequence toward diaryloxadiazole receptor, oxazid.




Figure 1. Fluorimetric titration of oxazid (1×10^{-6} M) with incremental Zn²⁺ (0 – 10^{-4} M) in MeCN–H₂O (7:3 v/v, buffered by 10 mM Tris–HCl).

measured with respect to anthracene. The low Φ_f may be attributable in part to the nonradiative aryl-ring-rotation processes in free probe. The emission profile of oxazid (1×10^{-6} M) was not significantly perturbed when 500 equiv of Li⁺, Na⁺, K⁺, Mg²⁺, Ca²⁺, Ba²⁺, Co²⁺, Ni²⁺, Cu²⁺, Cd²⁺, Hg²⁺, and Pb²⁺ as their perchlorates were introduced.⁹ In contrast, addition of just 100 equiv of Zn²⁺ induced marked changes both in the intensity and energy of the probe's emission.

Fluorimetric titration of oxazid (Figure 1) revealed linear enhancement in emission intensity with incremental addition of Zn²⁺. At saturating 10^{-4} M of Zn²⁺, a high 16-fold emission enhancement ($\Phi_f = 0.264$), accompanied with 17 nm red shift was observed. Figure 2a shows that relative to other metal ions examined, only Zn²⁺ is capable of inducing high fluorescence “off-on” signaling response. To further demonstrate the selective interaction with Zn²⁺, initially the fluorescence of the probe (1×10^{-6} M) in the presence of a matrix consisting of 5×10^{-4} M each of Li⁺, Na⁺, K⁺, Mg²⁺, Ca²⁺, Ba²⁺, Co²⁺, Ni²⁺, Cu²⁺, Cd²⁺, Hg²⁺, and Pb²⁺ was recorded. As shown in Figure 2b, the matrix barely impacted the fluorescence intensity (<1.5-fold enhancement) of the probe. Subsequently, addition of just 10^{-4} M of Zn²⁺ induced emission enhancement by ca. 14-fold. This finding clearly validates the selective binding of Zn²⁺ even though other metals may be present in excess




Figure 2. a) Relative emission intensity of oxazid (1×10^{-6} M) with metal ions (100 equiv), b) competitive matrix experiment: emission spectra of oxazid alone (1 μ g), oxazid + 5×10^{-4} M each of Li^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Ba^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , Cd^{2+} , Pb^{2+} , and Hg^{2+} and oxazid + matrix + Zn^{2+} (10^{-4} M).

concentrations. The Job plot indicated 1:1 binding stoichiometry, and the stability constant, $\log K$ for Zn^{2+} , was determined to be 3.72.⁸ For other metal ions tested, the emission changes were relatively insignificant to allow a reliable measure of their $\log K_S$. The detection limit of Zn^{2+} determined from the fluorescence data was found to be 5.75×10^{-7} M.⁸

The shift of the IR band from 1694 cm^{-1} in the free receptor to 1600 cm^{-1} in the oxazid–zinc complex confirms the coordination of carboxylic acids with Zn^{2+} .⁸ The ^1H NMR spectrum of the probe displayed a singlet for $-\text{OCH}_2\text{CO}_2-$ at δ 4.76, but in the Zn^{2+} complex, this signal, while retaining its singlet character, was upfield shifted to δ 4.41. In addition, relative to the free probe, the aryl ring protons in the zinc complex were also upfield shifted in the range of δ 0.13 to 0.22.⁸ These NMR data together with Job's plot imply that both the $-\text{OCH}_2\text{CO}_2\text{H}$ groups are probably bound to a single, centrally located Zn^{2+} in a 1:1 complexation mode.

To theoretically study the energetic and geometric features of the oxazid– Zn^{2+} complex, calculations were performed using DFT (RI-BP86/def2-TZVP) within the program TURBOMOLE^{11a} version 5.10 taking into account solvent effects (water) using the COSMO approximation.^{11b} Depending on the orientations of the oxadiazole ring, several possible geometries of 1:1 complexation were analyzed. However, we found only one computationally suitable conformation which had the oxygen atom of the oxadiazole ring pointing to the Zn^{2+} (**1•ZnA**, Figure 3). Inclusion of an explicit water molecule in the inner coordination sphere of the Zn^{2+} (**1•ZnB**) afforded extra $5.4 \text{ kcal mol}^{-1}$ stabilization.¹² The ability of the probe to form stable octahedral complex could account for the selective zinc coordination.

Understandably, Cu^{2+} , Co^{2+} , and Ni^{2+} , which predominantly form either the square-planar or tetrahedral coordination, fail to interact. Moreover, in the Zn^{2+} octahedral complex, the rotations around aryl–oxadiazole bonds would be severely restricted. This phenomena might contribute to emission amplification by enhancing the radiative channels over thermal quenching processes prevailing in the conformationally flexible probe.⁸

In conclusion, we have reported a new, structurally simple diaryloxadiazole chelate which affords high fluorescence turn-on response for targeting Zn^{2+} . Noteworthy, many potentially interfering or quenching metal ions displayed no significant optical perturbations even in relatively higher concentrations. The present probe, carrying wholly “oxygen”-based ligand




Figure 3. Optimized structures of oxazid– Zn^{2+} complex and their difference in energies.

system should be a welcome new addition to the repertoire of the potentially useful zinc fluorescence sensors.

References and Notes

- a) A. P. de Silva, H. Q. N. Gunaratne, T. Gunnlaugsson, A. J. M. Huxley, C. P. McCoy, J. T. Rademacher, T. E. Rice, *Chem. Rev.* **1997**, *97*, 1515. b) *Fluorescent Chemosensors for Ion and Molecule Recognition in ACS Symposium Series 538*, ed. by A. W. Czarnik, American Chemical Society, Washington, D.C., **1993**. doi:10.1021/bk-1993-0538.
- a) J. M. Berg, Y. Shi, *Science* **1996**, *271*, 1081. b) B. L. Vallee, D. S. Auld, *Biochemistry* **1990**, *29*, 5647. c) J. J. R. F. da Silva, R. J. P. Williams, *The Biological Chemistry of the Elements: The Inorganic Chemistry of Life*, Clarendon, Oxford, **1991**.
- a) C. J. Frederickson, J.-Y. Koh, A. I. Bush, *Nat. Rev. Neurosci.* **2005**, *6*, 449. b) C. J. Frederickson, E. J. Kasarskis, D. Ringo, R. E. Frederickson, *J. Neurosci. Methods* **1987**, *20*, 91.
- a) Z. Xu, J. Yoon, D. R. Spring, *Chem. Soc. Rev.* **2010**, *39*, 1996. b) P. Jiang, Z. Guo, *Coord. Chem. Rev.* **2004**, *248*, 205. c) K. Kikuchi, K. Komatsu, T. Nagano, *Curr. Opin. Chem. Biol.* **2004**, *8*, 182.
- a) K. Kiyose, H. Kojima, Y. Urano, T. Nagano, *J. Am. Chem. Soc.* **2006**, *128*, 6548. b) K. Komatsu, K. Kikuchi, H. Kojima, Y. Urano, T. Nagano, *J. Am. Chem. Soc.* **2005**, *127*, 10197. c) S. C. Burdette, C. J. Frederickson, W. Bu, S. J. Lippard, *J. Am. Chem. Soc.* **2003**, *125*, 1778. d) C. R. Goldsmith, S. J. Lippard, *Inorg. Chem.* **2006**, *45*, 555. e) K. R. Gee, Z.-L. Zhou, W.-J. Qian, R. Kennedy, *J. Am. Chem. Soc.* **2002**, *124*, 776. f) M. Taki, J. L. Wolford, T. V. O'Halloran, *J. Am. Chem. Soc.* **2004**, *126*, 712. g) T. Hirano, K. Kikuchi, Y. Urano, T. Higuchi, T. Nagano, *Angew. Chem., Int. Ed.* **2000**, *39*, 1052.
- a) G. Ambrosi, M. Formica, V. Fusi, L. Giorgi, E. Macedi, M. Micheloni, P. Paoli, R. Pontellini, P. Rossi, *Inorg. Chem.* **2010**, *49*, 9940. b) X.-L. Tang, X.-H. Peng, W. Dou, J. Mao, J.-R. Zheng, W.-W. Qin, W.-S. Liu, J. Chang, X.-J. Yao, *Org. Lett.* **2008**, *10*, 3653.
- a) S. H. Mashraqui, S. Sundaram, A. C. Bhasikuttan, *Tetrahedron* **2007**, *63*, 1680. b) S. H. Mashraqui, S. Sundaram, T. Khan, A. C. Bhasikuttan, *Tetrahedron* **2007**, *63*, 11093. c) S. H. Mashraqui, S. Sundaram, A. C. Bhasikuttan, S. Kapoor, A. V. Sapre, *Sens. Actuators, B* **2007**, *122*, 347.
- Supporting Information is available electronically on the CSJ-Journal Web site, <http://www.csj.jp/journals/chem-lett/index.html>, and includes synthesis of oxazid, IR, NMR, and Mass data, UV-vis, Job's plot, stability constant, and detection limit.
- a) Z. Zhou, C. J. Fahrni, *J. Am. Chem. Soc.* **2004**, *126*, 8862. b) J. V. Mello, N. S. Finney, *Angew. Chem., Int. Ed.* **2001**, *40*, 1536. c) K. R. A. S. Sandanayake, K. Nakashima, S. Shinkai, *J. Chem. Soc., Chem. Commun.* **1994**, 1621.
- UV-vis profile of the probe remained essentially invariant to several metal ions, including Zn^{2+} , implying none or at best weak perturbation of the ground state of the probe.
- a) R. Ahlrichs, M. Bär, M. Häser, H. Horn, C. Kölmel, *Chem. Phys. Lett.* **1989**, *162*, 165. b) A. Klamt, G. Schüürmann, *J. Chem. Soc., Perkin Trans. 2* **1993**, 799.
- The interaction energy was computed as the difference between the optimized complex (with water) and the sum of the monomers (water and the complex without water).