

Efficient oxidative radical spirolactamization†

Tannya R. Ibarra-Rivera,^a Rocío Gámez-Montaña^{*a} and Luis D. Miranda^{*b}

Received (in Cambridge, UK) 11th April 2007, Accepted 22nd May 2007

First published as an Advance Article on the web 14th June 2007

DOI: 10.1039/b705397e

An efficient xanthate-based method for the preparation of azaspirocyclic cyclohexadienones *via* an *ipso* oxidative radical cyclization of *p*-oxygenated *N*-benzylacetamides and *N*-phenethylacetamide is described.

Azaspirocyclic cyclohexadienones are pivotal synthetic intermediates in the preparation of an extensive range of biologically active molecules.¹ Recently the azaspirocyclic core was shown to be present in annosqualine **1**, a natural product isolated from the stems of *Annona Squamosa* (Fig. 1).² Furthermore, the reduced congeners 2-azaspiro[4.5]decane and 2-azaspiro[5.5]undecane have a variety of biological activities.³ For instance, the azaspirane **2** has inhibitory action in KB cells and cells of human mammary cancer grown in tissue culture.⁴ The spiroperidine BL1743⁵ is an antiviral and L687384⁶ is a potent and selective σ -receptor ligand (Fig. 1).

Several strategies have been reported for the construction of azaspirocyclic cyclohexadienone systems. The formation of azaspirocyclics from *p*-MeO-benzene derivatives was observed in the acid-catalyzed cyclization of aromatic diazoacetamides⁷ and also in the intramolecular addition of stabilized enolates to (η^6 -arene)ruthenium complexes.⁸ Radical spirocyclization onto a *p*-MeO-aryl ring was first observed by Hey and Todd⁹ in 1967, but the process was of limited synthetic value. Zard and co-workers have observed the formation of one azaspirocyclic cyclohexadienone in moderate yield, by a nickel/acetic acid induced oxidative *ipso*-type radical cyclization onto *p*-MeO-benzenoid systems.¹⁰ Very recently, Gonzalez-Lopez de Turiso and Curran¹¹ reported a similar (TMS)₃SiH-mediated cyclization, but only low yields of the spirocyclic systems were obtained. Herein, we describe an efficient

method for the preparation of highly substituted azaspirocyclic cyclohexadienones *via* an *ipso* oxidative radical cyclization of *p*-oxygenated *N*-benzylacetamides and homologs thereof (*e.g.*, Scheme 1).¹² Our studies commenced with the readily available xanthate **5a**,¹³ which upon reaction with a slight excess (1.2 equiv.) of dilauroyl peroxide in 1,2-dichloroethane at reflux temperature, was converted exclusively into the spirocyclic derivative **6**.¹⁴ Under the same conditions, the *p*-hydroxylated compound **5b** also gave **6** in nearly quantitative yield.

To investigate the scope of the spirolactamization process, a series of *p*-alkoxylated *N*-benzyl and *N*-phenethylacetamido derivatives (Table 1) was prepared and subjected to the above reaction conditions. The expected spirolactams were obtained as the exclusive, or major products in all cases. It is noteworthy that the presence of one or more substituents (MeO, Me, Br) *ortho* or *meta* to the *p*-alkoxyl moiety does not derail the spirocyclization process (entries 1–5) and a *p*-isopropoxy group also leads to the expected product (entry 7). Furthermore, 3-azaspiro[5.5]undecane systems (*e.g.*, compounds **20a** and **20b**; entry 7) are readily generated starting from *N*-*p*-alkoxyphenethylacetamides. Secondary xanthates can also be used, in which case azaspirocyclohexadienones containing a methyl substituent α to the lactam carbonyl group are produced (entry 6). The remaining *N*-substituent in the starting materials does not have to be a *tert*-butyl group, since the *N*-isopropyl (entry 7) and the *N*-dimethoxyphenethyl (entry 11) compounds also gave rise to spiro products. The xanthate **21** is a particularly interesting substrate since the formation of both 2-azaspiro[4.5]decane and 3-azaspiro[5.5]undecane systems is possible (entry 8). Nevertheless, the only spiro product isolated (65% yield), containing the [4.5]decane system, was compound **22**. Finally, the formation of the spiro compound **24**, derived from the naphthalene precursor **23** indicates that this process does indeed have considerable scope (entry 9).

A possible mechanistic rationalization for the formation of the spiro compounds is depicted in Scheme 2. The cyclohexadienyl radical **26** produced from the *ipso*-cyclization, is then oxidized to the oxonium ion **27**, or the corresponding protonated species, by lauroyl peroxide. The oxonium ion would then be transformed into the observed product in the reaction medium or upon workup of the reaction mixture. The success of the reaction is dependent on





Fig. 1 Azaspirocyclics.

^aFacultad de Química, Universidad de Guanajuato, Noria Alta S/N, Guanajuato, México C.P. 36050

^bInstituto de Química, Universidad Nacional Autónoma de México, Circuito Exterior S. N., Ciudad Universitaria, Coyoacán, México D. F. 04510, México. E-mail: lmiranda@sevidor.unam.mx; Fax: (+52) 55 56 16 22 17; Tel: (+52) 55 56 22 44 40

† Electronic supplementary information (ESI) available: Experimental details and spectroscopic data for all new compounds. See DOI: 10.1039/b705397e




Scheme 1

Table 1

Entry	Xanthate	Azaspirocycle
1	7a R = OMe 7b, R = Me 7c R = Br	8a R = OMe 75% 8b, R = Me 83% 8c R = Br 75%
2	9	10, 85%
3	11	12, 70%
4	13	14, 65%
5	15	16, 74%
6	17	18, 68%
7	19a R = H 19b, R = OMe,	20a R = H 65% 20b, R = OMe 57%
8	21	22, 65%
9	23	24, 65%

^a Xtht = SC(S)OEt.



Scheme 2

the use of slightly more than one equivalent of the peroxide, and this stoichiometry is consistent with the proposed oxidation mechanism.

The efficacy of the *ipso* cyclization may be a consequence of both a polarity match between the attacking electrophilic radical and the nucleophilic *ipso* benzenoid carbon (*para* to the methoxyl group) as well as to the formation of the highly resonance stabilized radical **26**.

In short, we have developed a novel, simple, and efficient procedure for the synthesis of highly functionalized 2-azaspiro-[4.5]decan-3-ones and 3-azaspiro[5.5]undecan-4-ones. We are currently studying the synthetic consequences of, and the mechanistic questions which, have been raised by the results described herein.

We thank CONACYT (J42673Q) for financial support and Dr Joseph M. Muchowski for many helpful discussions. Also we thank R. Patiño, J. Pérez, L. Velasco, H. Rios, N. Zavala, E. Huerta and A. Peña, for technical support.

Notes and references

- (a) D. J. Wardrop, M. S. Burge, W. Zhang and J. A. Ortiz, *Tetrahedron Lett.*, 2003, **44**, 2587; (b) G. M. Rishton and M. A. Schwartz, *Tetrahedron Lett.*, 1988, **29**, 2643.
- Y. L. Yang, F.-R. Chang and Y.-C. Wu, *Helv. Chim. Acta*, 2004, **87**, 1392.
- (a) W. M. Kazmierski, E. Furfine, A. Spaltenstein and L. L. Wright, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 3431; (b) C. R. Albrightson, P. J. Bugelski, T. A. Berkhout, B. Jackson, W. D. Kerns, A. J. Organ and A. M. Badger, *J. Pharmacol. Exp. Ther.*, 1995, **272**, 689.
- (a) L. M. Rice, B. S. Sheth and J. W. Wheeler, *J. Heterocycl. Chem.*, 1973, **10**, 731; (b) A. M. Badger, T. M. Newman-Tarr and J. L. Satterfield, *Immunopharmacology*, 1997, **37**, 53.
- G. S. Gandhi, K. Shuck, J. D. Lear, G. R. Diekmann, W. F. DeGrado, R. A. Lamb and L. H. Pinto, *J. Biol. Chem.*, 1999, **274**, 5474.
- C. A. Maier and B. Wünsch, *J. Med. Chem.*, 2002, **45**, 438.
- G. M. Rishton and M. A. Schwartz, *Tetrahedron Lett.*, 1988, **29**, 2643.
- (a) F. C. Pigge, S. Fang and N. Rath, *J. Am. Chem. Soc.*, 2006, **128**, 3498, and references therein.
- D. H. Hey and A. R. Todd, *J. Chem. Soc. C*, 1967, 1518.
- J. Boivin, M. Yousfi and S. Z. Zard, *Tetrahedron Lett.*, 1997, **38**, 5985.
- F. Gonzalez-Lopez de Turiso and D. P. Curran, *Org. Lett.*, 2005, **7**, 151, and references cited therein.
- For references on ipso-type radical cyclization onto other substituted aromatic systems, see: furans: (a) S. Guindeuil and S. Z. Zard, *Chem. Commun.*, 2006, 665; (b) A. Demircan and P. Parsons, *Eur. J. Org. Chem.*, 2003, 1729; pyrroles: (c) K. Jones, T. C. T. Ho and J. Wilkinson, *Tetrahedron Lett.*, 1995, **36**, 6743; (d) C. Escolano and K. Jones, *Tetrahedron Lett.*, 2000, **41**, 8951; (e) C. Escolano and K. Jones, *Tetrahedron*, 2002, **58**, 1453; benzofurans: (f) A. S. Kyei, K. Tchabanenko, J. E. Baldwin and R. M. Adlington, *Tetrahedron Lett.*, 2004, **45**, 8931; indoles: T. Hilton, T. C. T. Ho, G. Pljevaljcic and K. Jones, *Org. Lett.*, 2000, **2**, 2639; benzene: H. Ohno, S.-I. Maeda, M. Okumura, R. Wakayama and T. Tanaka, *Chem. Commun.*, 2002, 316.

13 B. Quiclet-Sire and S. Z. Zard, *Chem. Eur. J.*, 2006, **12**, 6002.

14 *Typical procedure for radical spirocyclization.* A deaerated solution of the corresponding xanthate (1.0 mmol) in 1,2-dichloroethane (5 mL) was heated at reflux and 10% mol lauryl peroxide (1.2–1.5 mmol) was then added every 1 h until complete consumption of the xanthate was observed by TLC. The solvent was removed under reduced pressure and the residue purified by chromatography on a silica gel column

(hexane–EtOAc) to furnish the desired product. *Selected spectral data for 6.* IR (neat, cm^{-1}): 1676, 1628; ^1H NMR (300 MHz, CDCl_3) δ 6.96 (d, 2H, $J = 10.2$ Hz), 6.34 (d, 2H, $J = 10.2$ Hz), 4.49 (s, 2H), 2.54 (s, 2H), 1.43 (s, 9H); ^{13}C NMR (300 MHz, CDCl_3) δ 185.0, 171.6, 150.1, 129.3, 54.6, 53.0, 42.6, 40.8, 27.6; EI-MS m/z (%) 219 (M^+ , 65); HRMS (FAB+) m/z calc. for $\text{C}_{13}\text{H}_{18}\text{NO}_2$ ($\text{M} + \text{H}^+$) 220.1338, found 220.1337.



Looking for that **special** chemical biology research paper?

TRY this free news service:

Chemical Biology

- highlights of newsworthy and significant advances in chemical biology from across RSC journals
- free online access
- updated daily
- free access to the original research paper from every online article
- also available as a free print supplement in selected RSC journals.*

*A separately issued print subscription is also available.

Registered Charity Number: 207890

RSCPublishing

www.rsc.org/chembiology

22030681