


This article was downloaded by: [University of Tennessee, Knoxville]

On: 26 December 2014, At: 16:06

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

SYNTHESIS AND X-RAY CRYSTAL STRUCTURE OF THE ASYMMETRIC DITHIADIAZENE $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$

Joanna Griffiths^a & Jeremy M. Rawson^a

^a University of Cambridge, Cambridge, United Kingdom

Published online: 12 Aug 2010.

To cite this article: Joanna Griffiths & Jeremy M. Rawson (2004) SYNTHESIS AND X-RAY CRYSTAL STRUCTURE OF THE ASYMMETRIC DITHIADIAZENE $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$, Phosphorus, Sulfur, and Silicon and the Related Elements, 179:4-5, 883-886, DOI: [10.1080/10426500490427754](https://doi.org/10.1080/10426500490427754)

To link to this article: <http://dx.doi.org/10.1080/10426500490427754>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS AND X-RAY CRYSTAL STRUCTURE OF THE ASYMMETRIC DITHIADIAZENE $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$

Joanna Griffiths and Jeremy M. Rawson
University of Cambridge, Cambridge, United Kingdom

(Received August 17, 2003; accepted October 3, 2003)

The asymmetric dithiadiazene $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$ crystallizes in the space group P-1 and exhibits a 1D polar chain structure linked via structure-directing $\text{Br}\cdots\text{O}_2\text{N}$ synthons.

Keywords: Crystal engineering; dithiadiazene; fluorination; supra-molecular synthons


The design of polar and noncentrosymmetric crystal structures for nonlinear optical (NLO) applications relies on rational crystal engineering strategies to produce materials with desirable properties. Two key features are necessary for SHG (second harmonic generation); firstly a strong molecular hyperpolarisability which is usually afforded by a polar molecule with donor and acceptor groups linked via a π -conjugated spacer group,¹ secondly, these molecules must crystallise in noncentrosymmetric space groups.² Our design strategy has focussed on the use of perfluorinated donor (D)—acceptor (A) substituted aromatics, specifically compounds of general formula $\text{DC}_6\text{F}_4\text{A}$. Results on these single ring systems³ have shown that the formation of polar chains can be encouraged by the selection of donor and acceptor groups which are well-recognized as forming linear motifs in the solid-state (supramolecular synthons). Perfluorination tends to minimize interactions between these polar chains and so favor the formation of polar acentric structures.⁴ As an extension of this work, we have begun an examination of thiazyl chains as π -delocalized spacer groups between donor-acceptor substituted perfluorinated aromatic rings. The extended delocalization is expected to enhance the molecular NLO response.

We are grateful to the EPSRC for funding (J. G.) and J. E. Davies for data collection.

Address correspondence to J. M. Rawson, Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, UK. E-mail: jmr31@cus.cam.ac.uk

RESULTS AND DISCUSSION

Synthetic routes to asymmetric dithiadiazenes are not well documented. Syntheses involving chain-scrambling of symmetrically substituted thiazyl chains have been used with some success,⁵ although these reactions have a tendency to be uncontrollable and give a mixture of products. Conversely, a single report indicated that condensation reactions between sulfenyl chlorides (RSCl) and trimethylsilyl protected sulfur diimides can be used to produce asymmetric dithiadiazenes.⁵ The methodology detailed in Scheme 1 represents a clean stepwise synthesis of the target compound, $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$ (**1**) in reasonable yield, and has been applied to prepare several other similar compounds in this laboratory.



SCHEME 1 Synthesis of **1**.

Red needle-like crystals of **1** suitable for x-ray diffraction studies were grown from a concentrated THF solution at 4°C. The asymmetric unit contains a single molecule, which is essentially planar. Within the structure, molecules are linked into polar chains by $\text{Br} \cdots \text{O}_2\text{N}$ synthons ($d_{\text{Br} \cdots \text{O}} = 3.118 \text{ \AA}$, compared to the sum of the van der Waals radii 3.08–3.38 Å). Hydrogen bonding between a nitro group in one chain and aromatic protons in an adjacent chain ($d_{\text{C-H} \cdots \text{O}} = 2.509 \text{ \AA}$, 146.8°) results in the antiparallel chain alignment seen in Figure 1.

Successive layers also align antiparallel, with short $\text{S} \cdots \text{N}$ contacts between layers ($d_{\text{S} \cdots \text{N}} = 3.411 \text{ \AA}$, compared to the sum of the van der Waals radii 3.20–3.63 Å) being reinforced by $\pi_{\text{F}} \cdots \pi_{\text{H}}$ stacking.




FIGURE 1 Single layer illustrating antiparallel chains in 1.

CONCLUSIONS

Whilst the formation of one-dimensional polar chains is encouraged through the use of the structure-directing $\text{Br}\cdots\text{O}_2\text{N}$ synthon, interactions between polar chains generate a centric structure which is therefore NLO inactive. Key intermolecular interactions promoting the centric structure are $\pi_{\text{F}}\cdots\pi_{\text{H}}$ and $\text{C}\text{--}\text{H}\cdots\text{O}$ interactions. Current attempts to synthesize perfluorinated derivatives in which these features are absent are underway.

EXPERIMENTAL

Crystal data for **1**: $\text{C}_{12}\text{H}_4\text{BrF}_4\text{N}_3\text{O}_2\text{S}_2$, $M = 442.21$, triclinic, space group $P\text{-}1$, $a = 7.5372(15)$, $b = 7.9664(16)$, $c = 13.805(3)$ $\alpha = 103.79(3)$, $\beta = 90.42(3)$, $\gamma = 114.63(3)$, $U = 726.4(3)$ \AA^3 , $T = 180(2)\text{K}$, $Z = 2$, $D = 2.022$ Mg/m^3 . Of 8474 reflections measured, 3315 were unique ($R(\text{int}) = 0.0435$) and were used in all calculations. The final $wR2 = 0.0794$ (all data), $R1 [F^2 > 2\sigma(F^2)] = 0.0326$. CCDC *****

Preparation of $\text{BrC}_6\text{F}_4\text{NSNSC}_6\text{H}_4\text{NO}_2$

SOCl_2 (1.8 ml, 2.716 g, 22.8 mmol) in toluene (20 ml) was added dropwise to a rapidly stirred solution of $p\text{-BrC}_6\text{F}_4\text{NH}_2$ (5.56 g, 22.8 mmol) in toluene (20 ml) to yield a yellow precipitate. The suspension was heated to reflux for 5 h then solvents were removed in vacuo to yield $p\text{-BrC}_6\text{F}_4\text{NSO}$ as a yellow crystalline solid which was used without further purification. A solution of $p\text{-BrC}_6\text{F}_4\text{NSO}$ (2.90 g, 10 mmol) in THF (20 ml) was added dropwise to a stirred solution of $\text{LiN}(\text{SiMe}_3)_2$ (1.67 g, 10 mmol) in THF (20 ml) at -40°C to form a deep red solution. After stirring for 1 h, the reaction mixture was allowed to warm to 0°C and

stirred for a further 1 h. After cooling again to -40°C , a solution of $p\text{-O}_2\text{NC}_6\text{H}_4\text{SCl}$ (1.85 g, 10 mmol) in THF (20 ml) was added dropwise to the stirred reaction mixture to give a red/brown precipitate. After warming to room temperature and stirring for a further 16 h, the reaction mixture was filtered to remove insoluble impurities. The solvents were removed from the filtrate in vacuo to give the **1** as a red/brown solid (1.24 g, 27.9%). Anal. Calc. for $\text{C}_{12}\text{H}_4\text{F}_4\text{O}_2\text{BrN}$: C 32.6, H 0.9, N 9.5. Found: C 32.4, H 0.9, N 9.5. EI-MS: 442.9, 440.9 (M⁺). UV-vis (CHCl_3): 457.9 nm, $\epsilon = 7.4 \times 10^4 \text{ mol dm}^{-3} \text{ cm}^{-1}$.

REFERENCES

- [1] L.-T. Cheng, W. Tam, S. H. Stevenson, G. R. Meredith, G. Rikken, and S. R. Marder, *J. Phys. Chem.*, **95**, 10631 (1991).
- [2] Ch. Bosshard, K. Sutter, Ph. Prêtre, et al., *Organic Nonlinear Optical Materials* (Gordon and Breach, Basel, 1995).
- [3] A. D. Bond, J. Griffiths, J. M. Rawson, and J. Hulliger, *J. Chem. Soc., Chem. Commun.*, 2488 (2001).
- [4] J. Hulliger, P. J. Langley, and S. W. Roth, *Cryst. Eng.*, **1**, 177 (1998).
- [5] K. Bestari, R. T. Oakley, and A. W. Cordes, *Can. J. Chem.*, **69**, 94 (1991).