This article was downloaded by: [Akdeniz Universitesi] On: 22 December 2014, At: 15:08 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

# Novel Communication: Some Novel Organosulfur Amino Acid Derivatives as Potential Antiparasitic Agents

Ragab A. EI-Sayed <sup>a</sup>

<sup>a</sup> Chemistry Department, Al-Azhar University, Nasr City, Cairo, Egypt Published online: 20 Apr 2007.

To cite this article: Ragab A. El-Sayed (2007) Novel Communication: Some Novel Organosulfur Amino Acid Derivatives as Potential Antiparasitic Agents, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:5, 1143-1151, DOI: 10.1080/10426500601142221

To link to this article: http://dx.doi.org/10.1080/10426500601142221

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



### Novel Communication: Some Novel Organosulfur Amino Acid Derivatives as Potential Antiparasitic Agents

Ragab A. El-Sayed

Chemistry Department, Al-Azhar University, Nasr City, Cairo, Egypt

The chalones 1, 14, and 26 reacted with chlorosulfonic acid to give corresponding sulfonyl chlorides 2, 15, and 27, which condensed with nucleophiles to give amino acid derivatives 3–6, 16–18, and 28–31. Some of the corresponding methyl esters 7, 8, 19–20, and 32–34 were prepared. Hydrazinolysis of these methyl esters yielded hydrazides 9, 10, 21, 22, and 35–36. Coupling reaction of some amino acid derivatives with amino acid methyl ester hydrochloride in THF-Et<sub>3</sub>N using dicyclohexylcarbodiimide method furnished the desired dipeptide methyl ester's 11–13, 23–25, and 37–38. Some spectroscopic evidence including NMR spectra analysis are briefly discussed.

**Keywords** Chalones amino acid derivatives; chlorosulfonation; sulfonyl chloride; reactions with nucleophiles as an essential amino acid

# INTRODUCTION

The work described in this article forms part of our general program concerned with the chemistry and biological evaluation of arysulfonyl derivatives.<sup>1–12</sup> Chalcone derivatives are of interest as potential biocides, since some naturally occurring antibiotics,<sup>13</sup> and aminochalcones<sup>14,15</sup> owe their activity to the presence of the  $\alpha$ ,  $\beta$ unsaturated carbonyl group. Chlorosulfonation provides a useful route to a variety of sulfonyl derivatives by nucleophilic displacement of the chlorine atom in the resulting sulfonyl chloride. These compounds are of interest for potential biocidal activity.<sup>16–19</sup>

Early workers<sup>20</sup> examined the sulfonation of chalcone and 4methoxychalcone and concluded that the latter gave sulfonic acid. Later, several chalcone-4-sulfonamides were prepared by a condensation of N-substituted P-sulfonyl benzaldiacetates with substituted acetophenones.<sup>21</sup> In chalcones (1 and 26) the 4- or 5-position should be active due to the activating influence of the olefinic double bond.

Received May 23, 2006; accepted October 17, 2006.

Address correspondence to Ragab A. El-Sayed, Chemistry Department, Faculty of Science, Al-Azhar University, Nasr City, Cairo, Egypt. E-mail:dr\_ragabaly@yahoo.com

We have demonstrated that the optimum conditions for the chlorosulfonation of chalcone (75% yield) were by a reaction with a reagent (6 mol. equivalent); for 3 weeks at r.t., the more reactive P-methoxy chalcone reacted faster under similar conditions (63% yield in one week or 75% yield in 24 h, respectively).

## DISCUSSION

The present work involves the chlorosulfonation of chalcones 1, 14, and 16 (Chart 1).

The chalcones were prepared by the standard procedure<sup>22,23</sup> involving base-catalyzed condensation of the appropriate aromatic aldehyde and accetophenone or the p-methoxy derivatives.

The different chalcones were reacted with chlorosulfonic acid, generally, 6 mol. equivalents or 3 mol. equivalents in excess thionyl chloride at r.t., to give the corresponding sulfonyl chlorides, which were condensed with a range of amino acid to give the sulfonamide listed in Table I.

Chalcones are active due to the powerful combined electron-donating methoxy group and the alkenic double bond in enhancing reactivity of the benzene ring with respect to electrophilic attack. The NMR spectra of chlacone which demonstrated that the alkenic H-proton ( $\delta$  7.79, 7.72) the most sensitive to substitutent effects.

The **NMR** spectra of 2-thienylidene sulfonylamino acid showed the resonance of thiophene protons as a doublet ( $\delta$  7.7). In chalcone PhCO.CH=CH moiety was estimated from the literature-data for PhCO and CH=CH group.<sup>24</sup>

The electron impact mass spectra of the majarity of the chalcone sulfonylamino acid derivatives showed the molecular ions  $M^+$ .

### EXPERIMENTAL

Melting points were taken on a Griffin melting point apparatus and are uncorrected. Infrared analysis of solid samples were run as a KBr disc on a Schimadzu model 440 spectrophotometer. <sup>1</sup>H NMR spectra were measured in DMSO-d<sub>6</sub> as a solvent unless otherwise stated using Fx 90 Q Fourier Transform <sup>1</sup>H NMR. Mass spectra were obtained using a Schimadzu (Japan). GC. M.S. QP 1000 Ex spectrometer using the direct inlet system. TLC analyses were carried out on Merek silica gel plates and developed with n-butanol-acetic acid-water (4:1:1) using iodine, ninhydrin, and benzidine as spraying agents.

Compounds 2, 15, and 27 were prepared according to the procedure described earlier.<sup>25,26</sup>



#### CHART 1

### Coupling Reaction 3–5, 16–18, and 28–31: General Procedure

To an amino acid (0.11 mol) in a water (25 mL) THF (15 mL) mixture was added triethylamine (5 mL), followed by a protionwise addition

4
01
6
December
Ц
2
òò
<u>.</u>
15
Ħ
~
esi
rersit
'niv
$\mathbf{c}$
niz
de
¥,
7
y
Ę
ĕ
эас
ЧC
M
õ
Ц

TABLE I Physical Data for the Chalconesulfonyl, Amino Acid Derivatives 2-13, 16-25, and 28-38

						Element	al Analysis	% Calculate	d/Found
Compound	ſ	M.P.	$\operatorname{Yield}_{\widetilde{\mathcal{X}}}$	¢	Molecular	2	:	2	2
No.	К	°C	%	Кf	Formula	% C	% Н	% N	% S
က	Gly	224 - 226	53	0.76	$\mathrm{C_{17}H_{15}NO_5S}$	59.13	4.35	4.06	9.28
						59.10	4.33	4.03	9.25
4	DL-Ala	264 - 266	64	0.73	$ m C_{18}H_{17}NO_5S$	60.17	4.74	3.90	9.91
						60.00	4.71	3.89	9.90
5	L-Val	230 - 232	70	0.80	$ m C_{20}H_{21}NO_5S$	62.02	5.43	3.62	8.27
						62.00	5.41	3.61	8.11
9	L-Leu	244 - 246	75	0.74	$ m C_{21}H_{23}NO_5S$	62.84	5.74	3.49	7.98
						62.80	5.71	3.41	7.96
7	L-Val-OMe	266-268	76	0.84	$ m C_{21}H_{23}NO_5S$	62.84	5.79	3.49	7.98
						62.84	5.73	3.45	7.96
8	L-Leu-OMe	268 - 270	79	0.77	$ m C_{22}H_{25}NO_5S$	63.61	6.02	3.37	7.71
						63.60	6.00	3.33	7.70
6	$L-Val-N_2H_3$	248 - 250	63	0.71	${ m C}_{20}{ m H}_{23}{ m N}_{3}{ m O}_{4}{ m S}$	59.85	5.74	10.47	7.98
						59.81	5.73	10.41	7.93
10	$ m L-Leu-N_2H_3$	259 - 261	66	0.77	$ m C_{21}H_{25}N_{3}O_{4}S$	60.72	6.02	10.12	7.71
						60.70	6.00	10.00	7.00
11	Gly-Gly-OMe	278 - 280	65	0.85	${ m C}_{20}{ m H}_{20}{ m N}_2{ m O}_6{ m S}$	57.69	4.81	6.73	7.69
						57.00	4.80	6.71	7.66
12	DL-Ala-Gly-OMe	255 - 257	77	0.94	${ m C}_{21}{ m H}_{22}{ m N}_2{ m O}_6{ m S}$	58.60	5.12	6.51	7.44
						58.58	5.10	6.48	7.41
13	L-Leuc-Gly-OMe	280 - 282	70	0.90	$\mathrm{C}_{24}\mathrm{H}_{28}\mathrm{N}_{2}\mathrm{O}_{6}\mathrm{S}$	61.02	5.93	5.93	6.79
						61.00	5.90	5.91	6.73
16	DL-Ala	209 - 211	55	0.61	$ m C_{19}H_{19}NO_6S$	58.61	4.88	3.60	8.23
						58.58	4.83	3.50	8.19

L-Val	212–214	50	0.63	$\mathrm{C}_{21}\mathrm{H}_{23}\mathrm{NO_6S}$	60.43 60.40	5.52 5.48	3.36 2 20	7.67
	L-Leu 194–196	60	0.77	$\mathrm{C}_{22}\mathrm{H}_{25}\mathrm{NO_6S}$	61.25	5.80 5.80	0.00 3.25	7.42
					61.21	5.76	3.18	7.33
al-OMe	214 - 216	99	0.80	$ m C_{22}H_{25}NO_6S$	61.25	5.80	3.25	7.42
					61.19	5.77	3.16	7.38
eu-OMe	196 - 198	67	0.84	$\mathrm{C}_{23}\mathrm{H}_{27}\mathrm{NO_6S}$	62.02	6.07	3.15	7.19
					62.00	6.00	3.00	7.00
$h_{3}$ -N $_{2}$ H $_{3}$	222 - 224	71	0.76	${ m C}_{21}{ m H}_{25}{ m N}_{3}{ m O}_{5}{ m S}$	58.47	5.80	9.74	7.42
					58.41	5.77	9.71	7.38
${ m eu-N_2H_3}$	230 - 232	69	0.74	${ m C}_{22}{ m H}_{27}{ m N}_{3}{ m O}_{5}{ m S}$	59.32	6.07	9.44	7.19
					59.30	6.00	9.33	7.00
vla-Gly-OMe	170 - 172	58	0.98	$\mathrm{C}_{22}\mathrm{H}_{24}\mathrm{N}_{2}\mathrm{O}_{7}\mathrm{S}$	57.39	5.22	6.09	6.96
					57.33	5.20	6.00	6.81
al-Gly-OMe	148 - 150	52	0.92	$\mathrm{C}_{24}\mathrm{H}_{28}\mathrm{N}_{2}\mathrm{O}_{7}\mathrm{S}$	59.02	5.74	5.74	6.56
					59.00	5.61	5.62	6.44
au-Gly-OMe	162 - 164	88	0.75	${ m C}_{25}{ m H}_{30}{ m N}_2{ m O}_7{ m S}$	59.76	5.98	5.58	6.37
					59.66	5.91	5.53	6.30
DL-Ala	170 - 172	59	0.81	$ m C_{16}H_{15}NO_5S_2$	52.60	4.11	3.83	17.53
					52.55	4.00	3.81	17.51
L-Val	151 - 153	49	0.78	$\mathrm{C_{18}H_{19}NO_5S_2}$	54.96	4.83	3.56	16.23
					54.91	4.81	3.52	16.28
L-Leu	162 - 164	58	0.76	$ m C_{19}H_{21}NO_5S_2$	56.02	5.16	3.44	15.72
					56.00	5.00	3.41	15.70
						(Coni	inued on r	ext page)

7
Ξ
<u>o</u>
$\sim$
<u> </u>
5
×
4
Е
5
×.
×.
č.
н
<b>^</b>
11
2
$\sim$
ž
$\mathbf{C}$
·~
41
· · · ·
÷
а
_
÷
ŝ
e
:=:
ίΩ.
<u>9</u>
e
>
·=
_ <b>=</b>
N
•=
9
e
Ċ.
Ň
7
<,
نب
5
5
Ч,
5
2
Ĕ
2
3
0
T
5
Ô
č.
( )

TABLE I Physical Data for the Chalconesulfonyl, Amino Acid Derivatives 2–13, 16–25, and 28–38(Continued)

i

ī

						Element	tal Analys	is % Calcula	ted/Found
Jompound Vo.	R	M.P. °C	$_\%^{\rm Yield}$	${ m R_{f}}$	Molecular Formula	% C	Н %	N %	% S
31	$ m N\text{-}Et_2$	94–96	53	0.61	$\mathrm{C}_{17}\mathrm{H}_{17}\mathrm{NO}_3\mathrm{S}_2$	58.45	5.44	4.01	18.34
0	DI ALCOMO	111 001	62			58.41 59.00	5.42	4.00 2.60	18.34
4	AIMO-BIN-TIC	TTT-POT	20		017111710502	53.80	4.43 4.44	3.63	16.81
33	L-Val-OMe	117 - 119	50	0.71	$ m C_{19}H_{21}NO_5S_2$	56.02	5.16	3.44	15.72
						56.00	5.10	3.41	15.70
34	L-Leu-OMe	129 - 131	56	0.77	$ m C_{20}H_{33}NO_5S_2$	57.01	5.46	3.33	15.20
						57.00	5.41	3.30	15.00
35	$L-Val-N_2H_3$	100 - 102	69	0.75	$ m C_{18}H_{21}N_{3}O_{4}S_{2}$	53.07	5.16	10.32	15.72
						53.00	5.11	10.31	15.71
36	$ m L-Leu-N_2H_3$	112 - 114	71	0.80	$ m C_{19}H_{23}N_{3}O_{4}S_{2}$	54.16	5.46	9.98	15.20
						54.00	5.41	9.91	15.13
37	L-Val-Gly-OMe	208 - 210	62	0.84	$C_{21}H_{24}N_2O_6S_2$	54.31	5.17	6.03	13.79
						54.31	5.11	6.00	13.71
38	L-Leu-Gly-OMe	200 - 202	79	0.87	$ m C_{22}H_{26}N_2O_6S_2$	55.23	5.44	5.86	13.39
						55.21	5.41	5.81	13.33

of sulfonyl chlorides (0.1 mol) during 30 min. The temperature of the reaction mixture during the process of the addition was kept at 10°C. Stirring continued for 4 h at 20°C. Tetrahydrofuran was removed by concentration of the reaction mixture under reduced pressure. Water (50 mL) was then added and acidified with 2M HC1 to pH5. The crude products were filtered and recrystallized (ethanol-water). All of products **3–5**, **16–18**, and **28–31** were chromatographicallv homogeneous by iodine and benzidine development in TLC, Table I and Chart 1.

IR of 4:	$v3400 \text{ cm}^{-1}$ (NH), $v1690 \text{ cm}^{-1}$ (CO), $v1420$ , 1360
	$cm^{-1}$ (SO <sub>2</sub> -NH), v1600 cm <sup>-1</sup> (Ar-C=C), v1320,
	$1160 \text{ cm}^{-1} (\text{SO}_2)$
IR of 17:	$\nu$ 1680 cm <sup>-1</sup> (CO), $\nu$ 1600 cm <sup>-1</sup> (Ar-C=C), $\nu$ 1320,
	$1180 \text{ cm}^{-1} (\text{SO}_2)$
IR of 30:	$\nu$ 3200 cm <sup>-1</sup> (NH), $\nu$ 1680 cm <sup>-1</sup> (CO), $\nu$ 1610 cm <sup>-1</sup>
	(Ar-C=C), $v1350$ , 1160 cm <sup>-1</sup> (SO <sub>2</sub> )
<sup>1</sup> H NMR of 4:	$(DMSO-d_6): \delta 1.1-1.2 (s, 3H, \underline{CH}_3), \delta 4.2-4.6 (s, \alpha H,$
	<u>CH</u> ), $\delta 4.7 - 5.2$ (s, H, <u>NH</u> ), $\delta 7.8 - 7.7$ (9H, ArH), $\delta 8.1$ ,
	7.4 (alkenic $H_a$ , $H_b$ ), $\delta 10.9$ (s, H, COOH). <b>MS of 4:</b>
	$m/z 359 (M^+)$
<sup>1</sup> H NMR of 17:	$(DMSO-d_6): \delta 0.9[s, 6H, (CH_3)_2], \delta 1.97 (s, H, \beta CH),$
	$\delta 4.26$ (s, H, $\alpha$ CH), $\delta 3.97$ (s, 3H, O–CH <sub>3</sub> ), $\delta 4.8-5.2$
	(s, H, NH), $\delta 8.30-7.03$ (10H, ArH and alkenic H <sub>a</sub> ,
	$H_{b}$ , $(H_{a})$ 7.79, 7.72, $(H_{b})$ 7.53, 7.49, $\delta 10.9$ (s, H,
	COOH). <b>MS of 17:</b> m/z 417 (M <sup>+</sup> )
<sup>1</sup> H NMR of 30:	$(DMSO-d_6): \delta 0.9[s, 6H, (CH_3)_2], \delta 4.5(s, 2H, CH_2), $
	$\delta$ 8.4–7.5 (9H, ArH), and alkenic H <sub>a</sub> , H <sub>b</sub> , $\delta$ 9.8 (s, H,
	COOH), δ10.49 (s, H, SO <sub>2</sub> NH). <b>MS of 30:</b> m/z 407
	$(\mathrm{M}^+)$
<sup>1</sup> H NMR of 31:	(DMSO-d <sub>6</sub> ): $\delta$ 0.9[s, 6H, (CH <sub>3</sub> ) <sub>2</sub> ], $\delta$ 8.5–7.5(9H, ArH
	and CH=CH), $\delta H_a$ , 7.72, $\delta H_b$ 7.53, $\delta$ 7.7 (s, 2H, thio-
	phene 3,4-H), $\delta$ 3.4–3.2 (4H, N · CH <sub>2</sub> · <u>CH<sub>3</sub></u> ), $\delta$ 1.3–
	1.15(6H, N-CH <sub>2</sub> - <u>CH<sub>3</sub></u> ). <b>MS of 31:</b> 349 (M <sup>+</sup> ), 277
	$(M-NEt_2)$ , signals removed by $D_2O$ tratment 213
	$(M-SO_2-NEt2).$

# Synthesis of Chalcone Sulfonylamino Acid Methyl Esters 7–8, 19–20, and 33–34: General Procedure

A suspension of coupling reaction products **5–6**, **17**, **18 28**, and **30** (0.2 mole) in absolute methanol (100 mL) was cooled to  $-10^{\circ}$ C, and pure

thionyl chloride (2.2 mL) was added dropwise during 1 h. The reaction mixture was stirred for an additonal 3–4 h at r.t. after standing overnight, and the solvent was removed by vacuum distillation. The residual solid material was recrystallized (methanol-water). (cf., Table I, Chart 1).

IR of 8:	$\nu$ 3300, 3180 cm <sup>-1</sup> (NH), $\nu$ 1600, 1610 cm <sup>-1</sup> (CO),
	$\nu$ 1370, 1170 cm <sup>-1</sup> (SO <sub>2</sub> -NH), $\nu$ 1610 cm <sup>-1</sup> (ArC-C),
	$\nu$ 1350, 1150 cm <sup>-1</sup> (SO <sub>2</sub> ), $\nu$ 1360 cm <sup>-1</sup> (COOCH <sub>3</sub> )
<sup>1</sup> H NMR of 19:	(DMSO-d <sub>6</sub> ): $\delta 8.2-7.0$ (Ar-H and alkenic H <sub>a</sub> , H <sub>b</sub> ),
	$\delta 3.51 - 3.41(s, 3H, COOCH_3)$
<sup>1</sup> H NMR of 32:	(DMSO-d <sub>6</sub> ): δ8.04(s, H, <u>NH</u> ), δ3.75(s, 3H, COOCH <sub>3</sub> ),
	$\delta 4.4(H, \alpha \underline{CH}), \delta 1.2(3H, \beta \underline{CH}_3)$ absence of OH proton,
	and other peak in support of their structure

# Synthesis of Chalcone Sulfonylamino Acid Hydrazides 9, 10, 19, 20, and 35, 36: General Procedure

The methyl esters **7**, **8**, **19**, **20**, **33**, and **34** (0.2 mol) were dissolved in ethanol (100 mL), and hydrazine hydrate 85% (0.2 mol) was added. The reaction mixture was stirred for 3 h at 20°C and left for 24 h at r. t. The crystalline products **19**, **10**, **19**, **20**, **35**, and **36** were filtered off, washed with water, and recrystallized (ethanol–water).

IR of 35:	$\nu$ 3340, 3125 cm <sup>-1</sup> (NH), $\nu$ 1640 cm <sup>-1</sup> (CO), $\nu$ 1600,
	1550 cm <sup>-1</sup> Ar C=C, $\nu$ 1340, 1180 cm <sup>-1</sup> (SO <sub>2</sub> )
<sup>1</sup> H NMR of 36:	(DMSO-d <sub>6</sub> ): δ9(H, SO <sub>2</sub> <u>NH</u> ), δ8.2–7.5(Ar-H), δ5.50 (H,
	$\underline{\text{NH}}$ ), $\delta 5.61 (2\text{H}, \underline{\text{NH}}_2)$

# Synthesis of Chalcone Sulfonylamino Dipeptide Methyl Esters 11–13, 23–25, and 37–38: General Procedure

To a solution of amino acid methyl ester hydrochloride (0.11 mol) in THF (100 mL) was added trethylamine (5 mL). The solution was stirred at 20°C for 30 min and cooled to 0°C, where the sulfonyl amino acid (0.001 mol) and dicyclohexylcarbodiimide DCC (1.62 g) were added to the mixture. The reaction mixture was stirred for 2 h at 0°C and for another 2 h at r.t. The precipitated dicyclohexylurea was filtered off, and acetic acid (2 mL) was added to the solution, which was left standing overnight. A precipitate was filtered off and the remaining solution was distilled under vacuum. The remaining solid was recrystallized (ethanol-water). The products were chromatographically homogeneous.

IR of 12:	$v3300, 3100 \text{ cm}^{-1}$ (NH, CONH), $v1750 \text{ cm}^{-1}$ (CO),
	$v1320 \text{ cm}^{-1} (\text{COOCH}_3)$
<sup>1</sup> H NMR of 12:	$(DMSO-d_6): \delta 7.82 (s, 2H, SO_2 \underline{NH}), \delta 8.04 (s, H, CO \underline{NH}),$
	δ3.8(s, 3H, COOCH <sub>3</sub> ), δ3.65(s, 2H, <u>CH<sub>2</sub></u> ), δ4.32(s, H,
	$\alpha$ - <u>CH</u> -alalyl) $\delta$ 1.21(s, 3H, CH <sub>3</sub> alalyl) and other bands
	supporting the structure of dipeptide
<sup>1</sup> H NMR of 37:	(DMSO-d <sub>6</sub> ): δ3.84(s, 3H, COOCH <sub>3</sub> ), δ4.26(s, H, α <u>CH</u>
	valyl), $\delta$ 1.79(s, H, $\beta$ CH-valyl), $\delta$ 0.95[s, 6H, (CH <sub>3</sub> ) <sub>2</sub> ],
	$\delta 3.65(s, 2H, CH_2)$ and other bands supporting the
	structure of dipeptide

### REFERENCES

- [1] R. A. El-Sayed, J. Serb. Chem. Soc., 56, 311 (1991).
- [2] R. A. El-Sayed, N. S. Khalaf, F. A. Kora, and Y. A. Abbas, Pak. J. Sci. Industrial Research, 34, 369 (1991).
- [3] R. A. El-Sayed, N. S. Khalaf, F. A. Kora, and M. F. Badie, J. Chem. Soc. Pak., 14 49, (1992).
- [4] R. A. El-Sayed, N. S. Khalaf, F. A. Kora, and M. H. El-Haki, Proc. Ind. Nat. Sci. Acad., 58, 389 (1992).
- [5] R. A. El-Sayed, J. Ind. Chem. Soc., 69, 618 (1992).
- [6] R. A. El-Sayed, N. S. Khalaf, F. A. Kora, and M. A. El-Gazzar, J. Serb. Chem. Soc., 59, 727 (1994).
- [7] R. A. El-Sayed, Ind. Chem. Soc., 75, 53 (1998)
- [8] R. A. El-Sayed, Phosphorus, Sulfur, and Silicon, 131, 207 (1997).
- [9] R. A. El-Sayed, Chemistry of Heterocyclic Compounds, 7, 821 (1998).
- [10] R. A. El-Sayed, Chemistry of Heterocyclic Compounds, 1, 95 (2001).
- [11] R. A. El-Sayed, Phosphorus, Sulfur, and Silicon, 179, 237 (2004).
- [12] R. A. El-Sayed, Phosphorus, Sulfur, and Silicon, 181, (2006).
- [13] W. B. Geiger and J. E. Conn, J. Amer. Chem. Soc., 67, 112 (1945).
- [14] D. H. Marrian, P. B. Russell, and A. R. Todd, J. Chem. Soc., 1419 (1947).
- [15] D. N. Dhar, Chemistry of Chalcenes, (Wiley, New York, 1981).
- [16] L. Weinstein, Sulfonamides in the Pharmacological Basic of Therapeutics L. S. Goodman and A. Goodman, Eds. (Macmillan, London, and New York, pp. 1177–1203, 1970).
- [17] R. J. Cremyln, J. Chem. Soc., 1132 (1965).
- [18] R. J. Cremyln, J. Chem. Soc., 77 (1967).
- [19] R. J. Cremyln, J. Chem. Soc. (C), 1341 (1969).
- [20] P. Pfeiffer and P. A. Negreanu, Ber., 50, 1465 (1917).
- [21] K. K. Hsu, M. S. Lo, and F. C. Chen, J. Chinese Chem. Soc., 16, 91 (1969); Chem. Abs., 72, 31393V (1970).
- [22] Vogels, Practical Organic Chemistry, 5th, ed., pp. 1034 (Longnnan, Harlow, 1989).
- [23] N. R. El-Rayyes, G. H. Havakeemian, and H. S. Hmoud, J. Chem. Eng. Data., 29, 225 (1984).
- [24] E. Pretch, T. Clero, J. Seibe, and W. Simon, Spectral Data for Structure Determination of Organic Compounds, (Springer-Verleg, Berlin, 1989).
- [25] F. J. Swinbourne and O. O. Shode, J. Chinese Chem. Soc., 31, 383 (1984).
- [26] F. J. Swinbourne, P. Bassin, D. Dane, and Mitchell, *Phosphorus, Sulfur, and Silicon*, 63, 385 (1991).