

Synthesis of (–)-Isofagomine

Irma Panfil

Institute of Organic Chemistry of the Polish Academy of Sciences, Warsaw, Poland

Jolanta Solecka

National Institute of Hygiene, Warsaw, Poland

Marek Chmielewski

Institute of Organic Chemistry of the Polish Academy of Sciences, Warsaw, Poland

1,3-Dipolar cycloaddition of *N*-benzyl nitrone **2** to *D*-threo δ -lactone **15** proceeded with excellent stereoselectivity to provide only one adduct **16**. Cycloadduct **16** was subsequently subjected to a sequence of reactions involving rearrangement to γ -lactone, glycolic cleavage/reduction, protection of the terminal hydroxymethyl group, reduction of the lactone, desilylation/mesylation, and hydrogenolysis of the N-O bond providing (–)-isofagomine and its *N*-substituted derivatives. The biologic activity of *N*-substituted (–)-isofagomines toward commercially available α - and β -glucosidases, α -D-mannosidase, α -L-fucosidase, β -D-glucuronidase, and β -D-galactosidase was tested.

Keywords (–)-Isofagomine, Iminosugars, 1,3-Dipolar cycloaddition, Nitrones, Unsaturated aldolactones

INTRODUCTION

Glycosidases and glycosyl transferases play an important role as carbohydrate-modifying enzymes. Carbohydrate mimics, such as iminosugars, carbasugars, or *C*-glycosides, display properties as specific and competitive inhibitors of these enzymes.^[1] Consequently, they display many interesting biologic and therapeutic activities such as antiviral, anticancer, antidiabetic, and antihyperglycemic.^[1–7] The discovery of isofagomine (**1**) by a Danish group^[8] as one of the most potent β -glucosidase inhibitors has propelled, during the past years, a growing interest of many laboratories in the synthesis of iminosugars with nitrogen at the anomeric position.^[9–21]

Received August 28, 2006; accepted November 9, 2006.

Address correspondence to Marek Chmielewski, Institute of Organic Chemistry of the Polish Academy of Sciences, 01-224, Warsaw, Poland. E-mail: chmiel@icho.edu.pl

Recently we reported on 1,3-dipolar cycloaddition of *N*-benzyl nitrene **2** to *D*-glycero γ -lactone **3**, *L*-erythro δ -lactone **4**, and *D*-threo δ -lactone **5**.^[22] The cycloaddition performed with **4** afforded two stereoisomers **7** and **8** in a ratio of about 2.5:1. Cycloaddition of the same nitrene **2** to lactones **3** and **5** yielded, in each case as a sole product, **6** and **9**, respectively (Chart 1).

Cycloadducts **6–9** were subsequently subjected to a sequence of reactions involving hydrogenolysis of the N-O bond and intramolecular alkylation of the nitrogen atom by the C-4 or C-5 carbon atom of the sugar backbone to afford iminosugars **10–14** with a hydroxymethyl group at the C-2 carbon atom.^[23] This strategy proceeded with inversion of configuration at the secondary carbon atom (C-4 or C-5) of the starting sugar. In addition, the interchange of the C-1 carbon atom by the aminomethyl group caused inversion of configuration (according to the carbohydrate nomenclature) at the tertiary carbon atom (C-2).^[16]

RESULTS AND DISCUSSION

In all cases, the high preference for the *anti* addition of the nitrene to the terminal methyl or substituted methyl groups in lactones was observed.^[22–25] Low stereoselectivity of cycloadditions of simple nitrenes to *erythro* lactones




Chart 1

of **26** followed by hydrogenolysis of the *N*-benzyl substituent afforded (–)-isofagomine (**1-ent**). Compound **1-ent** and its *N*-benzyl derivative **25** have been synthesized before from 1-tert-butoxycarbonyl-1,2,3,6-tetrahydropyridyl-3-methanol^[13] and (+)-L-tartaric acid (Scheme 1).^[15b]

The biologic activity of compounds **1-ent**, **25** and **28**, toward commercially available α - and β -glucosidases, α -D-mannosidase, α -L-fucosidase, β -D-glucuronidase, and β -D-galactosidase were tested. (–)-*N*-Acetyl-isofagomine (**28**) was found to be a better inhibitor of tested enzymes than (–)-isofagomine (**1-ent**). It showed moderate/weak activity against bovine kidney α -L-fucosidase (50% inhibition required 3.5 mM), bovine liver β -D-galactosidase (50% inhibition required at 6.6 mM), and bovine liver β -D-glucuronidase (50% inhibition required at 9.1 mM), and little or no activity against rice α -, almond β -D-glucosidases, and jack bean α -D-mannosidase. (–)-Isifagomine (**1-ent**) showed moderate/weak activity against α -L-fucosidase, β -D-galactosidase, and β -D-glucuronidase, and a little or no activity against α -D-glucosidases and α -D-mannosidase. (–)-*N*-Benzyl-isofagomine (**25**) showed only inhibition of α -L-fucosidase (50% inhibition at 5.1 mM).

CONCLUSION

It was demonstrated that 1,3-dipolar cycloaddition of the simple *N*-benzyl nitron to unsaturated *threo* 1,5-lactone **15** proceeded exclusively *anti* to the substituents in the sugar. The adduct **16** after rearrangement to the γ -lactone **17** and removal of the terminal hydroxymethyl group could be the attractive substrate for the synthesis of enantiomer of natural isofagomine (**1-ent**). Examination of the glycosidases' inhibitory activity showed that (–)-*N*-acetyl-isofagomine (**28**) was a better inhibitor of tested enzymes than (–)-isofagomine (**1-ent**).

EXPERIMENTAL

Melting points were determined on a Koeffler hot-stage apparatus. ¹H NMR spectra were recorded using Bruker Avance 500 and Varian Mercury 400 instruments. Absorptions of aromatic protons were not reported. IR spectra were recorded on a Perkin-Elmer FTIR Spectrum 200 spectrophotometer. Mass spectra were recorded using AMD-604 Inectra GmbH and HPLC-MS with Mariner and API 356 detectors. Optical rotations were measured using a JASCO P 3010 polarimeter at 22 ± 3°C. Column chromatography was performed using E. Merck Kiesel Gel (230–400 mesh).

(3aR, 6R, 7R, 7aS)-7-Acetoxy-6-acetoxymethyl-2-benzyl-4-oxo-tetrahydropyrano[3,4-d]isoxazolidine (16). Compound **16** was obtained according to the procedure described earlier.^[22] Syrup; $[\alpha]_D + 44.2$ (c 1.5, CH₂Cl₂); IR

(film): 1748 cm^{-1} ; ^1NMR (500 MHz, toluene- d_8 , 100°C) δ : 5.09 (ddt, 1H, J 1.6, 3.3 Hz, H-7), 4.88 (dt, 1H, J 1.6, 6.3 Hz, H-6), 4.11 (dd, 1H, J 3.3, 8.0 Hz, H-7a), 4.11 (dd, 1H, J 6.3, 11.4 Hz, CHHOAc), 4.09 (dd, 1H, J 6.3, 11.4 Hz, CHHOAc), 3.69, 3.63 (2d, 2H, J 13.3 Hz, NBn), 3.22 (dt, 1H, J 5.4, 8.0 Hz, H-3a), 2.89 (bs, 1H, H-2), 2.83 (bt, 1H, H-2'); HRMS (ESI) m/z $[\text{M} + \text{Na}]^+$, Calcd. for $\text{C}_{18}\text{H}_{21}\text{NO}_7\text{Na}$: 386.1210. Found: 386.1230.

(3aR, 6R, 6aS, 1'R)-2-Benzyl-6-(1',2'-dihydroxyethyl)-4-oxo-isoxazolidino[4,5-c]tetrahydrofuran (17). Adduct **16** (0.36 g, 1 mmol) was dissolved in methanol (10 mL) containing 1.5% of ammonia and left at rt for 24 h. Subsequently, solvent was evaporated and the crude product was purified by chromatography using AcOEt as an eluent to afford **17** (0.26, 95%). m.p. 95–97°C; $[\alpha]_{\text{D}} + 25.2$ (c 0.5, CH_2Cl_2); IR (film): 3588, 1777 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 4.99 (bt, 1H, H-6a), 4.65 (dd, 1H, J 3.7, 6.5 Hz, H-6), 4.10, 3.89 (2d, 2H, J 12.4 Hz, NBn), 4.06 (dd, 1H, J 5.1, 9.6 Hz, CHHOH), 3.74–3.56 (m, 4H, CHOH , CHHOH , H-3, H-3a), 2.82 (bm, 1H, H-3'); HRMS (LSIMS) m/z $[\text{M} + \text{H}]^+$, Calcd. for $\text{C}_{14}\text{H}_{18}\text{NO}_5$: 280.1185. Found: 280.1197.

(3aR, 6R, 6aS, 1'R)-6-(1',2'-Diacetoxyethyl)-2-benzyl-4-oxo-isoxazolidino[4,5-c]tetrahydrofuran (17Ac). Syrup, $[\alpha]_{\text{D}} + 1.15$ (c 0.8, CH_2Cl_2); IR (film): 1779, 1744 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, toluene- d_8) δ : 5.50 (ddd, 1H, J 3.1, 4.2, 8.9 Hz, CHOAc), 4.22 (dd, 1H, J 3.1, 12.4 Hz, CHHOAc), 4.20 (dd, 1H, J 5.4, 7.1 Hz, H-6a), 4.16 (dd, 1H, J 5.4, 8.9 Hz, H-6), 4.10 (dd, 1H, J 4.2, 12.4 Hz, CHHOAc), 3.63, 3.58 (2d, 2H, J 13.2 Hz, NBn), 3.17 (bd, 1H, J 9.3 Hz, H-3), 2.74 (t, 1H, J 7.1 Hz, H-3a), 2.33 (bt, 1H, H-3') 1.76, 1.70 (2s, 6H, 2Ac); HRMS (ESI) m/z $[\text{M} + \text{Na}]^+$, Calcd. for $\text{C}_{18}\text{H}_{21}\text{NO}_7\text{Na}$: 386.1210. Found: 386.1232.

(3aR, 6R, 6aR)-2-Benzyl-6-hydroxymethyl-4-oxo-isoxazolidino[4,5-c]tetrahydrofuran (18). Compound **17** (0.36 g, 1.0 mmol) was dissolved in methanol/water 2:1 v/v (15 mL) and treated with sodium metaperiodate at rt (0.24 g, 1.1 mmol). After 1.5 h the reaction was completed. After evaporation of the solvent, water was added into the flask and the resulting solution was extracted with AcOEt. Subsequently, the extract was evaporated and the residue was dissolved in dry CH_2Cl_2 (15 mL) and reduced with 1.5 molar equiv. of $\text{NaBH}(\text{OAc})_3$ (0.29 g). Standard workup provided the product, which was purified on a silica gel column using AcOEt as an eluent to afford **18** (0.16 g, 60%). Syrup; $[\alpha]_{\text{D}} + 31.0$ (c 0.84, CH_2Cl_2); IR (film): 3564, 1778 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ : 4.94 (bt, 1H, H-6a), 4.22 (dt, 1H, J 5.1, 5.8, 5.8 Hz, H-6), 3.99, 3.95 (2bd, 2H, J 12.9 Hz, Bn), 3.89 (dd, 1H J 4.7, 12.3 Hz, CHHOH), 3.83 (dd, 1H J 5.6, 12.3 Hz, CHHOH), 3.66–3.54 (m, 2H, H-3, H-3'), 2.80 (bs, 1H, H-3a); HRMS (EI) m/z M^+ , Calcd. for $\text{C}_{13}\text{H}_{15}\text{NO}_4$: 249.1001. Found: 249.1007.

(3aR, 6R, 6aR)-6-Acetoxymethyl-2-benzyl-4-oxo-isoxazolidino[4,5-c]tetrahydrofuran (18Ac). m.p. 108–110°C; $[\alpha]_{\text{D}} - 5.14$ (c 1.0, CH_2Cl_2); IR

(film): 1778, 1741 cm^{-1} ; ^1H NMR (500 MHz, toluene- d_8 , 100°C) δ : 4.32 (dd, 1H, J 4.1, 12.2 Hz, CHHOAc), 4.24 (dd, 1H, J 7.6, 12.2 Hz, CHHOAc), 4.15 (ddd, 1H, J 4.1, 5.5, 7.6 Hz, H-6), 4.11 (dd, 1H, J 5.5, 7.2 Hz, H-6a), 3.66, 3.58 (2d, 2H, J 13.2 Hz, NBn), 3.14 (bd, 1H, J 9.3 Hz, H-3), 2.71 (dt, 1H, J 1.6 7.2 Hz, H-3a), 2.31 (bt, 1H, H-3'), 1.69 (s, 3H, Ac); HRMS (LSIMS) m/z $[\text{M} + \text{H}]^+$, Calcd. for $\text{C}_{15}\text{H}_{18}\text{NO}_5$: 292.1185. Found: 292.1192.

(3aR, 6R, 6aR)-2-Benzyl-6-*t*-butyldiphenylsiloxymethyl-4-oxo-isoxazolidino[4,5-*c*]tetrahydrofurane (19). Compound **18** (0.25 g, 1 mmol) dissolved in CH_2Cl_2 (15 mL) was treated with *t*-butyldiphenylchlorosilane (0.302, 1.1 mmol) and DMAP (1.1 mmol). The mixture was left overnight. Subsequently, the solvent was evaporated and the residue was purified by chromatography using hexane/AcOEt 4:1 v/v as an eluent to give **19** (0.35 g, 73%). Syrup, $[\alpha]_D + 26.3$ (c 0.9, CH_2Cl_2); IR (film): 1772 cm^{-1} ; ^1H NMR (500 MHz, toluene- d_8 , 100°C) δ : 4.22–4.16 (m, 2H, H-6, H-6a), 4.03 (dd, 1H, J 6.4, 11.1 Hz, CHHOTBDPS), 3.98 (dd, 1H, J 5.1, 11.1 Hz, CHHOTBDPS), 3.57, 3.54 (2d, 2H, J 13.3 Hz, NBn), 3.13 (bd, 1H, J 9.3 Hz, H-3), 2.73 (dt, 1H, J 1.7, 7.0 Hz, H-3a), 2.33 (bt, 1H, H-3'), 1.14 (s, 9H, *t*-Bu); HRMS (ESI) m/z $[\text{M} + \text{H}]^+$, Calcd. for $\text{C}_{29}\text{H}_{33}\text{NO}_4\text{Si}$: 488.2252. Found: 488.2251.

(3aR, 8R, 8aR)-2-Benzyl-8-*tert*-butyldiphenylsiloxymethyl-6,6-dimethyl-5,7-dioxa-isoxazolidino[4,5]cycloheptane (21). Compound **18** (0.49 g, 1 mmol) was dissolved in dry THF and treated with LiAlH_4 (0.04 g, 1 mmol). The mixture was stirred for 0.5 h and then, consecutively, AcOEt (1 mL) and water (1 mL) were added. Subsequently, the mixture was filtered through Celite and solvents were evaporated. The residue was dissolved in brine (5 mL) and extracted with AcOEt. The extract was dried and evaporated. The residue (crude **20**) was dissolved in dimethoxypropane (10 mL), treated with catalytic amount of *p*-TsOH, and refluxed for 2 h. Subsequently, the mixture was neutralized with Na_2CO_3 and evaporated. The crude product was purified by chromatography using hexane/AcOEt 7:3 v/v as an eluent to afford **21** (0.38 g, 71%). m.p. 84–86°C; $[\alpha]_D + 24.5$ (c 1.0, CH_2Cl_2); ^1H NMR (500 MHz, toluene- d_8 , 100°C) δ : 4.17–4.12 (m, 2H, H-8, H-8a), 3.95 (d, 2H, CH_2OTBDPS), 3.87 (t, 1H, J 10.4, 11.9 Hz, H-4), 3.84, 3.82 (2d, 2H, J 13.1 Hz, NBn), 3.20 (dd, 1H, J 5.2, 11.9 Hz, H-4'), 2.76 (bt, 1H, J 7.8, 9.1 Hz, H-3), 2.38 (m, 1H, H-3a), 2.19 (bdd, 1H, H-3'), 1.31, 1.29 (2s, 6H, 2 CH_3), 1.14 (s, 9H, *t*-Bu); HRMS (ESI) m/z $[\text{M} + \text{H}]^+$, Calcd. for $\text{C}_{32}\text{H}_{42}\text{NO}_4\text{Si}$: 532.2878. Found: 532.2860.

(3S,4S,5S) *N*-Benzyl-3,4-diacetoxy-5-(acetoxymethyl)piperidine [(-)-*N*-Benzyl-tri-*O*-acetyl-isofagomine] (26). Compound **21** (0.266 g, 0.5 mmol) was dissolved in THF (10 mL), treated with TBAF (0.16 g, 0.5 mmol), and stirred at rt for 3 h. The reaction mixture was evaporated and crude product was passed through a silica gel column using AcOEt as an

eluent. Subsequently, the solvent was evaporated and the product (**22**, 0.13 g, 90%) was dissolved in CH₂Cl₂ (10 mL), treated with Et₃N (0.06 g) and mesyl chloride (0.06 g, 0.5 mmol), and left for 1 h at rt. The mixture was then washed with water and evaporated. The product was dissolved in methanol (10 mL) and acetic acid (2 mL) was added, the resulting solution was refluxed for 10 min and evaporated, and the residue acetylated with acetic anhydride/pyridine 1:1 mixture (2 mL). After standard workup and evaporation, the crude mixture was dissolved in methanol (15 mL), treated with K₂CO₃ (20 mg), and hydrogenated using Degussa 10% Pd/C at rt for 24 h. The mixture was filtered through Celite, evaporated, and acetylated using 1:1 acetic anhydride/pyridine mixture. After standard workup, the crude product was purified by chromatography using hexane/AcOEt 1:1 v/v mixture to afford **26** (0.07 g, 50%). Syrup, [α]_D – 10.5 (*c* 0.9, CH₂Cl₂); IR (CHCl₃): 1738 cm^{–1}; ¹H NMR (500 MHz, toluene-d₈, 100°C) δ : 5.07 (ddd, 1H, *J* 5.0, 8.9, 9.7 Hz, H-3), 4.91 (dd, 1H, *J* 8.9, 10.1 Hz, H-4) 3.98 (dd, 1H, *J* 6.2, 11.4 Hz, CHHOAc), 3.83 (dd, 1H, *J* 3.8, 11.4 Hz, CHHOAc), 2.96 (ddd, 1H, *J* 2.1, 5.0, 10.9 Hz, H-2), 2.68 (ddd, 1H, *J* 2.1, 4.1, 11.5 Hz, H-6), 2.02 (m, 1H, H-5), 1.97 (dd, 1H, *J* 9.7, 10.9 Hz, H-2'), 1.87 (t, 1H, *J* 11.1 Hz, H-6'), 1.74, 1.68, 1.67 (3s, 12H, 3Ac); HRMS (ESI) *m/z* [M + H]⁺, Calcd. for C₁₉H₂₆NO₆: 364.1755. Found: 364.1748.

(–)-*N*-Benzyl-isogagomine (**25**). Deacetylation of **26** with 1% of ammonia in methanol gave **25** (90%). Colorless crystals; m.p. 147–150°C; [α]_D – 16.5 (*c* 0.4, CH₂Cl₂), lit.^[15b] [α]_D – 13.2 (*c* 1.1, ethanol); ¹H NMR (400 MHz, methanol-d₄) δ : 3.78 (dd, 1H, *J* 3.7, 11.0 Hz, CHHOH), 3.58, 3.53 (2d, 2H, *J* 12.8 Hz, Bn), 3.51 (m, 1H, H-3), 3.49 (dd, 1H, *J* 6.9, 11.0 Hz, CHHOH), 3.07 (dd, 1H, *J* 8.8, 10.3 Hz, H-4), 3.01 (ddd, 1H, *J* 2.3, 3.8, 11.4 Hz, H-2), 2.96 (ddd, 1H, *J* 2.3, 4.8, 10.8 Hz, H-3), 1.89 (t, 1H, *J* 11.4 Hz, H-2'), 1.86 (t, 1H, *J* 10.8 Hz, H-3'), 1.76 (m, 1H, H-5); HRMS (ESI) *m/z* [M + H]⁺, Calcd. for C₁₃H₂₀NO₃: 238.1438. Found: 238.1428.

(–)-*N*-Acetyl-tri-*O*-acetyl-isogagomine (**27**). Compound **26** was hydrogenated in methanol under standard condition. After evaporation of the solvent, the residue was acetylated with acetic anhydride pyridine mixture, which after evaporation was purified on a silica gel column using AcOEt/MeOH 4:1 v/v as an eluent to afford **27** (90%) as a mixture of two rotamers in a ratio of about 2:1. Colorless crystals; m.p. 89–90°C; [α]_D – 14.7 (*c* 0.9, CH₂Cl₂); IR (film): 1774, 1649 cm^{–1}; ¹H NMR (500 MHz, CDCl₃) δ : major rotamer 5.03 (bm, 1H, H-3), 4.73 (bm, 1H, H-4), 4.49 (bd, 1H, CHHOAc), 4.09 (m, 1H, CHHAc), 4.03 (bm, 1H, H-2), 4.01 (bm, 2H, H-2, H-6), 3.11 (m, 1H, H-2'), 2.81 (bt, 1H, H-6'), 2.05 (m, 1H, H-5), 2.13, 2.11, 2.06, 2.05 (4s, 12H, 4Ac); minor rotamer 5.02 (bs, 1H, H-3), 4.84 (bm, 1H, H-4), 4.38 (bm, 1H, CHHOAc), 4.11 (m, 1H, CHHOAc), 3.77 (bm, 1H, H-3), 3.21 (m, 1H, H-6), 3.08 (bm, 1H, H-3'), 3.05 (m, 1H, H-6'), 1.56 (bm, 1H, H-5), 2.10, 2.09, 2.08, 2.03 (4s, 12H,

4Ac); HRMS (ESI), for the mixture of rotamers, m/z $[M + Na]^+$, Calcd. for $C_{14}H_{21}NO_7Na$: 338.1210. Found: 338.1226.

(-)-***N*-Acetyl-isofagomine (28)**. Deacetylation of compound **27** with 1% of ammonia in methanol gave **28** (90%) as a mixture of two rotamers in a ratio of about 1:1. Colorless crystals; m.p. 60–62°C; $[\alpha]_D - 8.7$ (*c* 0.7, methanol); IR (film): 1637 cm^{-1} ; 1H NMR (500 MHz, methanol- d_4) δ : 4.53, 4.50 (2 m, 1H, H-4a and b), 3.95, 3.89 (2ddd, 1H, *J* 2.4, 4.4, 13.8 Hz, H-2a and b), 3.86, 3.78 (2dd, 1H, *J* 3.7, 11.1 Hz, CHHOHa and b), 3.60 (2dd, 0.5H, *J* 6.7, 11.1 Hz, CHHOHa), 3.55 (2dd, 0.5H, *J* 7.5, 11.0 Hz, CHHOHb), 3.39 (ddd, 0.5H, *J* 4.8, 8.3, 10.1 Hz, H-6a), 3.35–3.24 (m, 1.5H, H-4a and b, H-6b), 2.96 (dd, 0.5H, *J* 10.3, 13.3 Hz, H-2'a), 2.93 (2dd, 0.5H, *J* 11.5, 13.1 Hz, H-2'b), 2.56 (dd, 0.5H, *J* 11.5, 13.1 Hz, H-6'a), 2.47 (dd, 0.5H, *J* 10.3, 12.7 Hz, H-6'b), 2.11, 2.10 (2s, 3H, Ac a and b), 1.64, 1.55 (2 m, 1H, H-5a and b); HRMS (ESI), for the mixture of rotamers, m/z $[M + Na]^+$, Calcd. for $C_8H_{15}NO_4Na$: 212.0893. Found: 212.0890.

(-)-**Isfagomine (1-ent)**. Compound **1-ent** was obtained from **25** by hydrogenation in methanol in the presence of 10% Pd/C (Degussa). Standard workup and purification on a silica gel column using *i*-PrOH/ NH_3 aq 3:1 v/v as an eluent afforded **1-ent** in 90% yield. Syrup; $[\alpha]_D - 17.8$ (*c* 0.5, methanol), lit. ^[15b] $[\alpha]_D - 15.7$ (*c* 0.19, ethanol); IR (film): 3631, 3533, 3415 cm^{-1} ; 1H NMR (500 MHz, methanol- d_4) δ : 3.78 (dd, 1H, *J* 3.8, 11.0 Hz, CHHOH), 3.55 (dd, 1H, *J* 6.8, 11.0 Hz, CHHOH), 3.40 (ddd, 1H, *J* 5.0, 8.7, 10.6 Hz, H-3), 3.20 (dd, 1H, *J* 8.8, 10.2 Hz, H-4), 3.08 (m, 2H, H-2, H-6), 2.39 (dd, 1H, *J* 11.8, 12.8 Hz, H-2'), 2.37 (dd, 1H, *J* 10.7, 12.4 Hz, H-6'), 1.64 (m, 1H, H-5); HRMS (EI) m/z M^+ , Calcd. for $C_6H_{13}NO_3$: 147.0895. Found: 147.0892.

Measurements of Enzymes Inhibition

The following hydrolases were used: α -L-glucosidase from rice (type V, 63.43 U/mg, 1.34 mg/mL, Sigma), β -D-glucosidase from almonds (25.8 U/mg, 95.4% protein, Sigma), α -D-mannosidase from jack bean (6.2 mg prot./mL, 22 U/mg, Sigma), α -L-fucosidase from bovine kidney (28.0 U/mg, 0.55 mg prot./mL, Sigma), β -D-galactosidase from bovine liver (0.148 U/mg, Sigma) solution (0.562 U/mL) and β -D-glucuronidase from bovin liver (2630 U/mg, Sigma) solution (4909 U/mL) (Table 1).

These enzymes were assayed with appropriate p-nitrophenyl glycoside substrates (phenolphthalein β -glucuronide for β -glucuronidase), which were purchased from Sigma. Hydrolase activities were measured by modification of the procedures described previously.^[24–29] The reaction mixtures (for each enzyme) consisted of 25 μ L of 16.5 mM p-nitrophenyl α -D-glucopyranoside, 403 μ L of 0.1 M acetate buffer (pH 4.0), 50 μ L of inhibitor solution (water or methanol), and 22 μ L of α -L-glucosidase solution (10 \times diluted) for

Table 1: Effect of (-)-isufagomine (**1-ent**) and (-)-*N*-acetyl-isufagomine (**28**) on various glycosidases.

Enzyme tested	(-)-Isufagomine (1-ent)		(-)- <i>N</i> -Acetyl-isufagomine (28)	
	Concentration of inhibitor (mM)	Enzyme inhibition (%)	Concentration of inhibitor (mM)	Enzyme inhibition (%)
α -L-Fucosidase	12.0	50	3.5	50
β -D-Galactosidase	8.5	50	6.6	50
β -D-Glucuronidase	15.0	50	9.1	50
α -D-Glucosidase	10.0	lack	10.0	Lack
β -D-Glucosidase	9.0	50	50.0	8.5
α -D-Mannosidase	11.0	lack	6.9	16

α -glucosidase activity; 100 μ L of 14 mM *p*-nitrophenyl β -D-glucopyranoside, 250 μ L of 0.2 M acetate buffer (pH 4.6), 50 μ L of inhibitor solution, and 100 μ L of β -D-glucosidase solution (30 μ g/mL) for β -D-glucosidase activity; 5 μ L of 20 mM *p*-nitrophenyl α -D-mannopyranoside, 200 μ L of 0.1 M acetate buffer (pH 4.5), 25 μ L of inhibitor solution, and 20 μ L of α -D-mannosidase solution (100 \times diluted) for α -D-mannosidase activity; 20 μ L of α -L-fucosidase solution (100 \times diluted), 25 μ L of inhibitor solution, 155 μ L of 0.1 M acetate buffer (pH 5.0), and 50 μ L of 20 mM 4-nitrophenyl- α -L-fucopyranoside for α -L-fucosidase activity; 90 μ L of 4 mM phenolphthalein β -D-glucuronide, 95 μ L of 0.1 M acetate buffer (pH 5.0), 25 μ L of inhibitor solution, and 40 μ L of β -D-glucuronidase solution for β -D-glucuronidase activity; 25 μ L of 20 mM 4-nitrophenyl β -D-galactopyranoside, 195 μ L of 0.1 M acetate buffer (pH 5.0), 25 μ L of inhibitor solution, and 5 μ L of β -D-galactosidase solution for β -D-galactosidase activity. Generally, enzyme and inhibitor were mixed in buffer and reaction was started by the addition of the substrate. Each reaction mixture was incubated for 30 min (α - and β -glucosidases for 15 min) at 37°C and was terminated by adding 0.250 mL of 2% sodium carbonate. The absorbance of liberated *p*-nitrophenol was measured at 405 nm (liberated phenolphthalein at 540 nm).

(-)-*N*-Benzyl-isufagomine (**25**) showed 50% inhibition of α -L-fucosidase activity in concentration 5.1 mM, trace inhibition of β -D-glucosidase activity in concentration 18 mM, and lack of inhibition of α -D-glucosidase, β -D-galactosidase, β -D-glucuronidase, and α -D-mannosidase.

REFERENCES

- [1] (a) Fleet, G.W.J.; Fellows, L.E.; Winchester, B. Plagiarizing plants: aminosugars as a class of glycosidase inhibitors. In *Bioactive Compounds from Plants*; Chadwick, P.J., March, J., Eds.; Wiley & Sons: Chichester, 1990; 112–125; (b) Asano, N.; Nash, R.J.; Molyneux, R.J.; Fleet, G.W.J. Sugar-mimic glycosidase

- inhibitors: natural occurrence, biological activity and prospects for therapeutic application. *Tetrahedron: Asymm.* **2000**, *11*, 1645–1680.
- [2] Chapleur, Y., Ed. *Carbohydrate Mimics*; Wiley-VCH: Weinheim, 1998.
- [3] Stütz, A.E., Ed. *Iminosugars as Glycosidase Inhibitors*; Wiley-VCH: Weinheim, 1999.
- [4] Martin, O.R.; Compain, P.H. Eds., Design, synthesis and biological evaluation of iminosugar-based glycosyltransferase inhibitors *Curr. Top. Med. Chem.* **2003**, *3*, 541–560, and references cited therein.
- [5] (a) El Nemr, A. Synthetic methods for the stereoisomers of Swainsonine and its analogues. *Tetrahedron* **2000**, *56*, 8579–8629; (b) Felpin, F.-X.; Lebreton, J. Recent advances in the total synthesis of piperidine and pyrrolidine natural alkaloids with ring-closing metathesis as a key step. *Eur. J. Org. Chem.* **2003**, 3693–3712; (c) Cossy, J.; Vogel, P. Hydroxylated indolizidines and their synthesis. In *Studies in Natural Products Chemistry. Stereoselective Synthesis (Part H)*; Attar-Rahman, Eds.; Elsevier: Amsterdam, **1993**; Vol. 12, 275–364.
- [6] Herczegh, P.; Kovacs, I.; Sztaricskai, F. Chemistry of biologically important hydroxylated indolizidines: synthesis of swainsonine, castanospermine and slaframine. In *Recent Progress in Chemical Synthesis of Antibiotics and Related Microbial Products*; Lukacs, G., Ohno, M., Eds.; Springer-Verlag: Berlin, 1993; 751–828.
- [7] Cipolla, L.; La Ferla, B.; Nicotra, F. General methods for iminosugar synthesis. *Curr. Top. Med. Chem.* **2003**, *3*, 485–511.
- [8] (a) Jespersen, T.M.; Dong, W.; Sierks, M.R.; Skrydstrup, T.; Lundt, I.; Bols, M. Isofagomine, a potent, new glycosidase inhibitor. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1778–1779; (b) Jespersen, T.M.; Bols, M.; Sierks, M.R.; Skrydstrup, T. Synthesis of isofagomine, a novel glycosidase inhibitor. *Tetrahedron.* **1994**, *50*, 13449–13460.
- [9] Lundt, I.; Madsen, R. Iminosugars: glycosidase inhibitors with nitrogen at the anomeric position. In *Iminosugars as Glycosidase Inhibitors*; Stütz, A.E., Ed.; Wiley-VCH: Weinheim, 1999; 112–124.
- [10] (a) Bols, M. 1-Aza sugars, apparent transition state analogues of equatorial glycoside formation/cleavage. *Acc. Chem. Res.* **1998**, *31*, 1–8; (b) Bülow, A.; Plesner, I.W.; Bols, M. A large difference in the thermodynamics of binding of Isofagomine and 1-Deoxynojirimycin to β -glucosidase. *J. Am. Chem. Soc.* **2000**, *122*, 8567–8568; (c) Liu, H.; Liang, X.; SØhoel, H.H.; Bülow, A.; Bols, M. Noeuromycin, a glycosyl cation mimic that strongly inhibits glycosidases. *J. Am. Chem. Soc.* **2001**, *123*, 5116–5117.
- [11] (a) Ichikava, Y.; Igarashi, Y.; Ichikava, M.; Suhara, Y. 1-N-Iminosugars: potent and selective inhibitors of β -glycosidases. *J. Am. Chem. Soc.* **1998**, *120*, 3007–3018; (b) Ichikava, Y.; Igarashi, Y.; Ichikava, M. Facile synthesis of glucose-type 1-N-iminosugars: new inhibitor of glycolipid biosynthesis. *Tetrahedron Lett.* **1995**, *36*, 1767–1770; (c) Ichikava, Y.; Igarashi, Y. An extremely potent inhibitor for β -galactosidase. *Tetrahedron. Lett.* **1995**, *36*, 4585–4586; (d) Igarashi, Y.; Ichikava, M.; Ichikava, Y. Synthesis of a potent inhibitor of β -glucuronidase. *Tetrahedron Lett.* **1996**, *37*, 2707–2708; (e) Igarashi, Y.; Ichikava, M.; Ichikava, Y. Synthesis of a new inhibitor of α -fucosidase. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 553–558; (f) Kim, Y.J.; Ichikava, M.; Ichikava, Y. Highly selective synthesis of 1-N-iminosugars of the D-glucose and -glucuronic acid types. *J. Org. Chem.* **2000**, *65*, 2599–2602.

- [12] (a) Hansen, A.; Tagmose, T.M.; Bols, M. Synthesis of a new potent α -fucosidase inhibitor. *J. Chem. Soc., Chem. Commun.* **1996**, 2649–2650; (b) Hansen, A.; Bols, M. Synthesis of (\pm)-isogomine and its stereoisomers from arecoline. *J. Chem. Soc. Perkin Trans. 1*, **2000**, 911–915; (c) Bols, M.; Hazell, R.G.; Thomsen, I. 1-Azafagomine: a hydroxyhexahydropyridazine that potently inhibit enzymatic glycosidase cleavage. *Chem. Eur. J.* **1997**, *3*, 940–947; (d) Liang, X.; Lohse, A.; Bols, M. Chemoenzymatic synthesis of isogalactofagomine. *J. Org. Chem.* **2000**, *65*, 7432–7437; (e) Anders, J.; Bols, M. Efficient synthesis of isogomine and Noeuromycin. *Chem. Eur. J.* **2001**, *7*, 3744–3747.
- [13] Jakobsen, P.; Lundbeck, J.M.; Kristiansen, M.; Breinholt, J.; Demuth, H.; Pawlas, J.; Torres Candela, M.P.; Andersen, B.; Westergaard, N.; Lundgren, K.; Asano, N. Iminosugars: potential inhibitors of liver glycogen phosphorylase. *Bioorg. Med. Chem.* **2001**, *9*, 733–744.
- [14] (a) Nishimura, Y.; Satoh, T.; Kudo, T.; Kondo, S.; Takeuchi, T. Synthesis and activity of 1-N-iminosugar inhibitors, siastatin B analogues for α -N-acetylgalactosaminidase and β -N-acetylglucosaminidase. *Bioorg. Med. Chem.* **1996**, *4*, 91–96; (b) Nishimura, Y.; Shitara, E.; Adachi, H.; Toyoshima, M.; Nakajima, M.; Okami, Y.; Takeuchi, T. Flexible synthesis and biological activity of uronic acid-type gem-diamine 1-N-iminosugars: a new family of glycosidase inhibitors. *J. Org. Chem.* **2000**, *65*, 2–11; (c) Shitara, E.; Nishimura, Y.; Kojima, F.; Takeuchi, T. gem-Diamine 1-N-iminosugars of L-fucose-type, the extremely potent L-fucosidase inhibitors. *Bioorg. Med. Chem.* **2000**, *8*, 343–352.
- [15] (a) Pandey, G.; Kapur, M. A general strategy towards the synthesis of 1-N-iminosugar type glycosidase inhibitors: demonstration by the synthesis of D- as well as L-glucose type iminosugars (isogomines). *Tetrahedron Lett.* **2000**, *41*, 8821–8824; (b) Pandey, G.; Kapur, M. A novel approach to both the enantiomers of potent glycosidase inhibitor Isogomine via PET-promoted cyclization of 1-[benzyl(trimethylsilyl)methylamino]-1,4,5-trideoxy-2,3-O-(1-methylethylidene)-threo-pent-4-ynitol. *Synthesis* **2001**, 1263–1267; (c) Pandey, G.; Kapur, M.; Khan, M.I.; Gaikwad, S.M. A new access to polyhydroxy piperidines of the azasugar class: synthesis and glycosidase inhibition studies. *Org. Biomol. Chem.* **2003**, *1*, 3321–3326.
- [16] Mehta, G.; Mohal, N. A norbornyl route to azasugars: stereoselective synthesis of isogomine analogues. *Tetrahedron Lett.* **2000**, *41*, 5747–5751.
- [17] Zhao, G.; Deo, U.C.; Ganem, B. Selective fowler reductions: asymmetric total syntheses of isogomine and other 1-azasugars from methyl nicotinate. *Org. Lett.* **2001**, *3*, 201–203.
- [18] Best, W.M.; MacDonald, J.M.; Skelton, B.W.; Steck, R.V.; Tilbrook, D.M.G.; White, A.H. The synthesis of a carbohydrate-like dihydrooxazine and tetrahydrooxazine as putative inhibitors of glycoside hydrolases: a direct synthesis of isogomine. *Can. J. Chem.* **2002**, *80*, 857–865.
- [19] Guanti, G.; Riva, R. Asymmetrized tris(hydroxymethyl)methane as precursor of iminosugars: application to the synthesis of Isogomine. *Tetrahedron Lett.* **2003**, *44*, 357–360.
- [20] Ouchi, H.; Mihara, Y.; Watanabe, H.; Takahata, H. A short and concise synthesis of Isogomine, Homoisogomine, and 5'-Deoxyisogomine. *Tetrahedron Lett.* **2004**, *45*, 7053–7056.
- [21] Altenbach, H.-J.; Blanda, G. A novel building block for the synthesis of isogomine analogues. *Tetrahedron: Asymm.* **1998**, *9*, 1519–1524.

- [22] Panfil, I.; Urbańczyk-Lipkowska, Z.; Chmielewski, M. Synthesis of iminosugars from α,β -unsaturated lactones and N-benzyl nitronone. *Polish. J. Chem.* **2005**, *79*, 239–249.
- [23] Panfil, I.; Bełzecki, C.; Urbańczyk-Lipkowska, Z.; Chmielewski, M. 1,3-Dipolar cycloaddition of nitrones to sugar enlactones. *Tetrahedron.* **1991**, *47*, 10087–10094.
- [24] Paśniczek, K.; Socha, D.; Jurczak, M.; Frelek, J.; Suszczyńska, A.; Urbańczyk-Lipkowska, Z.; Chmielewski, M. Double asymmetric induction in 1,3-dipolar cycloaddition of nitrones to 2,3-unsaturated sugar 1,5-lactones. *J. Carbohydr. Chem.* **2003**, *22*, 613–629.
- [25] Stecko, S.; Paśniczek, K.; Jurczak, M.; Urbańczyk-Lipkowska, Z.; Chmielewski, M. Double asymmetric induction in 1,3-dipolar cycloaddition of five-membered cyclic nitrones to 2-(5H)-furanones. *Tetrahedron: Asymm.* **2006**, *17*, 68–78.
- [26] Niva, T.; Inouye, S.; Tsuruoka, T.; Koaze, Y.; Niida, T. Nojirimycin as a potent inhibitor of glucosidase. *Agr. Biol. Chem.* **1970**, *34*, 966–968.
- [27] Tsuruoka, T.; Fukuyasu, H.; Ishii, M.; Usui, T.; Shibahara, S.; Inouye, S. Inhibition of mouse tumor metastasis with Nojirimycin-related compounds. *J. Antibiot.* **1996**, *49*, 155–161.
- [28] Legler, G. Glycoside hydrolases: mechanistic information from studies with reversible and irreversible inhibitors. *Adv. Carbohydr. Chem. Biochem.* **1990**, *48*, 319–384.
- [29] Faber, Ch.N. Glew, R.H. α -D-mannosidase. In *Meth. Enzym. Anal. 3rd Ed.*; Bergmeyer, H.U., Eds.; Verlag Chemie: Weinheim, **1984**; Vol. 4, 230–240.
- [30] Li, Y.-T. Studies on the glycosidases in jack bean meal. I. Isolation and properties of α -mannosidase. *J. Biol. Chem.* **1967**, *242*, 5474–5480.
- [31] Saul, R.; Chambers, J.P.; Molyneux, R.J.; Elbein, A.D. Castanospermine, a tetrahydroxylated alkaloid that inhibits β -glucosidase and β -glucocerebrosidase. *Archiv. Biochem. Biophys.* **1983**, *221*, 593–597.