This article was downloaded by: [Case Western Reserve University] On: 31 October 2014, At: 12:49 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry Publication details, including instructions for authors and

subscription information: http://www.tandfonline.com/loi/lcar20

Synthesis of Oligosaccharide Fragments of the Lipoarabinomannan from Rhodococcus ruber

Gladys C. Completo^a, Jessica A. Ponto^a & Todd L. Lowary^a ^a Alberta Ingenuity Centre for Carbohydrate Science and Department of Chemistry, The University of Alberta, Gunning-Lemieux Chemistry Centre, Edmonton, AB, Canada Published online: 16 Aug 2006.

To cite this article: Gladys C. Completo , Jessica A. Ponto & Todd L. Lowary (2005) Synthesis of Oligosaccharide Fragments of the Lipoarabinomannan from Rhodococcus ruber , Journal of Carbohydrate Chemistry, 24:4-6, 517-527

To link to this article: http://dx.doi.org/10.1081/CAR-200067114

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Synthesis of Oligosaccharide Fragments of the Lipoarabinomannan from *Rhodococcus ruber*

Gladys C. Completo, Jessica A. Ponto, and Todd L. Lowary

Alberta Ingenuity Centre for Carbohydrate Science and Department of Chemistry, The University of Alberta, Gunning-Lemieux Chemistry Centre, Edmonton, AB, Canada

The synthesis of two oligosaccharide fragments (1 and 2) of the lipoarabinomannan from *Rhodococcus ruber* is reported. Thioglycoside donors were used to assemble these glycans, which were prepared as their 8-(methoxycarbonyl)octyl glycosides for potential incorporation into neoglycoconjugates.

Keywords Arabinofuranosides, Glycosylation, Lipoarabinomannan, Thioglycosides

INTRODUCTION

A major immunomodulatory molecule in mycobacteria, including the human pathogen *Mycobacterium tuberculosis*, is lipoarabinomannan (LAM), a cell wall polysaccharide composed of mannopyranose and arabinofuranose residues bound to a phosphatidylinositol moiety.^[1,2] Structural differences in LAM molecules are observed among different mycobacterial species,^[3-6] and an increasing number of recent papers have reported the structures of LAM molecules from other actinomycetes.^[7-14] Among these^[9] is the LAM from *Rhodococcus ruber* (RruLAM), a species closely related to the opportunistic human pathogen *Rhodococcus equi*.

The structure of RruLAM is similar to that of mycobacterial LAM in that the mannan domain consists of an α - $(1 \rightarrow 6)$ -linked chain of mannopyranose residues. However, unlike mycobacterial LAM, which has a large arabinan

Dedicated to the memory of Jacques H. van Boom.

Received February 7, 2005; accepted March 9, 2005.

Address correspondence to Todd L. Lowary, Alberta Ingenuity Centre for Carbohydrate Science and Department of Chemistry, The University of Alberta, Gunning-Lemieux Chemistry Centre, Edmonton, AB, T6G 2G2 Canada. E-mail: tlowary@ualberta.ca

518 G. C. Completo, J. A. Ponto, and T. L. Lowary

motif attached to this mannan backbone, RruLAM has no separate arabinan domain per se. Instead, the arabinofuranose residues are found as single "capping" units attached α - $(1 \rightarrow 2)$ to the mannan backbone. Approximately 45% of the residues in the mannan core are capped at O-2, and the majority of these capping motifs are α -arabinofuranose residues, although in a small percentage of cases the α -arabinofuranose cap is replaced with an α -mannopyranose moiety. The proposed structure of RruLAM is shown in Figure 1.

We have a long-standing interest in the synthesis of oligosaccharide fragments of cell wall polysaccharides from mycobacteria and related organisms.^[15–19] As part of this program, we were interested in the synthesis of RruLAM fragments, functionalized to allow future preparation of neoglycoconjugates. Reported here is the preparation of disaccharide **1** and trisaccharide **2** (Fig. 2), as their 8-(methoxycarbonyl)octyl glycosides. Another similarly functionalized fragment of RruLAM, disaccharide **3**, has previously been synthesized.^[20]

RESULTS AND DISCUSSION

The synthesis of **1** and **2** relied on the use of the previously reported protected 8-methoxycarbonyloctyl glycosides $4^{[20]}$ and $5^{[21]}$ and the known thioglycosides $6^{[22]}$ and $7^{[23]}$. With these building blocks in hand, the preparation of the two targets was straightforward. The synthesis of **1** (Sch. 1) involved first the



Figure 1: Proposed structure of the lipoarabinomannan from *Rhodococcus ruber* (RruLAM). Approximately 45% of the α -(1 \rightarrow 6)-linked D-Manp residues are capped. The values for A, B, and C are approximately ≤ 1 , 15, and 11, respectively.



Figure 2: Synthetic targets 1 and 2, related structure 3, and monosaccharide building blocks (4-7) required for synthesis of 1 and 2.

coupling of alcohol **4**, with thioglycoside **7** promoted by *N*-iodosuccinimide and silver triflate.^[24] The expected disaccharide, **8**, was obtained in 75% yield. The anomeric stereochemistry of the arabinofuranosyl residue could be clearly established by NMR spectroscopy. In the ¹H NMR spectrum of **8**, the resonance for the anomeric hydrogen appeared as a singlet, which is consistent with the α -arabinofuranose stereochemistry.^[25] Had the β -isomer been formed, this signal would have appeared as a doublet with a 4–5 Hz coupling constant. Similarly, in the ¹³C NMR spectrum of **8**, the chemical shift of the anomeric carbon was 106.8 ppm as would be expected for an α -arabinofuranoside. This disaccharide was deprotected in two steps. Treatment of **8** with sodium methoxide removed the benzoyl groups and gave **9** in 95% yield. Subsequent hydrogenation of the benzoyl ethers afforded a 92% yield of the final target, **1**.

The preparation of 2 followed a similar route, as illustrated in Scheme 2. Reaction of alcohol **5**, with thioglycoside **6**, in the presence of *N*-iodosuccinimide



Scheme 1: (a) NIS, AgOTf, CH₂Cl₂, 0°C, 75%; (b) NaOCH₃, CH₃OH, rt, 95%; (c) H₂, Pd/C, CH₃OH, rt, 92%.



Scheme 2: (a) NIS, AgOTf, CH_2CI_2 , 0°C, 65%; (b) AcCl, CH_3OH , 0°C \rightarrow rt, 60%; (c) 7, NIS, AgOTf, CH_2CI_2 , 0°C, 68%; (d) NaOCH₃, CH_3OH , rt, 85%; (e) H_2 , Pd(OH)₂, CH_3OH , rt, 82%.

and silver triflate afforded the product disaccharide 10, in 65% yield. The acetyl group was then selectively cleaved in the presence of the benzoate esters upon treatment of 10 with methanolic HCl.^[26] The product, disaccharide alcohol 11, was obtained in 60% yield. The arabinofuranose residue was introduced by way of thioglycoside 7 under conditions identical to those used for the other glycosylations and the protected trisaccharide 12 was obtained in 68% yield. The *a*-stereochemistry of the arabinofuranose moiety could be established from the chemical shift of the anomeric carbon of this residue, which was 106.6 ppm.^[25] Deprotection of the product proceeded without incident. Compound 12 was first treated with sodium methoxide, which provided, in 85% yield, a product, 13, in which all of the acyl groups had been cleaved. Trisaccharide 2 was obtained in 82% yield by hydrogenation of the benzyl ethers in 13.

In summary, we have completed the synthesis of two fragments of LAM from *Rhodoccus ruber*. LAM fragments of known structure are expected to be useful tools in elucidating the biologic role of these glycans. The ester functionality present in the aglycone of these oligosaccharides will facilitate the preparation of neoglycoconjugates for use in, for example, assays requiring microtiter plates.

EXPERIMENTAL

Solvents were distilled from the appropriate drying agents before use. Unless stated otherwise, all reactions were carried out at rt and under a positive pressure of argon and were monitored by TLC on silica gel 60 F_{254} (0.25 mm, E. Merck). Spots were detected under UV light by charring with 10% H₂SO₄ in ethanol or by charring with anisaldehyde in ethanol. During the work-up of reaction mixtures, crude products in organic solvents were washed with

equal volumes of aqueous solutions. Solvents were evaporated under reduced pressure and below 40°C (bath). Column chromatography was performed on silica gel 60 (40–60 μ M) or Iatrobeads, which refers to a beaded silica gel 6RS–8060 manufactured by Iatron Laboratories (Tokyo). The ratio between silica gel and crude product ranged from 100 to 50:1 (w/w). Optical rotations were measured at 21 ± 2°C and are in units of degrees · mL/g · dm. ¹H NMR spectra were recorded at 400, 500, or 600 MHz, and chemical shifts are referenced to TMS (0.0 ppm, CDCl₃), CH₃OH (4.78 ppm, CD₃OD), or HOD (4.78 ppm, D₂O). ¹³C NMR spectra were recorded at 100 MHz, and ¹³C chemical shifts are referenced to CDCl₃ (77.00 ppm, CDCl₃), CD₃OD (49.15 ppm, CD₃OD), or external dioxane (68.11 ppm, D₂O). Electrospray mass spectra were recorded on samples suspended in THF or CH₃OH with added NaCl.

8-(Methoxycarbonyl)octyl 2-O-(2,3,5-tri-O-benzoyl- α -D-arabinofuranosyl)-3,4,6-tri-O-benzyl- α -D-mannopyranoside (8). Alcohol 4 (200 mg, 0.322 mmol), thioglycoside 7 (237 mg, 0.418 mmol), and powdered 4A molecular sieves (0.50 g) were dried overnight under vacuum with P_2O_5 . Freshly distilled CH_2Cl_2 (5 mL) was added and the mixture was cooled to 0°C before N-iodosuccinimide (108 mg, 0.483 mmol) and AgOTf (17 mg, 0.066) were added. The reaction mixture was stirred for 30 min and then neutralized with triethylamine, before being filtered through Celite and concentrated. The resulting residue was purified by column chromatography (6:1, hexanes: EtOAc) to yield 8 (257 mg, 75%) as a clear oil. R_f 0.44 (3:1, hexanes: EtOAc); $[\alpha]_{\rm D}$ + 13.1 (*c* 0.9, CHCl₃); ¹H NMR (500 MHz, CDCl₃): $\delta_{\rm H}$, 8.10–8.00 (m, 6 H, aromatic H), 7.70-7.48 (m, 3 H, aromatic H), 7.44-7.15 (m, 23 H, aromatic H), 5.73 (d, 1 H, J = 1.2 Hz, H-2'), 5.60 (d, 1 H, J = 1.2, 4.5 Hz, H-3'), 5.58 (s, 1 H, H-1'), 4.99 (d, 1 H, J = 1.8 Hz, H-1), 4.87 (d, 1 H, J = 10.8 Hz, $PhCH_2$), 4.84 (dd, 1 H, J = 5.1, 11.6 Hz, H-5a'), 4.76 (d, 1 H, J = 12.0 Hz, $PhCH_2$), 4.73 (d, 1 H, J = 12.0 Hz, $PhCH_2$), 4.70–4.68 (m, 2 H, H-4', H-5b'), 4.58-4.53 (m, 3 H, PhCH₂), 4.16-4.12 (s, 1 H, H-2'), 3.98 (dd, 1 H, J = 2.9, 9.3 Hz, H-3), 3.95 (dd, 1 H, J = 9.6, 9.6 Hz, H-4), 3.84–3.81 (ddd, 1 H, J = 1.7, 5.6, 9.6 Hz, H-5), 3.73 (dd, 1 H, J = 1.8, 10.8 Hz, H-6a), 3.71-3.68(m, 2 H, H-6b, aglycone OCH₂), 3.67 (s, 3 H, OCH₃), 3.39 (dt, 1 H, J = 6.6, 9.5 Hz, aglycone OCH₂), 2.31 (t, 2 H, J = 7.5 Hz, CH₂C=O), 1.65-1.55 (m, 4 H, aglycone CH₂), 1.35–1.31 (m, 8 H, aglycone CH₂); $^{13}\mathrm{C}$ NMR (125 MHz, $CDCl_3$): δ_C , 175.1, 166.2, 165.8, 165.2, 138.5, 138.4, 133.4, 133.0, 130.1, 129.9, 129.8, 129.8, 129.3, 129.3, 128.5, 128.4 (2), 128.3 (3), 128.2 (2), 128.1, 127.6 (2), 127.5, (2), 106.8, 99.1, 82.0, 81.2, 80.0, 77.7, 75.2, 75.1, 74.4, 73.2, 72.2, 75.1, 74.4, 73.2, 72.2, 75.1, 74.4, 73.2, 72.2, 75.1, 74.4, 73.2, 72.2, 75.1, 74.4, 73.2, 72.2, 75.1, 74.4, 73.2, 75.2, 75.1, 74.4, 73.2, 75.2, 75.1, 74.4, 73.2, 75.2, 75.1, 74.4, 75.2, 75.1, 74.4, 75.2, 75.1, 74.4, 75.2, 75.1, 75.2, 75.1, 75.1, 75.2, 75.1, 75.2, 75.1, 75.2, 75.1, 75.2, 75.1, 75.2, 75.1, 75.2, 75.1, 75.2, 75.2, 75.1, 75.2, 75.2, 75.1, 75.2, 75.2, 75.1, 75.271.8, 69.5, 67.8, 63.7, 51.4, 34.1, 29.5, 29.3, 29.2, 29.1, 26.1, 24.9. HRMS (ESI) calcd. for $(M + Na) C_{63}H_{68}O_{15}$: 1087.4450; found 1087.4448.

8-(Methoxycarbonyl)octyl 2-O-(D-arabinofuranosyl)-3,4,6-tri-Obenzyl-α-D-mannopyranoside (9). Disaccharide 8 (219 mg, 0.21 mmol) was

522 G. C. Completo, J. A. Ponto, and T. L. Lowary

dissolved in CH_3OH (10 mL) and the solution was stirred for 10 min before solid $NaOCH_3$ was added in small portions until the pH reached 9. The reaction mixture was stirred for 2 hr and then neutralized with acetic acid. The crude product was purified by column chromatography $(17:1, CH_2Cl_2:CH_3OH)$ to yield **9** as a colorless oil (150 mg, 95%). R_f 0.37 (17:1 CH₂Cl₂:CH₃OH); $[\alpha]_{\rm D}$ + 63.6 (c 0.6, CHCl₃); ¹H NMR (500 MHz, CDCl₃): $\delta_{\rm H}$, 7.37–7.25 (m, 15 H, aromatic H), 7.15–7.13 (m, 2 H, aromatic H), 5.14 (s, 1 H, H-1'), 4.90 (d, 1 H, J = 2.2 Hz, H-1), 4.80 (d, 1 H, J = 10.7 Hz, PhCH₂), 4.72 (d, 1 H, J = 10.7 Hz, PhCH $11.7 \text{ Hz}, \text{PhC}H_2$, 4.67 (d, 1 H, $J = 11.7 \text{ Hz}, \text{PhC}H_2$), 4.62 (d, 1 H, $J = 12.4 \text{ Hz}, \text{PhC}H_2$) PhCH₂), 4.48 (d, 1 H, J = 12.4 Hz, PhCH₂), 4.39 (d, 1 H, J = 10.7 Hz, PhCH₂), 4.22–4.21 (m, 1 H, H-4'), 4.19 (s, 1 H, H-2'), 4.01 (s, 1 H, H-3'), 3.93 (dd, 1 H, J = 2.8, 9.3 Hz, H-3), 3.89-3.80 (m, 4 H, H-5a', H-5b', H-5, H-2),3.69-3.61 (m, 7 H, aglycone OCH₂, H-4, C-6a, C-6b, OCH₃), 3.39 (dt, 1 H, $J = 6.6, 9.5 \text{ Hz}, \text{ aglycone OCH}_2), 2.31 (t, 2 H, J = 7.5 \text{ Hz}, CH_2C=O), 1.65-$ 1.55 (m, 4 H, aglycone CH_2), 1.35–1.31 (m, 8 H, aglycone CH_2); ¹³C NMR $(125 \text{ MHz}, \text{ CDCl}_3): \delta_C, 174.4, 138.3, 138.2, 138.1, 128.5, 128.4, 128.3, 128.0$ (2), 127.9, 127.6 (2), 109.4, 98. 8, 87.6, 79.2, 78.8, 78.2, 75.2, 74.8, 74.5, 73.3, 72.4, 71.4, 68.5, 67.9, 62.1, 51.5, 34.1, 29.4, 29.3, 29.2, 29.1, 26.0, 24.9. HRMS (ESI) calcd. for $(M + Na) C_{42}H_{56}O_{12}$: 775.3664; found 775.3664.

8-(Methoxycarbonyl)octyl 2-O-(α-D-arabinofuranosyl)-α-D-mannopyranoside (1). Disaccharide 9 (100 mg, 0.13 mmol) was dissolved in CH_3OH (5 mL) and 10% Pd/C (20 mg) was added. The solution was stirred under a H_2 atmosphere for 8 hr and the reaction mixture was then filtered through Celite and concentrated. The crude product was purified by column chromatography (10:1, $CH_2Cl_2:CH_3OH$) to yield **10** as a colorless oil (58 mg, 92%). R_f 0.30 (9:1, CH₂Cl₂:CH₃OH); $[\alpha]_{\rm D}$ + 56.0 (c 0.4, CH₃OH); ¹H NMR $(500 \text{ MHz}, \text{CD}_3\text{OD}): \delta_{\text{H}}, 5.04 \text{ (d, 1 H, } J = 1.8 \text{ Hz}, \text{H-1'}), 4.91 \text{ (s, 1 H, H-1)}, 4.06$ (dd, 1 H, J = 1.8, 5.9 Hz, H-2'), 3.98-3.94 (m, 1 H, H-3), 3.82-3.57 (m, 11 H, H-3)H-2', H3', H-4', H-4, H5, H-6a, H-6b, aglycone OCH₂, OCH₃), 3.51–3.48 (m, 2 H, H-5a', H-5b'), 3.39 (dt, 1 H, J = 6.6, 9.5 Hz, aglycone OCH₂), 2.30 (t, 2 H, J = 7.4 Hz, $CH_2C=0$, 1.61–1.56 (m, 4 H, aglycone CH_2), 1.39–1.31 (m, 8 H, aglycone CH₂); ¹³C NMR (125 MHz, CD₃OD): $\delta_{\rm C}$, 176.1 (C=O), 111.4 (C-1'), 100.7 (C-1), 85.5 (C-4'), 83.2 (C-2'), 79.4 (C-3'), 78.4 (C-2), 74.6 (C-4), 72.3 (C-3), 69.0 (C-5), 68.6 (aglycone OCH₂), 63.0 (C-5'), 62.9 (C-6), 52.0 (OCH₃), 34.8 (aglycone CH₂), 30.6 (aglycone CH₂), 30.4 (aglycone CH₂), 30.3 (aglycone CH_2), 30.1 (aglycone CH_2), 27.3 (aglycone CH_2), 26.0 (aglycone CH_2). HRMS (ESI) calcd. for $(M + Na) C_{21}H_{38}O_{12}$: 505.2256; found 505.2256.

8-(Methoxycarbonyl)octyl 6-O-(2-O-acetyl-3,4,6-tri-O-benzyl- α -D-man nopyranosyl)-2,3,4-tri-O-benzoyl- α -D-mannopyranoside (10). Alcohol 5 (680 mg, 1.03 mmol) was glycosylated with thioglycoside 6 (459 mg, 0.86 mmol) in CH₂Cl₂ (30 mL) using N-iodosuccinimide (289 mg, 1.28 mmol) and silver triflate (5 mg, 0.21 mmol) in the presence of powdered 4Å molecular

sieves (1.2 g) as described for the preparation of **8**. The product was purified by column chromatography (4:1, hexanes: EtOAc) to yield 10 (636 mg, 65%) as an oil. R_f 0.33 (3:1, hexanes: EtOAc); $[\alpha]_D$ -26.9 (c 1.0, CHCl₃); ¹H NMR $(500 \text{ MHz}, \text{ CDCl}_3)$: δ_H , 8.11–7.82 (m, 6 H, aromatic H), 7.61–7.14 (m, 24 H, aromatic H), 5.93 (dd, 1 H, J = 10.0, 10.0 Hz, H-4), 5.87 (dd, 1 H, J = 3.3, 10.2 Hz, H-3), 5.64 (dd, 1 H, J = 1.8, 3.3 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 3.9 Hz, H-2), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8, 5.38 (dd, 1 H, J = 1.8), 5.38 (dd, 1 H, J = 1.8 3.4 Hz, H-2', 5.04 (d, 1 H, J = 1.7 Hz, H-1), 4.89 (d, 1 H, J = 1.7 Hz, H-1'), 4.83 (d, 1 H, J = 10.9 Hz, PhCH₂), 4.57 (d, 1 H, J = 12.1 Hz, PhCH₂), 4.51 $(d, 1 H, J = 10.9 Hz, PhCH_2), 4.44 (d, 1 H, J = 10.9 Hz, PhCH_2), 4.37 (d, 1 H, J = 10.9 Hz, PhCH_2$ $J = 12.1 \text{ Hz}, \text{ Ph}CH_2$, 4.34 (d, 1 H, $J = 10.9 \text{ Hz}, \text{ Ph}CH_2$), 4.26–4.20 (m, 1 H, H-5), 3.98-3.92 (m, 2 H, H-5', H-3'), 3.86 (dd 1 H, J = 9.7, 9.7 Hz, H-4'), 3.78-3.72 (m, 2 H, aglycone OCH₂), 3.72-3.65 (m, 6 H, OCH₃, H-6a, H-6a', H-6b), 3.56-3.49 (m, 2 H, H-2, H-6b'), 2.31 (dd, 2 H, J = 7.5, 7.5 Hz, $CH_2C=0$, 2.12 (s, 3 H C(0) CH_3), 1.70–1.60 (m, 4 H, aglycone CH_2), 1.45– 1.30 (m, 8 H, aglycone CH₂); ¹³C NMR (125 MHz, CDCl₃): $\delta_{\rm C}$, 170.2, 166.2, $165.6,\ 165.5,\ 165.4,\ 138.6,\ 138.2,\ 138.0,\ 133.4,\ 133.3,\ 133.1,\ 129.9,\ 128.$ 129.7 (2), 129.5, 129.2, 128.4, 128.3 (2), 128.2, 127.8, 127.5, 127.3, 98.0, 97.5, 78.5, 75.0, 74.2, 73.3, 71.8, 71.5, 70.8, 70.2, 69.2, 68.6 (2), 68.5, 67.5, 66.8, 51.4, 34.1, 29.4, 29.3, 29.2, 29.1, 26.1, 25.0, 21.1. HRMS (ESI) calcd. for $(M + Na) C_{66}H_{72}O_{17}$: 1159.4667; found 1159.4681.

8-(Methoxycarbonyl)octyl 6-O-(3,4,6-tri-O-benzyl-α-D-mannopyra**nosyl**)-2,3,4-tri-O-benzoyl-α-D-mannopyranoside (11). A solution of methanolic HCl was prepared by dissolving acetyl chloride (1.5 mL) in CH_3OH (48.5 mL) at 0°C. The entirety of this solution was used to dissolve disaccharide **10** (614 mg, 0.54 mmol) and the reaction mixture was stirred for 6 hr. The solution was partially concentrated, diluted with CH_2Cl_2 , and washed successively with a saturated aqueous solution of NaHCO₃, water, and brine. The organic phase was dried with MgSO₄, filtered through cotton, and concentrated, and the resulting residue was purified by column chromatography (4:1, hexanes: EtOAc) to afford 11 (281 mg, 60%) as an oil. $R_f 0.25 (3:1, 1)$ hexanes: EtOAc); $[\alpha]_D = 15.1$ (c 1.0, CHCl₃); ¹H NMR (500 MHz, CDCl₃): δ_H , 8.10-7.83 (m, 6 H, aromatic H), 7.56-7.14 (m, 24 H, aromatic H), 5.97 (dd, 1 $J = 1.8, 3.3 \,\text{Hz}, \text{H-2}), 5.05 \,(\text{d}, 1 \,\text{H}, J = 1.7 \,\text{Hz}, \text{H-1}), 5.01 \,(\text{d}, 1 \,\text{H}, J = 1.5 \,\text{Hz})$ H-1'), 4.80 (d, 1 H, J = 10.9 Hz, PhCH₂), 4.54 (d, 1 H, J = 12.2 Hz, PhCH₂), 4.52 (d, 1 H, J = 10.9 Hz, PhCH₂), 4.49 (d, 1 H, J = 10.9 Hz, PhCH₂), 4.54 (d, 1 H, J = 12.2 Hz PhCH₂), 4.39 (d, 1 H, J = 10.9 Hz, PhCH₂), 4.23-4.20 (m, 2 H, H-5, H-5'), 4.02 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, J = 1.2, 2.6 Hz, H-2'), 3.94 (dd, 1 H, H-2'), 3.94 ($J = 4.6, 11.2 \text{ Hz}, \text{H-6a}, 3.85 - 3.70 \text{ (m, 5 H, aglycone OCH}_2, \text{H-3'}, \text{H-4'}, \text{H-6a'}),$ 3.66 (s, 3 H, OCH₃), 3.62 (dd, 1 H, J = 4.4, 10.8 Hz, H-6b'), 3.50-3.53 (m, 2 H, H-6b, H-6a), 2.32 (dd, 2 H, J = 7.5, 7.5 Hz, $CH_2C=0$), 1.59–1.72 (m, 4 H, aglycone CH_2), 1.30–1.42 (m, 8 H, aglycone CH_2); ¹³C NMR (125 MHz,

524 G. C. Completo, J. A. Ponto, and T. L. Lowary

 $\begin{array}{l} {\rm CDCl_3)\colon \delta_C,\ 170.3,\ 165.6,\ 165.5,\ 165.4,\ 138.5,\ 138.2,\ 137.9,\ 133.4,\ 133.3,\ 133.1, \\ 129.9,\ 129.8,\ 129.8,\ 129.5,\ 129.3,\ 129.2,\ 128.6,\ 128.5,\ 128.4,\ 128.3,\ 127.9, \\ 127.8,\ 127.7,\ 127.6,\ 127.5,\ 99.4,\ 97.6,\ 80.3,\ 75.0,\ 74.1,\ 73.3,\ 71.8,\ 71.2,\ 70.8, \\ 70.3,\ 69.4,\ 68.7,\ 68.6,\ 68.1,\ 67.6,\ 66.4,\ 51.4,\ 34.1,\ 29.4,\ 29.3,\ 29.2,\ 29.1,\ 26.1, \\ 25.0,\ HRMS\ (ESI)\ calcd.\ for\ (M+Na)\ C_{64}H_{70}O_{16}\colon 1117.4561;\ found\ 1117.4576. \end{array}$

8-(Methoxycarbonyl)octyl 6-O-[2-O-(2,3,5-tri-O-benzoyl-α-D-arabinofuranosyl)-3,4,6-tri-O-benzyl-α-D-mannopyranosyl]-2,3,4-tri-O-benzoyl- α -D-mannopyranoside (12). Alcohol 11 (261 mg, 0.24 mmol) was glycosylated with thioglycoside 7 (163 mg, 0.29 mmol) in CH₂Cl₂ (20 mL) using N-iodosuccinimide (80 mg, 0.36 mmol) and silver triflate (2 mg, 0.06 mmol) in the presence of powdered 4A molecular sieves (500 mg) as described for the preparation of 8. The product was purified by column chromatography (4:1, hexanes: EtOAc) to yield 12 (250 mg, 68%) as an oil. R_f 0.46 $(2:1, hexanes:EtOAc); [\alpha]_D - 19.8 (c 0.6, CHCl_3); _1H NMR (500 MHz, c) = 0.6 (c) + 0.6 (c) +$ $CDCl_3$: δ_H , 8.12–7.81 (m, 12 H, aromatic H), 7.62–7.11 (m, 33 H, aromatic H), 5.98 (dd, 1 H, J = 10.1, 10.1 Hz, H-4), 5.87 (dd, 1 H, J = 3.3, 10.1 Hz, H-3), 5.66–5.62 (m, 2 H, H-2", H-2), 5.56 (dd, 1 H, J = 1.0, 3.4 Hz, H-3"), 5.43 (s, 1 H, H-1), 5.05 (s, 2 H, H-1', H-1"), 4.80 (d, 1 H, J = 11.0 Hz, PhCH₂), 4.80-4.75 (m, 1 H, PhCH₂), 4.64-4.60 (m, 2 H, PhCH₂), 4.52 (d, 1 H, J =11.5 Hz, $PhCH_2$), 4.50 (d, 1 H, J = 11.5 Hz, $PhCH_2$), 4.47 (d, 1 H, J = 11.0 Hz, PhCH₂), 4.43 (d, 1 H, J = 12.0 Hz, PhCH₂), 4.22 (d, 1 H, J = 12.0 Hz, PhCH₂), 4.24–4.20 (m, 1 H, H-5), 4.10–4.13 (m, 1 H, H-5'), 4.00–3.88 (m, 5 H, H-4", H-3', H-4', H-2', H-6b'), 3.80-3.66 (m, 3 H, aglycone OCH₂, H-5a", H'5b"), 3.64 (s, 3 H, OCH₃), 3.58-3.48 (m, 4 H, H-6a, H-6a', H-6b, aglycone OCH₂), 2.31 (dd, 2 H, J = 7.5, 7.5 Hz, CH₂C=O), 1.60–1.70 (m, 4 H, aglycone CH₂), 1.20–1.44 (m, 8 H aglycone CH₂); ¹³C NMR (125 MHz, CDCl₃): $\delta_{\rm C}$, 170.2, 166.2, 165.7, 165.6, 165.5, 165.3, 165.1, 138.6, 138.4, 138.3, 133.4, 133.3, 133.2, 133.1, 132.9, 130.0, 130.0, 129.9 (2), 129.8 (2), 129.8, 129.7, 129.5, 129.3, 129.2, 128.6, 128.5, 128.4 (2), 128.3 (2), 128.2 (2), 128.0, 127.6, 127.4 (3), 106.6, 99.3, 97.5, 82.0, 81.2, 80.1, 77.6, 75.0, 74.8, 73.7, 73.0, 72.0, 71.9, 70.9, 69.4, 69.2, 68.6, 67.6, 66.5, 66.5, 63.6, 51.1, 34.1, 29.4, 29.3, 29.2, 29.1, 26.1, 25.0. HRMS (ESI) calcd. for $(M + Na) C_{90}H_{90}O_{23}$; 1561.5770, found 1561.5774.

8-(Methoxycarbonyl)octyl 6-O-[2-O-(α -D-arabinofuranosyl)-3,4,6tri-O-benzyl- α -D-mannopyranosyl]- α -D-mannopyranoside (13). Trisaccharide 12 (249 mg, 0.16 mmol) was deacylated in CH₃OH (30 mL) with sodium methoxide as described for the preparation of 9. The product was purified by column chromatography (2:1, EtOAc:hexanes) to yield 13 (125 mg, 85%) as an oil. R_f 0.20 (2:1, EtOAc:hexanes). The product was characterized by ¹H NMR spectroscopy and mass spectrometry and used in the subsequent reaction. ¹H NMR (400 MHz, CD₃OD): $\delta_{\rm H}$, 7.36–7.04 (m, 15 H, aromatic H), 5.19 (s, 1 H, H-1), 5.09 (s, 1 H, H-1'), 4.72–4.58 (m, 5 H, H-1", $4 \times PhCH_2$), 4.44 (d, 1 H, J = 12.2 Hz, PhC H_2), 4.32 (d, 1 H, J = 10.9 Hz, PhC H_2), 4.18–4.10 (m, 3 H, H-3', H-5', H-4"), 3.98–3.50 (m, 18 H, H-2, H-3, H-4, H-6a, H-6b, H-2', H-4', H-6a', H-6b, aglycone OC H_2 , OC H_3 , H-2", H-3", H-5a", H-5b"), 3.28 (1 H, ddd, J = 7.5, 7.5, 9.6 Hz, H-5), 2.28 (dd, 2 H, J = 7.5, 7.5 Hz, CH_2C =O), 1.62–1.58 (m, 4 H, aglycone CH_2), 1.32–1.20 (m, 8 H, aglycone CH_2). HRMS (ESI) calcd. for (M+Na) C₄₈H₆₆O₁₇: 937.4192; found 937.4198.

8-(Methoxycarbonyl)octyl 6-O-[2-O-(α -D-arabinofuranosyl)- α -Dmannopyranosyl]- α -D-mannopyranoside (2). Trisaccharide 13 (96 mg, 0.11 mmol) was dissolved in CH₃OH (5 mL) and Pd(OH)₂ (60 mg) was added. The solution was stirred under H_2 for 4 hr and then filtered through Celite and concentrated. The product was purified on Iatrobeads (7:3, $CH_2Cl_2:CH_3OH$) to yield 2 (55 mg, 82%) as an oil. R_f 0.19 (4:1, $CH_2Cl_2: CH_3OH$; $[\alpha]_D + 29.5$ (c 1.0, CH_3OH); ₁H NMR (600 MHz, D_2O): δ_H , 5.23 (d, 1 H, $J = 1.8 \,\text{Hz}$, H-1"), 5.10 (d, 1 H, $J = 1.8 \,\text{Hz}$, H-1'), 4.92 (d, 1 H, J = 1.8 Hz, H-1), 4.26 (dd, 1 H, J = 1.8, 3.6 Hz, H-2"), 4.17 (ddd, 1 H, J = 3.6, $5.4, \; 9.0 \, Hz, \; H\text{-}4''), \; 4.05 - 3.76 \; (m, \; 19, \; H\text{-}2, \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6a, \; H\text{-}6b, \; H\text{-}2', \; H\text{-}3, \; H\text{-}4, \; H\text{-}5, \; H\text{-}6b, \; H\text{-}6b, \; H\text{-}6b, \; H\text{-}6b, \; H\text{-}10, \; H\text{-}10,$ H-3', H-4', H-6a', H-6b', H-3", H-5a", H-5b", OCH₂ aglycone, OCH₃), 3.58 (ddd, 1 H, J = 7.5, 7.5, 9.6 Hz, H-5'), 2.39 (dd, 2 H, J = 7.5, 7.5 Hz) $CH_2C=0$, 1.67–1.54 (m, 4 H, aglycone CH_2), 1.39–1.25 (m, 8 H, aglycone CH_2); ¹³C NMR (125 MHz, D_2O): δ_C , 178.8 (C=O), 110.3 (C-1"), 100.4 (C-1'), 99.6 (C-1), 84.5 (C-2"), 82.1 (C-4"), 78.4, 77.4, 73.5, 71.9, 71.7, 71.2, 71.0, 68.9, 67.8, 67.5, 66.9 (C-1), 62.0 (C-5"), 61.6 (C-1'), 52.9 (OCH₃), 34.6 (aglycone CH_2), 29.3 (aglycone CH_2), 29.2 (aglycone CH_2), 29.1 (aglycone CH_2), 29.0 (aglycone CH_2), 26.2 (aglycone CH_2), 25.2 (aglycone CH_2). HRMS (ESI) calcd. for $(M + Na) C_{27}H_{48}O_{17}$: 667.2784, found 667.2788.

ACKNOWLEDGMENTS

This work was supported by the Natural Sciences and Engineering Research Council of Canada, The Alberta Ingenuity Centre for Carbohydrate Science and The University of Alberta.

REFERENCES

- Nigou, J.; Gilleron, M.; Puzo, G. Lipoarabinomannans: from structure to biosynthesis. Biochimie 2003, 85, 153-166.
- [2] Briken, V.; Porcelli, S.A.; Besra, G.S.; Kremer, L. Mycobacterial lipoarabinomannan and related lipoglycans: from biogenesis to modulation of the immune response. Mol. Microbiol. 2004, 53, 391–403.
- [3] Guérardel, Y.; Maes, E.; Briken, V.; Chirat, F.; Leroy, Y.; Locht, C.; Strecker, G.; Kremer, L. Lipomannan and lipoarabinomannan from a clinical isolate of

Mycobacterium kansasii—novel structural features and apoptosis-inducing properties. J. Biol. Chem. **2003**, *278*, 36637–36651.

- [4] Guérardel, Y.; Maes, E.; Elass, E.; Leroy, Y.; Timmerman, P.; Besra, G.S.; Locht, C.; Strecker, G.; Kremer, L. Structural study of lipomannan and lipoarabinomannan from *Mycobacterium chelonae*—presence of unusual components with α -(1 \rightarrow 3)-mannopyranose side chains. J. Biol. Chem. **2002**, *277*, 30635–30648.
- [5] Torrelles, J.B.; Khoo, K.-H.; Sieling, P.A.; Modlin, R.L.; Zhang, N.; Marques, A.M.; Treumann, A.; Rithner, C.D.; Brennan, P.J.; Chatterjee, D. Truncated structural variants of lipoarabinomannan in *Mycobacterium leprae* and an ethambututolresistant strain of *Mycobacterium tuberculosis*. J. Biol. Chem. **2004**, 279, 41227-41239.
- [6] Khoo, K.-H.; Dell, A.; Morris, H.R.; Brennan, P.J.; Chatterjee, D. Inositol phosphate capping of the non-reducing termini from rapidly growing strains of *Mycobacterium*. J. Biol. Chem. **1995**, 270, 12380–12389.
- [7] Gibson, K.J.C.; Gilleron, M.; Constant, P.; Puzo, G.; Nigou, J.; Besra, G.S. Identification of a novel mannose-capped lipoarabinomannan from *Amycolatopsis sulphurea*. Biochem. J. **2003**, 372, 821–829.
- [8] Gibson, K.J.C.; Gilleron, M.; Constant, P.; Brando, T.; Puzo, G.; Besra, G.S.; Nigou, J. *Tsukamurella paurometabola* lipoglycan, a new lipoarabinomannan variant with proinflammatory activity. J. Biol. Chem. 2004, 279, 22973–22982.
- [9] Gibson, K.J.C.; Gilleron, M.; Constant, P.; Puzo, G.; Nigou, J.; Besra, G.S. Structural and functional features of *Rhodococcus ruber* lipoarabinomannan. Microbiology **2003**, 149, 1437–1445.
- [10] Garton, N.J.; Gilleron, M.; Brando, T.; Dan, H.-H.; Giguère, S.; Puzo, G.; Prescott, J.F.; Sutcliffe, I.C. A novel lipoarabinomannan from the equine pathogen *Rhodococcus equi*—structure and effect on macrophage cytokine production. J. Biol. Chem. **2002**, 277, 31722–31733.
- [11] Gilleron, M.; Garton, N.J.; Nigou, J.; Brando, T.; Puzo, G.; Sutcliffe, I.C. Characterization of a truncated lipoarabinomannan from the actinomycete *Turicella otitidis*. J. Bacteriol. 2005, 187, 854–861.
- [12] Sutcliffe, I.C. Characterisation of a lipomannan lipoglycan from the mycolic acid containing actinomycete *Dietzia maris*. Antonie Van Leeuwenhoek 2000, 78, 195–201.
- [13] Flaherty, C.; Sutcliffe, I.C. Identification of a lipoarabinomannan-like lipoglycan in Gordonia rubropertincta. Syst. Appl. Microbiol. 1999, 22, 530–533.
- [14] Flaherty, C.; Minnikin, D.E.; Sutcliffe, I.C. A chemotaxonomic study of the lipoglycans of *Rhodococcus rhodnii* N445 (NCIMB 11279). Zentralbl. Bakteriol. 1996 (285), 11–19.
- [15] Subramaniam, V.; Lowary, T.L. Synthesis of oligosaccharide fragments of mannosylated lipoarabinomannan from *Mycobacterium tuberculosis*. Tetrahedron 1999, 55, 5965–5976.
- [16] Han, J.; Gadikota, R.R.; McCarren, P.R.; Lowary, T.L. Synthesis of octyl arabinofuranosides as substrates for mycobacterial arabinosyltransferases. Carbohydr. Res. 2003, 338, 581–588.
- [17] Yin, H.; D'Souza, F.W.; Lowary, T.L. Arabinofuranosides from mycobacteria:synthesis of a highly branched hexasaccharide and related fragments containing β -arabinofuranosyl residues. J. Org. Chem. **2002**, *67*, 892–903.

- [18] D'Souza, F.W.; Ayers, J.D.; McCarren, P.R.; Lowary, T.L. Arabinofuranosyl oligosaccharides from mycobacteria: synthesis and effect of glycosylation on ring conformation and hydroxymethyl group rotamer populations. J. Am. Chem. Soc. 2000, 122, 1251-1260.
- [19] Gadikota, R.R.; Callam, C.S.; Appelmelk, B.J.; Lowary, T.L. Synthesis of oligosaccharide fragments of mannosylated lipoarabinomannan appropriately functionalized for neoglycoconjugate preparation. J. Carbohydr. Chem. 2003, 22, 459–480.
- [20] Tsui, D.S.; Gorin, P.A.J. Preparation of 8-(methoxycarbonyl)octyl glycosides of α-D-mannopyranose, 2-O-α-mannopyranosyl-α-D-mannopyranose, β-D-galactofuranose, and 3-O-β-D-galactofuranosyl-α-D-mannopyranoside. Carbohydr. Res. 1986, 156, 1–8.
- [21] Srivastava, O.P.; Hindsgaul, O. Synthesis of 6'-O-phosphorylated O- α -D-manno-pyranosyl (1 \rightarrow 3)- and (1 \rightarrow 6)- α -D-mannopyranosides. Carbohydr. Res. **1987**, 161, 324–329.
- [22] Callam, C.S.; Lowary, T.L. Synthesis and conformational investigations of methyl 4a-carba-D-arabinofuranosides. J. Org. Chem. 2001, 66, 8961–8972.
- [23] Callam, C.S.; Gadikota, R.R.; Lowary, T.L. Sensitivity of ${}^{1}J_{C1-H1}$ magnitudes to anomeric stereochemistry in 2,3-anhydro-O-furanosides. J. Org. Chem. **2001**, 66, 4549–4558.
- [24] Konradsson, P.; Udodong, U.E.; Fraser-Reid, B. Iodonium-promoted reactions of disarmed thioglycosides. Tetrahedron Lett. 1990, 31, 4313-4316.
- [25] Mizutani, K.; Kasai, R.; Nakamura, M.; Tanaka, O.; Matsuura, H. NMR spectral study of α and β -L-arabinofuranosides. Carbohydr. Res. **1989**, *185*, 27–38.
- [26] Byramova, N.E.; Ovchinnikov, M.V.; Backinowsky, L.V.; Kochetkov, N.K. Selective removal of O-acetyl groups in the presence of O-benzoyl groups by acid-catalysed methanolysis. Carbohydr. Res. 1983, 124, C8–C11.