


than by the usual carbonium ion structure



The rapid formation of the ether suggests that the latter formulation is of considerable importance.

Experimental

The spectra were run on a Beckman model DU spectrophotometer; 1-cm. thick quartz cells were used.

Crystal Violet Preparation.—The carbinol was prepared by dissolving the chloride salt of crystal violet (a product of the National Aniline Division, Allied Chemical and Dye) in water and precipitating the carbinol by addition of solid sodium hydroxide pellets. After filtration, the solid carbinol was blue in color. Attempts to purify the product by recrystallization from ligroin and benzene were futile. The method of purification was that followed by Heertjes, *et al.*,⁶ who also reported that simple recrystallization from organic solvents failed. The carbinol was dissolved in acetone and precipitated by the slow addition of an aqueous sodium hydroxide solution. The carbinol, collected by filtration, was much less colored than originally. Repetition of this process several times yielded a white solid. In order to remove traces of sodium hydroxide from the carbinol, it was dissolved in a small volume of anhydrous ether, and the solution filtered. The carbinol, after evaporation of the ether, was stored under vacuum since exposure to air causes the compound to turn blue slowly. The compound melted at 200° dec.; a melting point of 205° has been reported.⁶

Anal. Calcd. for $\text{C}_{25}\text{H}_{31}\text{ON}_3$: C, 77.08; H, 8.02. Found: C, 77.02; H, 8.11.

Methyl Ether Preparation.—A 3-g. sample of crystal violet was dissolved in 10 ml. of concd. sulfuric acid, giving an intense orange-brown solution. This solution was added dropwise to 50 ml. of absolute methanol which was cooled in a

(6) P. M. Heertjes, J. C. Van Kerkhof and K. A. De Vries, *Rec. trav. chim.*, **62**, 737 (1943).




Fig. 1.—Absorption spectra of triphenylcarbonium ions in H_2SO_4 : —, tri-*p*-dimethylaminophenylcarbinol; ·····, tri-*p*-nitrophenylcarbinol¹; - - -, triphenylcarbinol.¹ Crystal violet solution was $2.5 \times 10^{-3} M$. Spectrum for methyl ether of crystal violet was identical with that for crystal violet.

Dry Ice-acetone-bath during the addition. The resulting solution was immediately added to 200–300 ml. of an ice-water mixture and sufficient sodium hydroxide solution added to make the solution basic. The methyl ether was extracted from this solution with two 100-ml. portions of ether. Evaporation of this solution yielded a blue solid which was purified in the same manner as that described for the carbinol. The yield of crude material was approximately 50%. The purified ether melted with decomposition at 145–150°.

Anal. Calcd. for $\text{C}_{25}\text{H}_{33}\text{ON}_3$: C, 77.38; H, 8.24. Found: C, 77.16; H, 8.19.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF TENNESSEE
KNOXVILLE, TENNESSEE

COMMUNICATIONS TO THE EDITOR

TRACER STUDY OF THE OXYGEN IN PRECIPITATED HgO AND Ag_2O ; O^{18} ISOTOPE EFFECT IN THE REACTION OF OH^- WITH Hg^{++} AND Ag^+

Sir:

O^{18} experiments have been used to investigate the source of the oxygen in HgO and Ag_2O precipitated by OH^- . Further, an $\text{O}^{16}/\text{O}^{18}$ effect was found in the reactions of Hg^{++} and Ag^+ with OH^- ; O^{18} concentrates preferentially in the precipitated oxide.

Tracer experiments were as follows. One ml. of a solution of M NaOH , enriched in O^{18} (1.4 atom %) was added to 100 ml. of *ca.* M $\text{Hg}(\text{NO}_3)_2$ in normal water (0.20 atom % O^{18}); the precipitated HgO was washed, dried and pyrolyzed. The $\text{O}^{18}/\text{O}^{16}$ ratio of the liberated oxygen was 1% lower than the ratio for normal water. Thus the oxygen in the HgO is derived exclusively from the original aquo-cation, which apparently exchanges only slowly with solvent water. Proton transfer from hydrate water to OH^- is a satisfactory explanation. However, in similar experiments with

AgNO_3 (1–9 M), the Ag_2O was found to be substantially enriched in O^{18} , ranging from 0.67 atom % (no stirring, high $[\text{Ag}^+]$) to 0.26 atom % (vigorous stirring, low $[\text{Ag}^+]$).¹ A reason for the difference between Hg^{++} and Ag^+ may be the tighter binding of the coordinated water around the more polarizing Hg^{++} .²

The $\text{O}^{16}/\text{O}^{18}$ isotope effect was studied with normal O^{18} abundance materials. Oxygen from the pyrolysis of the oxides was compared mass spectrometrically with reference oxygen which differed by +2.3% in $\text{O}^{18}/\text{O}^{16}$ ratio from distilled water. Fractionation factors with respect to water were computed: $\alpha = (\text{O}^{16}/\text{O}^{18})_{\text{oxide}} / (\text{O}^{16}/\text{O}^{18})_{\text{water}}$.

When a solution of Hg^{++} was added to a solution containing excess OH^- , α was 1.018. For the

(1) Auxiliary experiments showed negligible exchange between water and HgO or Ag_2O in 1 hr.

(2) H. Taube, *et al.* (discussed in *J. Phys. Chem.*, **58**, 523 (1954)) studied exchange between hydrated cations and solvent water and found differences in lability for various (other) cations.

reverse procedure (OH^- into Hg^{++}), α was the same. When Ag^+ was added to excess OH^- , α was 1.016. For the reverse addition, α was significantly smaller; all values were less than 1.013. Thus, for a given sample to Ag_2O , it appears possible to deduce the order in which the reactants were originally mixed by measurement of the $\text{O}^{18}/\text{O}^{16}$ ratio.

These results are suggestive of a kinetic isotope effect. For the Hg^{++} aquo-ion, the preferential rupture of $\text{O}^{16}\text{-H}$ relative to $\text{O}^{18}\text{-H}$ would concentrate O^{16} in the HgO . For Ag^+ , the situation is more complex. An alternative mechanism for the fractionation involves the equilibrium isotopic exchange between OH^- and H_2O , favoring the concentration of O^{16} in the OH^- .³

Further work is in progress to extend the data and examine other cations. We appreciate the discussions with R. W. Parry, the assistance of W. E. Barrett with the mass spectrometer, and the financial support of the A.E.C. and the Michigan Memorial-Phoenix Project.

(3) The ionic hydration fractionation effect of Feder and Taube (*J. Chem. Phys.*, **20**, 1335 (1952)) wherein O^{18} concentrates in the aquo-ion relative to the solvent water would probably be of negligible importance since most of the experiments were carried out at low concentrations.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF MICHIGAN
ANN ARBOR, MICHIGAN


RICHARD B. BERNSTEIN
HARRY G. PARS

RECEIVED JUNE 14, 1955

ON THE STEREOCHEMISTRY AND MECHANISM OF THE DIELS-ALDER REACTION¹

Sir:

We wish to report the first conclusive evidence that isomerization of a bicyclic Diels-Alder adduct of *endo*-stereochemistry to the *exo*-isomer can occur without dissociation into kinetically free addends.² When the *endo* cyclopentadiene-maleic anhydride adduct (N), radioactively labeled with C^{14} in the carbonyl carbons, was heated with an equimolar quantity of non-radioactive maleic anhydride (MA) in boiling decalin (190°), rapid exchange of N with MA occurred simultaneously with slower conversion of N to the *exo*-isomer (X). (Exchange of X with MA was small at short reaction times.)




The variation of *endo*-activity with time was determined by re-isolating N from quenched reaction mixtures. Since isolation of X directly from reaction mixtures at small concentrations of X was not feasible, the activity and concentration

- (1) Sponsored by the Office of Ordnance Research, U. S. Army.
- (2) For previous discussion of the mechanism of such isomerizations see (a) R. B. Woodward and H. Baer, *THIS JOURNAL*, **66**, 645 (1944); (b) K. Alder and W. Trimborn, *Ann.*, **566**, 58 (1950); (c) D. Craig, *THIS JOURNAL*, **73**, 4889 (1951); (d) D. Craig, J. J. Shipman, J. Kiehl, F. Widmer, R. Fowler and A. Hawthorne, *ibid.*, **76**, 4573 (1954); (e) J. A. Berson and R. Swidler, *ibid.*, **75**, 1721 (1953).

of X were determined by an isotope dilution technique.

The activities theoretically to be expected of X arising exclusively by an "internal" path from N (path A) and by an "external" path (path B) were calculated by a graphical integration method.



The data for activity of N *vs.* time and for moles of X formed *vs.* time were plotted. A reaction period of t minutes was subdivided into arbitrarily small time increments. If X_1 is the number of moles of X formed during time increment 1, X_2 during time increment 2, etc., and X_t during time t , α_0 the original activity of N, α_1 the average activity of N during time increment 1, etc., then the activity of X to be expected by the "internal" path (A) is $\alpha_1 X_1/X_t + \alpha_2 X_2/X_t + \dots + \alpha_t X_t/X_t$. Since the only source of activity in the system is N,³ the activity of MA at any time is $\alpha_0 - \alpha_j$ where α_j is the average activity of N during any given time increment. The activity of X to be expected by the "external" path (B) is therefore $(\alpha_0 - \alpha_1)X_1/X_t + (\alpha_0 - \alpha_2)X_2/X_t + \dots + (\alpha_0 - \alpha_t)X_t/X_t$. Application of these calculations to the data show that after three minutes and ten minutes 31.4% and 45.4% respectively of the X present has been formed by path A.⁴

Although epimerization of N to X *via* enolic species formally satisfies the requirement that part of the isomerization occur by path A, we consider this an unlikely mechanism, since the isomerization does not appear to be catalyzed by bases and does not occur with the dihydro derivative of N.^{2c} The results also do not formally exclude a mechanism involving intermediate formation of a diradical followed by hydrogen transfer, as proposed by Craig,^{2c} or a diffusion-controlled secondary recombination of original partners.⁵ However, we consider the most probable detailed description of path A to be conversion of N to a complex in which MA and cyclopentadiene are relatively free to rotate with respect to each other, followed by recombination of the fragments in either stereochemically possible sense. On the basis of the present evidence, this is indistinguishable from primary recombination in a Franck-Rabinowitch solvent cage.

We are indebted to the Office of Ordnance Research for financial support and to Professors A. W. Adamson and S. W. Benson for helpful discussion.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES 7, CALIFORNIA
JEROME A. BERSON
ROSALIE D. REYNOLDS
RECEIVED JULY 11, 1955

(3) This is true if the exchange of X with MA is neglected. The correction for this is small. Its omission leads to a slightly more conservative estimate of the importance of the "internal" path (A).

(4) The ratio of the amounts of X formed by the two paths is $(k_3 N - k_3 X)/(k_4 M - k_4 X)$. This need not be time invariant if $k_3 \neq k_4$ or if C_6H_6 is being irreversibly removed from the system, *e.g.*, by polymerization.

(5) *Cf. inter alia*, R. M. Noyes, *J. Chem. Phys.*, **22**, 1349 (1954).